

Institute for Chemical Research Kyoto University

ICR ANNUAL REPORT 2016 (Volume 23) - ISSN 1342-0321 -

This Annual Report covers from 1 January to 31 December 2016

Editors:

Professor: TERANISHI, Toshiharu Professor: WATANABE, Hiroshi

Associate Professor: SAKAMOTO, Masanori

Professor: OGATA, Hiroyuki

Editorial Staff:

Public Relations Section: INOUE, Junko

TAKEHIRA, Tokiyo NAKAMURA, Kaori NAKANO, Yukako HAMAOKA, Meri

Published and distributed by:

Institute for Chemical Research (ICR), Kyoto University

Copyright © 2017 Institute for Chemical Research, Kyoto University Enquiries about copyright and reproduction should be addressed to: *ICR* Annual Report Committee, Institute for Chemical Research, Kyoto University

Note: *ICR* Annual Report available from the *ICR* Office, Institute for Chemical Research, Kyoto University, Gokasho, Uji-city, Kyoto 611-0011, Japan

Tel: +81-(0)774-38-3344 Fax: +81-(0)774-38-3014 E-mail koho@scl.kyoto-u.ac.jp

URL http://www.kuicr.kyoto-u.ac.jp/sites/icr/

Uji Library, Kyoto University Tel: +81-(0)774-38-3010 Fax: +81-(0)774-38-4370

E-mail ujibunkan@mail2.adm.kyoto-u.ac.jp

URL http://lib.kuicr.kyoto-u.ac.jp/homepage/japanese/index.htm


Printed by

Zenis Co., Ltd.

67-10 Karahashi-Akakanecho, Minami-Ku, Kyoto 601-8463, Japan

TEL: +81-(0)75-671-7100 FAX: +81-(0)75-671-7107

ICR ANNUAL REPORT 2016



Institute for Chemical Research Kyoto University

Volume 23



Preface

The Institute for Chemical Research (ICR) was launched in 1926 as the first research institute at Kyoto University, and celebrates its 90th Anniversary in 2016. The philosophy at the time of its foundation was to "Excel in the Investigation of Basic Principles of Chemistry and Their Applications," and studies on special medicinal substances, organometallic chemistry, incendiary reagents, and other topics were undertaken at nine research laboratories. Over its lifetime, the ICR has continuously produced outstanding research achievements. Today, the organization is in five research divisions: Synthetic Chemistry, Materials Chemistry, Biochemistry, Environmental Chemistry, and Multidisciplinary Chemistry and three research centers on Beam Science, Elements Science (IRCELS), and Bioinformatics. About 120 faculty members, 210 graduate students and 60 researchers are engaged in research activities in 30 laboratories in total supervised by full-time professors. Further, five laboratories are supervised by visiting professors.

The research within the ICR encompasses the fields of chemistry, physics, biology, and informatics. The chemical studies core covers fields including physical chemistry, inorganic chemistry, organic chemistry, materials chemistry, and biochemistry. The graduate schools encompassing the laboratories accept students in diverse fields of science, engineering, agriculture, pharmaceutical sciences, medicine, and informatics. These laboratories are spearheading leading-edge research, and yielding outstanding results in their own and related research areas with publications such as: 1) Synthesis of a Distinct Water Dimer inside Fullerene C₇₀, 2) Formation of Pseudomorphic Nanocages from Cu₂O Nanocrystals through Anion Exchange Reactions, 3) Tuning Magnetic Anisotropy by Interfacially Engineering the Oxygen Coordination Environment in a Transitionmetal Oxide, 4) Discovery of 12-mer Peptides That Bind to Wood Lignin, and 5) Experimental Observation of Snell's Law for Magnetostatic Spin Waves in Thin Ferromagnetic Permalloy Films, all achieved last year. The legacy of our founding philosophy continues to the present day and describes the essence of our research activities. The ICR has entrusted its members, with this vision in mind, to choose and pursue research topics at the forefront of advanced chemistry with bottom-up paradigms. Sustainable and sustained growth of the human race is a key issue of this century, and we must reform our country from various perspectives. Hence, the ICR encourages its members to be actively involved in research projects and to value the development of unique interdisciplinary research projects, in order to contribute to the future of our society from materials-related fields.

The ICR also collaborates with domestic and international universities and research organizations (with 70 official international collaboration agreements as of January 1, 2017) and functions as a Joint Usage/Research Center proclaiming the Frontier and Interdisciplinary Research Core for Deepening Investigation and Promoting Collaboration in Chemistry-oriented Fields supported by MEXT (2016-2021); its activity during the former term received high commendation on the term-end assessment in 2016. In addition, the ICR collaborates with the Research Institute for Sustainable Humanosphere and the Institute of Advanced Energy at Kyoto University to start another MEXT-supported joint research project on bio-inspired smart materials as of April 1, 2015. Furthermore, the ICR—IRCELS in particular—is making a significant contribution to the MEXT Project of Integrated Consortium on Chemical Synthesis (2016-2021), in collaboration with the Catalysis Research Center at Hokkaido University, the Research Center for Materials Science at Nagoya University, and the Institute for Materials Chemistry and Engineering at Kyushu University. We also fully strive to foster and secure young researchers through these activities. For instance, we provide an in-house annual grant system named "ICR Grant for Encouraging Promoting Integrated Research." The strong collaboration basis so far constructed in-house and with outside researchers ensures that the Institute will continue to serve as a core of global research propellers in chemistry-oriented fields. With this consideration, we would appreciate your continued encouragement and support.

January 2017

TOKITOH, Norihiro Director

ICR News 2016

ICR Celebrates 90th Anniversary



Chairperson of ICR 90th Anniversary Committee Prof KAWABATA, Takeo

Ever since it was formally founded on 4 October 1926 to conduct "fundamental and applied researches on specific subjects in chemistry", ICR has carried out extensive studies in an extremely wide range of basic and applied chemistry, reaching its 90th anniversary last fall. We celebrated the anniversary on November 11, 2016 by holding Commemorative Lecture Meeting, Ceremony, and Banquet at the Clock Tower Centennial Hall (Centennial Hall and International Conference Hall), Kyoto University. We had another commemorative event, Poster Exhibition showing current activities

of ICR. In the Lecture Meeting, lectures were delivered by Director N. Tokitoh on the past, present, and future perspective of ICR and by four ICR members, Professors H. Ogata, Y. Shimakawa, S. Sakabe, and S. Yamago on the frontier science. The Commemorative Ceremony included an address by Director N. Tokitoh and greetings by President J. Yamagiwa, Kyoto University, Director-General Y. Komatsu, Research Promotion Bureau, the Ministry of Education, Culture, Sports, Science and Technology, Dr. M. Sasaki (on behalf of Dr. Z. Matsuura), Council for Research Institutes and Centers of

> Japanese National Universities, and President. H. Yamamoto, the Chemical Society of Japan. The

> Lecture Meeting and Ceremony had an approximate total of 500 participants including some 200 guests from academia and industry. The Banquet was also a success with over 300 participants, getting started with greetings by President Y. Hiraki, Joint Usage/Research Center, Director N. Oshiman, Kyoto University Research Coordination Alliance, Dr. K. Tamao, RIKEN, and Dr. K. Soda, Professor Emeritus, Kyoto University, and proceeding in a

friendly and enjoyable atmosphere.

Lecture given by Director Tokitoh, ICR



Exhibition of the latest researches at ICR



Commemorative Ceremony for 90th Anniversary



Speech by President Yamagiwa, **Kvoto University**



Speech by Director-General Komatsu, Research Promotion Bureau, the Ministry of Education, Culture, Sports, Science and Technology



KAGAMIBIRAKI ceremony at the banquet

International Research Network Weeks

Prof FUTAKI, Shiroh

ICR has built a community across the world including researchers, students and faculty. A remarkable event "International Research Network Weeks" was held on March 7–11, 2016. ICR invited global researchers and students who have been collaborated with ICR members for a week. The aim of this event is to strengthen partnerships and enhance international collaboration with institutions having MOU (Memorandum of Understanding) with ICR. At the beginning of the event, ICR built new ones with Department of Chemistry and Bioengineering, Notre Dame University (U.S.) and Department of Chemistry and Bioengineering, Ohio State University (U.S.). It is the 68th ICR's faculty level MOU.

On March 7–8, the opening event -Institute for Chemical Research International Symposium 2016 (ICRIS'16) "Research Network Based on ICR MOU"- was held at

Kihada Hall of Uji-Obaku Plaza. 13 investigators from Asia, Europe, and the United State and their Japanese counterparts, 7 investigators having domestic collaboration with ICR members and 6 faculty members of ICR, and 78 posters presented at the symposium. We hope that all presentations stimulated discussion among members of ICR's scientific community.



Materials Related with "Origin of Polyethylene Industry by the High-pressure Process in Japan" in the Institute for Chemical Research Were Approved as Chemical Heritage


Prof MURATA, Yasujiro

The engineering drawing, notebook, and reports for the industrial process in Kyoto University and the medal made by the polyethylene celebrating the first actual production in Japan were certified. Polyethylene (PE) is the most common synthetic polymers in petrochemical industry. There are three kinds of PEs depending on the production methods and the density. One of them is the low-density PE produced by high-pressure conditions, and it was industrialized in 1939 in UK. The PE has superior properties for insulation of radiofrequency, and therefore it was an essential material for radar devices during the World War II. In Japan, three groups started research works of PEs.

Although one of the group started small industrial process at Minamata in 1945, all facility were destroyed by an air raid. After the World War II, continuous production was conducted in Kyoto University during 1951 to 1953, with which the materials are related.

Chemical Heritage is one of the most important historical materials related to chemistry and chemical technology in Japan approved by the Chemical Society of Japan. The presentation ceremony of the award was held in the 96th Spring Annual Meeting of the Chemical Society of Japan at Doshisha University, Kyotanabe Campus on 26th March 2016.







v

CONTENTS

ICR Nev	vs 2016
ORGAN	IZATION
TOPICS	AND INTRODUCTORY COLUMNS OF LABORATORIES
Divi	sion of Synthetic Chemistry
Orga	anoelement Chemistry
Stru	ctural Organic Chemistry
Synt	hetic Organic Chemistry
Adv	anced Inorganic Synthesis
Divi	sion of Materials Chemistry
Che	mistry of Polymer Materials
Poly	mer Controlled Synthesis
Inor	ganic Photonics Materials
Nan	ospintronics
Divi	sion of Biochemistry
Biof	functional Design-Chemistry
Che	mistry of Molecular Biocatalysts
Mol	ecular Biology
Che	mical Biology
Divi	sion of Environmental Chemistry
Mol	ecular Materials Chemistry
Hyd	rospheric Environment Analytical Chemistry
Solu	tion and Interface Chemistry
Mol	ecular Microbial Science
Divi	sion of Multidisciplinary Chemistry
Poly	mer Materials Science
Mol	ecular Rheology
Mol	ecular Aggregation Analysis
Adv	anced Research Center for Beam Science
Part	icle Beam Science
Lase	er Matter Interaction Science
Elec	tron Microscopy and Crystal Chemistry
Stru	ctural Molecular Biology

	International Research Center for Elements Science	
	Synthetic Organotransformation	50
	Advanced Solid State Chemistry	52
	Organometallic Chemistry	54
	Nanophotonics	56
	Bioinformatics Center	
	Chemical Life Science	58
	Mathematical Bioinformatics	60
	Bio-knowledge Engineering	62
HAK	KUBI RESEARCHERS' ACTIVITIES IN ICR	65
ACT	TIVITIES OF JOINT USAGE/RESEARCH CENTER	67
	JURC Cooperative Research Subjects 2016	68
	JURC Publications (Selected Examples)	73
VISI	ITING PROFESSORS' ACTIVITIES IN ICR	75
PER	SONAL	77
	Awards	78
	Paper Awards	82
	Poster Awards	84
PUB	ELICATIONS	88
INT	ERNATIONAL RESEARCH COLLABORATIONS.	102
SEL	ECTED GRANTS	103
THE	CSES	110
THE	2 116TH ICR ANNUAL SYMPOSIUM	114
SEM	IINARS	118
MEE	ETINGS AND SYMPOSIA	121
IND	EX	123
	NAME INDEX.	124
	KEYWORD INDEX	127

– ${f A}$ bbreviations used in the columns –

Prof Em	Professor Emeritus	Guest Scholar	Guest Scholar
Prof	Professor	Guest Res Assoc	Guest Research Associate
Vis Prof	Vis Prof Visiting Professor		Special Post-Doctoral
Assoc Prof Associate Professor			Research Fellow (JSPS)
Vis Assoc Prof	Visiting Associate Professor	PD	Post-Doctoral Research Fellow
Senior Lect	Senior Lecturer	Res	Researcher
Assist Prof	Assist Prof Assistant Professor		Research Support Staff
Lect (pt)	Lect (pt) Lecturer (part-time)		Project Researcher
Specially Appointed Prof		RF	Research Fellow
	Specially Appointed Professor	D 1~3	Doctoral Course (Program) 1~3
Program-Specific Assoc Prof		M1~2	Master's Course (Program) 1∼2
	Program-Specific Associate	RS	Research Student
	Professor	UG	Undergraduate Student
Program-Specific Senior Lect		D Sc	Doctor of Science
	Program-Specific Senior	D Eng	Doctor of Engineering
	Lecturer	D Agr	Doctor of Agricultural Science
Program-Specific Assist Prof		D Pharm Sc	Doctor of Pharmaceutical Science
Program-Specific Assistant :		D Med Sc	Doctor of Medical Science
	Professor	D Inf	Doctor of Informatics
Program-Specific Res		D Human & Envirnmtl. Studies	
	Program-Specific Researcher		Doctor of Human and
Techn Staff	Technical Staff		Environmental Studies
Assist Res Staff	Assistant Research Staff	Ph D	Doctor of Philosophy
Assist Techn Staff Assistant Technical Staff		(pt)	part-time

ORGANIZATION

Institute for Chemical Research

5 Divisions and 3 Centers

Laboratories



TOPICS AND INTRODUCTORY COLUMNS OF LABORATORIES

Division of Synthetic Chemistry - Organoelement Chemistry -

http://oec.kuicr.kyoto-u.ac.jp/~tokitohlab/index-e.html



TOKITOH, Norihiro (D Sc)



Assoc Prof (D Sc)



Assist Prof SASAMORI, Takahiro MIZUHATA, Yoshiyuki (DSc)



Techn Staff HIRANO, Toshiko



PD GUO, Jing-Dong (Ph D)



PD MAJHI, Paresh Kumar (Ph D)

Students

HIRANO, Koki (D3) SUZUKI, Yuko (D2) FUJIMORI, Shiori (D1) SUGAHARA, Tomohiro (D1) HONJO, Kazuma (M2) NAKANO, Tomohiro (M2)

YANAGISAWA, Tatsuya (M2) OMATSU, Yamato (M1)

Scope of Research

Organic chemistry has mainly developed with the use of second-row elements such as carbon, oxygen, and nitrogen, while the synthesis and isolation of the heavier congeners of typical organic molecules as stable compounds has been a dream for organic chemists. Our main research interest is the elucidation of the similarities and differences in structures and reactivities between organic compounds and the corresponding heavier congeners. These studies are interesting and important from the standpoints of fundamental chemistry, and open the way to more extensive application of main group chemistry.

KEYWORDS

Steric Protection Kinetic Stabilization Low-coordinated Species Heteroatom

Transition Metal Complexes



Selected Publications

Nagata, K.; Murosaki, T.; Agou, T.; Sasamori, T.; Matsuo, T.; Tokitoh, N., Activation of Dihydrogen by Masked Doubly Bonded Aluminum Species, Angew. Chem. Int. Ed., 55, 12877-12880 (2016).

Suzuki, Y.; Sasamori, T.; Guo, J.-D.; Nagase, S.; Tokitoh, N., Isolation and Ambident Reactivity of a Chlorogermylenoid, Chem. Eur. J., 22, 13784-13788 (2016).

Agou, T.; Ikeda, S.; Sasamori, T.; Tokitoh, N., Synthesis and Structure of Lewis-Base-Free Phosphinoalumane Derivatives, Eur. J. Inorg. Chem. 2016, 623-627 (2016).

Sasamori, T.; Villalba Franco, J. M.; Guo, J.-D.; Sugamata, K.; Nagase, S.; Streubel, R.; Tokitoh, N., Selenium-Substituted Phosphaalkenes Obtained through 1,2-Elimination of Chlorosilanes from Selenenylchlorophosphines, Eur. J. Inorg. Chem., 2016, 678-684 (2016).

Sugahara, T.; Guo, J.-D.; Sasamori, T.; Karatsu, Y.; Furukawa, Y.; Ferao, A. E.; Nagase, S.; Tokitoh, N., Reaction of a Stable Digermyne with Acetylenes: Synthesis of a 1,2-Digermabenzene and a 1,4-Digermabarrelene, Bull. Chem. Soc. Jpn., 89, 1375-1384 (2016).

Activation of Dihydrogen by Masked Doubly Bonded Aluminum Species

Activation of dihydrogen (H₂) with transition-metal complexes is an important fundamental chemical transformation and has been extensively investigated for several decades. In contrast, main-group element compounds had been considered as ineffective for the splitting of H₂ for a long time, until the discoveries of the catalyst-free hydrogenation of heavier Group 14 triply bonded compounds, dimetallynes, and of the frustrated Lewis pair (FLP)-catalyzed hydrogenation. Thereafter, H₂ activation using multiply bonded and low-coordinate compounds of heavier main-group elements have been reported. Among various heavier main-group elements, aluminum is a promising candidate as the key component of the H₂ activation reagents, as shown by the recent reports on small-molecule activation with aluminum-based FLPs.

We have already reported the reactivity of barrelene-type dialumane 1 as the masked form of 1,2-diaryldialumene 2. At ambient temperature, 1 most certainly undergoes retro [2+4] cycloaddition to generate 2, which can be trapped by various trapping reagents. These findings are indicative of the possibility of H_2 activation by using 1 as the masked forms of the corresponding dialumenes 2.

Reactions of 1 with H₂ afforded dihydroalumanes ArAlH₂ at room temperature (Ar: bulky aryl groups).




Figure 1. Reactions of barrelene-type dialumane 1 with dihydrogen and the Structure of 3 (Ar = Tbb).

These dihydroalumanes form hydrogen-bridged dimers $[ArHAl(\mu-H)]_2$ 3 in the crystalline state, while a monomer–dimer equilibrium was suggested in solution. Dialumenes 2 generated from 1 are the putative active species in the cleavage of H_2 .

Isolation and Reactivity of a Chlorogermyl- enoid

Metallylenoids of the general formula R_2EXM (E =group 14 element, X =anionic leaving group, M =metal) represent the silicon analogs of carbenoids (R_2CXM), which may exhibit both carbanion (R_2CX^-) and carbene (R_2C :) character. Silylenoids (R_2SiXM) as well as silylenes (R_2Si :) have been postulated as important intermediates in the reduction of dihalosilanes towards silylenes and/or oligo- and polysilanes. In contrast to the variety and rich chemistry of halosilylenoids, heavier halogermylenoid (R_2GeXM) analogs and their associated chemistry still remains scarce.

Treatment of 2,5-di(3,5-tert-butylphenyl)-1-lithio-ferrocene (Fc*Li) with GeCl₂·dioxane afforded the corresponding chlorogermylenoid that exhibited an ambident reactivity in different solvents; it displayed a behavior characteristic for a dichlorogermylene anion in THF, while it exhibited the typical reactivity of a chlorogermylene in toluene. X-Ray diffraction analysis of a single crystal of this chlorogermylenoid, obtained from recrystallization in THF, revealed a separated-ion-pair structure in the solid state.




Figure 2. The structure of [Li⁺(thf)₄][Fc*GeCl₂]⁻.

Division of Synthetic Chemistry– Structural Organic Chemistry –

http://www.scl.kyoto-u.ac.jp/~kouzou/en/index.html



Prof MURATA, Yasujiro (D Eng)



Assoc Prof WAKAMIYA, Atsushi (D Eng)



Assist Prof MURATA, Michihisa (D Eng)



PD AHAREN, Tomoko (Ph D)



PD LEE, Jaehyun (D Eng)

Researchers(pt)

ARAKAWA, Noriko JUNG, Mina ISHIKURA, Yasuhisa

Assist Techn Staff SHIMAZAKI, Ai

Students

CHAOLUMEN (D3)
FUTAGOISHI, Tsukasa (D3)
SHIMOGAWA, Hiroyuki (D3)
ZHANG, Rui (D3)
HASHIKAWA, Yoshifumi (D2)
OZAKI, Masashi (D1)
NAKAMURA, Tomoya (D1)

KAJI, Shoji (M2) OKAZAKI, Shuhei (M1) TSUKAO, Masahiro (M1) MURATA, Naoki (UG) YAKUMARU, Shinya (UG) OKAMOTO, Shu (UG)

Scope of Research

Fundamental studies are being conducted for the creation of new functional π -systems with novel structures and properties, and for evaluation of their application as organic semi-conducting materials for photovoltaic and electroluminescent

devices. The major subjects are: 1) organochemical transformation of fullerenes C_{60} and C_{70} , specifically organic synthesis of endohedral fullerenes by the technique of molecular surgery; 2) generation of ionic fullerene species and their application for the synthesis of functional material; 3) synthesis of new carbon-rich materials by the use of transition metal complex; and 4) creation of new functional π -materials with unique photoelectric properties.



π-Conjugated Systems Functional Materials Organic Solar Cells Endohedral Fullerenes Perovskite-Based Solar Cells



Selected Publications

Zhang, R.; Murata, M.; Aharen, T.; Wakamiya, A.; Shimoaka, T.; Hasegawa, T.; Murata, Y., Synthesis of a Distinct Water Dimer inside Fullerene C₇₀, *Nat. Chem.*, **8**, 435-441 (2016).

Nishimura, H.; Fukushima, T.; Wakamiya, A.; Murata, Y.; Kaji, H., The Influence of Quasiplanar Structures of Partially Oxygen-Bridged Triphenylamine Dimers on the Properties of their Bulk Films, *Bull. Chem. Soc. Jpn.*, **89**, 726-732 (2016).

Hashikawa, Y.; Murata, M.; Wakamiya, A.; Murata, Y., Synthesis and Properties of Endohedral Aza[60] fullerenes: $H_2O@C_{59}N$ and $H_2@C_{59}N$ as Their Dimers and Monomers, *J. Am. Chem. Soc.*, **138**, 4096-4104 (2016).

Satou, M.; Nakamura, T.; Aramaki, Y.; Okazaki, S.; Murata, M.; Wakamiya, A.; Murata, Y., Near-Infrared Emissive Donor–Acceptor-Type Molecules Containing Thiazole-Fused Benzothiadiazole as an Electron-Acceptor Moiety, *Chem. Lett.*, **45**, 892-894 (2016).

Murata, M.; Kaji, S.; Nishimura, H.; Wakamiya, A.; Murata, Y., Efficient Synthesis of One- and Two-Dimensional Multimetallic Gold-Bis(dithiolene) Complexes, *Eur. J. Inorg. Chem.*, **2016**, 3228-3232 (2016).

Hashikawa, Y.; Murata, M.; Wakamiya, A.; Murata, Y., Water Entrapped inside Fullerene Cages: A Potential Probe for Evaluation of Bond Polarization, *Angew. Chem. Int. Ed.*, **55**, 13109-13113 (2016).

Synthesis of a Distinct Water Dimer inside Fullerene C₇₀

The water dimer is an ideal chemical species with which to study hydrogen bonds. Owing to the equilibrium between the monomer and oligomer structure, however, selective generation and separation of a genuine water dimer has not yet been achieved. We have developed a synthetic method that leads to the successful encapsulation of one or two water molecules inside fullerene C₇₀. These endohedral C₇₀ compounds offer the opportunity to study the intrinsic properties of a single water molecule without any hydrogen bonding, as well as an isolated water dimer with a single hydrogen bond between the two molecules. The structure of H₂O@C₇₀ was unambiguously determined by XRD analysis, showing that the H₂O molecule was located at the off-center position of C₇₀. Furthermore, in the case of (H₂O)₂@C₇₀, a hydrogen bond was observed between the two H₂O molecules, and their rapid interchange was supported by NMR measurements and theoretical calculations. Infrared spectroscopy suggested the possible qualitative recognition of each water-encapsulating C₇₀ species based on their unique and characteristic spectra.




Figure 1. The structures of $H_2O@C_{70}$ and $(H_2O)_2@C_{70}$.

Near-infrared Emissive Donor-Acceptor-type Molecules Containing Thiazole-fused Benzothiadiazole as an Electron-acceptor Moiety

Fluorophores with strong emission in the near-infrared (NIR) region have received attention as NIR organic light-emitting diodes (OLEDs), which find applications in night vision-readable displays and sensors, as well as in fluorescent probes for bioimaging. In addition to conventional dyes, donor (D)—acceptor (A)-type molecules have attracted considerable interest, as their optical properties can be easily tuned by varying the D and A units. In this work, two D–A-type molecules containing thiazole-fused benzothiadiazole (BT) as the acceptor moiety were designed and synthesized. The thiazole-fused BT skeleton enables

fine-tuning of the solubility and the electronic structure as well as two-dimensional extension of the π -conjugation. The combination with common donor units such as diphenylaminothiophene or triphenylamine-substituted thiophene resulted in the formation of D–A–D-type molecules, which exhibited fluorescence in the NIR region.




Figure 2. The structures of model compounds for the D-A-D type molecules based on a thiazole-fused BT skeleton.

Efficient Synthesis of One- and Two-Dimensional Multimetallic Gold-Bis (dithiolene) Complexes

Multimetallic complexes having π-conjugated dithiolate ligands have recently received attention because of their unique solid-state properties, which include high conductivity and magnetism, as well as their potential use in organic thermoelectric materials. We demonstrated efficient and versatile methods for the synthesis of 1D and 2D multimetallic gold-bis(dithiolene) complexes with well-defined structures by using a dichlorogold(III) complex having a benzene-1,2-dithiolate ligand as a terminal unit. By using benzene-1,2,4,5-tetrathiolate as a bridging unit, a 1D rod-shaped dimetallic complex was synthesized in good yield. Furthermore, on the basis of this synthetic strategy, we prepared a 2D trimetallic complex with a starburst structure by using a benzenehexathiolate bridging unit. The new synthetic methods can offer a way to prepare a wide variety of unprecedented 1D and 2D multimetallic gold-bis(dithiolene) complexes.




Figure 3. The structures of 1D dimetallic and 2D trimetallic gold—bis(dithiolene) complexes.

Division of Synthetic Chemistry – Synthetic Organic Chemistry –

http://www.fos.kuicr.kyoto-u.ac.jp/EnglishTop/English.top.html



Prof KAWABATA, Takeo (D Pharm Sc)



Assoc Prof FURUTA, Takumi (D Pharm Sc)



Assist Prof UEDA, Yoshihiro (D Pharm Sc)



Program-Specific Assist Prof YOSHIDA, Keisuke (D Pharm Sc)



Techn Staff FUJIHASHI, Akiko



PD HARISADHAN, Ghosh (Ph D)

Researcher(pt)

HAYASHI. Kazuhiro



PD YELLA, Ramesh (Ph D)

Students

HIRATA, Atsushi (D3) LU, Wenjie (D3) KASAMATSU, Koji (D3) KURIBAYASHI, Toshifumi (D3) HAYASAKA, Manami (D3) ARAI, Kenta (D2) OTSUKI, Haruka (D2) ORITA, Takuya (M2) GONDOH, Naruhiro (M2) SHIBAYAMA, Hiromitsu (M2) SHIMODA, Shotaro (M2) MATAYOSHI, Aki (M2) SHIMMIYA, Ruri (M1)
NAGAISHI, Masaru (M1)
NINOMIYA, Ryo (M1)
YAMAMOTO, Satoru (M1)
KUMEGAWA, Ryo (M1)
TAKASHIMA, Megumi (M1)
YANG, Baochen (RS)
CHEN, Gong (RS)
WANG, Shuo (RS)
HASHIMOTO, Hisashi (UG)
FUJIMOTO, Takumi (UG)
MURAI, Takuya (UG)

Scope of Research

The research interests of this laboratory include the development of advanced molecular transformation, total synthesis of biologically active products, and molecular recognition. Programs are active in the following areas: 1) asymmetric alkylation of carbonyl compounds based on "memory of chirality", 2) organocatalysis for fine organic syntheses, 3) synthesis of unusual amino acids and nitrogen heterocycles, 4) regioselective functionalization of carbohydrates, and 5) the structural and functional investigation of heterochiral oligomers.

KEYWORDS

Organocatalysis Regioselective Functionalization Dynamic Chirality Unusual Amino Acid Molecular Recognition

Selected Publications

Kawabata, T.; Moriyama, K.; Kawakami, S.; Tsubaki, K., Powdered KOH in DMSO: An Efficient Base for Asymmetric Cyclization via Memory of Chirality at Ambient Temperature, *J. Am. Chem. Soc.*, **130**, 4153-4157 (2008).

Kawabata, T.; Jiang, C.; Hayashi, K.; Tsubaki, K.; Yoshimura, T.; Majumdar, S.; Sasamori, T.; Tokitoh, N., Axially Chiral Binaphthyl Surrogates with an Inner N-H-N Hydrogen Bond, *J. Am. Chem. Soc.*, **131**, 54-55 (2009).

Yoshida, K.; Furuta, T.; Kawabata, T., Organocatalytic Chemoselective Monoacylation of 1, *n*-Linear Diol, *Angew. Chem. Int. Ed.*, **50**, 4888-4892 (2011).

Hamada, S.; Furuta, T.; Wada, Y.; Kawabata, T., Chemoselective Oxidation by Electronically Tuned Nitroxyl Radical Catalysts, *Angew. Chem. Int. Ed.*, **52**, 8093-8097 (2013).

Tomohara, K.; Yoshimura, T.; Hyakutake, R.; Yang, P.; Kawabata, T., Asymmetric α-Arylation of Amino Acid Derivatives by Clayden Rearrangement of Ester Enolates via Momory of Chirality, *J. Am. Chem. Soc.*, **135**, 13294-13297 (2013).

Yoshimura, T.; Tomohara, K.; Kawabata, T., Asymmetric Induction via Short-Lived Chiral Enolates with Chiral C-O Axis, *J. Am. Chem. Soc.*, 135, 7102-7105 (2013).

Takeuchi, H.; Mishiro, K.; Ueda, Y.; Fujimori, Y.; Furuta, T.; Kawabata, T., Total Synthesis of Ellagitannins via Regioselective Sequential Functionalization of Unprotected Glucose, *Angew. Chem. Int. Ed.*, **54**, 6177-6180 (2015).

Ueda, Y.; Furuta, T.; Kawabata, T., Final-Stage Site-Selective Acylation for the Total Syntheses of Multifidosides A-C, *Angew. Chem. Int. Ed.*, **54**, 11966-11970 (2015).

Late-stage Functionalization of Natural Products by Organocatalysis: Site-Selective Dimerization of 10-Deacetylbaccatin III

Late-stage site-selective functionalization of biologically active compounds has been receiving increasing attention because it enables diversification of biologically active compounds retaining their original activity. Here, we report application of our strategy for organocatalytic site-selective acylation to dimerization of 10-deacetylbaccatin III. 10-Deacetylbaccatin III is a natural terpenoid available in a relatively abundant amount and used as a key intermediate for clinically widely used antitumor agents, taxol and taxotere. Dimerization of bioactive compounds is potentially useful strategy toward the discovery of the agents with improved activity profiles including promising activity against the native-compound-resistant infections and cancers. The organocatalytic site-selective acylation of 10-Deacetylbaccatin III with a mixed anhydride prepared from undecanedioic acid and pivaloyl chloride took place at C(10)-OH in the presence of an organocatalyst to provide its dimer as a pure regioisomer in 40% yield. The site-selectivity was assumed to be controlled by the catalyst because the C(7)-OH has been known to be intrinsically more reactive hydroxy group.

Asymmetric α-Arylation of α-Amino Acid Derivatives via Memory of Chirality

We have studied asymmetric synthesis via memory of chirality (MOC). A salient feature of the MOC strategy is that asymmetric reactions take place via axially chiral enolates with limited half-lives of racemization. Because the rate of enolate-racemization strongly depends on the reaction temperature, the reactions via MOC are usually performed at low temperatures to minimize the enolate racemization. While we have successfully developed several intermolecular asymmetric reactions and *intramolecular* asymmetric α -arylation of α -amino acid derivatives via the MOC strategy, the development of asymmetric *intermolecular* α -arylation has been unsuccessful so far. This was assumed to be because usual methods for α -arylation of

enolates often requires long reaction times at high temperature, which should cause significant racemization of the intermediary chiral enolates. We describe here a solution to this long-standing problem in the MOC study. We employed arynes as reactive electrophilic aryl species at low temperatures, and successfully developed the target reaction. Treatment of a solution of the amino acid derivative and the aryne precursor with potassium hexamethyldisilazide (KHMDS) followed by phenyl lithium (PhLi) at -78 °C gave benozocyclobutenones with tetra-substituted carbon with retention of configuration in up to 99% ee. The reaction was assumed to proceed via intermolecular asymmetric α-arylation of axially chiral enolate A generated by the amino acid derivative and KHMDS with arynes generated by the aryne precursors and PhLi, followed by intramolecular C-acylation of resulting aryl metallic species **B**.

Asymmetric Intramolecular C-H Insertion Promoted by Dirhodium(II) Carboxylate Catalyst Bearing Axially Chiral Amino Acid Derivatives

The asymmetric intramolecular C–H insertion of a rhodium carbenoid intermediate has attracted much attention as a powerful synthetic tool for constructing chiral cyclic compounds. A dirhodium(II) carboxylate catalyst bearing axially chiral amino acid derivatives is prepared. This catalyst is effective in the asymmetric intramolecular C–H insertion of α -aryl- α -diazoacetates to α -aryl- β -substituted γ -lactones with a reasonable level of diastereo- and enantioselectivity, especially in the reaction of phenyl and β -naphthyl substituted substrates.

Division of Synthetic Chemistry - Advanced Inorganic Synthesis -

http://www.scl.kyoto-u.ac.jp/~teranisi/index_E.html



Prof TERANISHI, Toshiharu SAKAMOTO, Masanori (D Eng)



Assoc Prof (D Eng)



Assist Prof SATO, Rvota (D Sc)



SARUYAMA, Masaki (DSc)



Program-Specific Assist Prof Program-Specific Assist Prof TRINH, Thang Thuy (Ph D)



SPD KAWAWAKI, Tokuhisa (D Eng)



PD NISHINO, Toshio (D Sc)



PD HSU, Shih-Chen (D Sc)



PD CHEN, Lihui (D Sc)

Students

KIM, Sungwon (D3) EGUCHI, Daichi (D3) LIAN, Zichao (D2)

KIMURA, Masato (D1) MATSUMOTO, Kenshi (D1) TABUCHI, Michimasa (M2)

EGAWA, Shigehisa (M1) OKAMOTO, Yasuo (M1)


MIZUNO, Hiroki (M1) LI, Yanhui (M1)

Scope of Research

We are focusing on the precise synthesis of inorganic nanoparticles by controlling the primary (size, shape, composition, etc.) and secondary (spatial arrangement) structures to tune properties such as electron confinement, carrier oscillation, spin, and catalysis. These high-quality inorganic nanoparticles are applied to both high-performance nanodevices (e.g., single electron transistor, plasmon waveguide, and nanocomposite magnet) and photo-energy conversion materials (e.g., overall water splitting and solar cell).

KEYWORDS

Inorganic Nanoparticles Single Electronics Plasmonics Nanocomposite Magnet Photocatalysts



Selected Publications

Sakamoto, M.; Inoue, K.; Okano, M.; Saruyama, M.; Kim, S.; So, Y.-G.; Kimoto, K.; Kanemitsu, Y.; Teranishi, T., Light-Stimulated Carrier Dynamics of CuInS₂/CdS Heterotetrapod Nanocrystals, Nanoscale, 8, 9517-9520 (2016).

Wu, H.-L.; Sato, R.; Yamaguchi, A.; Kimura, M.; Haruta, M.; Kurata, H.; Teranishi, T., Formation of Pseudomorphic Nanocages from Cu₂O Nanocrystals through Anion Exchange Reactions, Science, 351, 1306-1310 (2016).

Nano-pseudomorphic Chemistry: Anionic Frameworks Dictate Crystal Systems in Sequential Ion Exchange Reactions

Ionic crystals have been widely used as photo-functional materials such as photocatalysts and photoelectric conversion materials, which are determined by the constituent elements, morphologies, and crystal structures. Because the stable crystal structures of the ionic crystals follow their phase diagrams, it has been difficult to chemically synthesize the high-temperature stable phases. In contrast to the bulk form, chemical conversion of ionic nanocrystals (NCs) via ion exchange reactions can overcome the difficulties associated with controlling the size, shape, chemical composition, and crystal structure in conventional syntheses. In our research, we discovered that the retained shape of the parent-NCs in sequential ion exchange reactions provided an opportunity to obtain non-equilibrium unique structures and even new structures of ionic NCs, these morphologically retained products being known as "pseudomorphs".

We investigated the crystal structures of the pseudomorphic nanocages, which were formed by the anion exchange ($O^{2-} \rightarrow S^{2-}$) of hexahedral Cu₂O NCs




Figure 1. (Left hand) The rhombic dodecahedral $Cu_{1.75}S$ nanocages with a triclinic phase are formed by the anion exchange $(O^{2-} \rightarrow S^{2-})$ and the subsequent etching of rhombic dodecahedral Cu_2O NCs enclosed with {110} planes at ambient conditions. Further cation exchange $(Cu^+ \rightarrow Cd^{2+}, Zn^{2+})$ gives the hexahedral CdS or ZnS with a hexagonal phase. (Right hand) The hexahedral $Cu_{1.8}S$ nanocages with a cubic phase are formed by the anion exchange $(O^{2-} \rightarrow S^{2-})$ and the subsequent etching of hexahedral Cu_2O NCs enclosed with {100} planes at ambient conditions. Further cation exchange $(Cu^+ \rightarrow Cd^{2+}, Zn^{2+})$ gives the hexahedral CdS or ZnS with a cubic phase.

enclosed with {100} planes and rhombic dodecahedral Cu₂O NCs enclosed with {110} planes at ambient conditions. It was discovered that the shape-dependent anionic framework (surface anion sublattice and stacking pattern) of Cu₂O NCs determined the crystal system of anion-exchanged products (Figure 1). In other words, the cubic system can transform into not only the cubic system but also the triclinic and hexagonal systems. Furthermore, we found that the resulting nanocages have the multiply-twinned structure. A high-temperature stable phase such as wurtzite ZnS (> 1020 °C) was also obtained with this method at ambient conditions. We envisage that this pseudomorphic transformation method could be applicable to a number of other ionic NCs and thin films at ambient temperatures to find the novel properties and functions.

Porphyrin Derivative-Protected Gold Cluster with a Pseudo-Tetrahedral Shape

The shape of a nanomaterial plays an important role in controlling the packing arrangement and properties of the assembly that it forms. In the case of metal clusters, the shape of the metal core is veiled by a flexible organic ligand with a comparable size to that of the metal core. Here, we controlled the overall shape of ligand-protected gold clusters (AuCs) using a rigid, planar molecule as a ligand by following a simple geometrical relationship between an inscribed sphere and a circumscribed polyhedron.

For the rigid, planar molecule, we synthesized a new porphyrin derivative that could strongly attach to the AuCs in a face-coordination fashion (Figure 2a). We synthesized AuCs face-coordinated by four ZnSC₂P-SS ligands with a core size of 1.0±0.2 nm. Structural assignment using MALDITOF MS, ICP-AES, and UV-vis-NIR absorption spectra indicated that the ZnSC₂P-SS/AuCs have a pseudo-tetrahedral shape (Figure 2b). Thus, we have demonstrated that a simple geometrical relationship between an inscribed sphere and a circumscribed polyhedron can be applied to control the shape (symmetry) of metal clusters with a size of less than 2 nm. The present results pave the way to a new concept for controlling the pseudo-symmetry of ligand-protected metal clusters.




Figure 2. (a) single X-ray crystal structures of SC_2P and $ZnSC_2P$ -SS from top view and side view. Hydrogen atoms have been omitted for clarity. (b) MALDI-TOF spectrum of $ZnSC_2P$ -SS/AuCs. Insert is schematic illustration of the $ZnSC_2P$ -SS/AuCs with pseudo-tetrahedron shape.

Division of Materials ChemistryChemistry of Polymer Materials

http://www.cpm.kuicr.kyoto-u.ac.jp/



Prof TSUJII, Yoshinobu (D Eng)



Assoc Prof OHNO, Kohji (D Eng)



Assist Prof SAKAKIBARA, Keita (D Agr)

Assist Techn Staffs NAKANISHI, Yohei TANISHIMA, Miho YAHATA, Yoshikazu

Researchers(pt)

MORIKI, Yoshito SAKAUE, Motokazu

NAKAJIMA, Yuuki YANADA, Mizuho NAKAMICHI, Kimiyo

Proj Res

MATSUKAWA, Kimihiro (D Eng)* *Res of Kyoto Inst Technol

Students

KINOSE, Yuji (D2) SHIMIZU, Yoshihiko (D2) EGI, Kohki (M2) YAMASHITA, Kohei (M2)

YASUDA, Shuma (M2) ZHAO, Chenzhou (M2) FUJIMORI, Tomoya (M1) HISATOMI, Tatsuya (M1) NISHIUMI, Kosuke (M1) HARAYAMA, Seiya (UG) MURAJI, Koichi (UG) NIWA, Ryota (UG)

Scope of Research

We perform kinetic and mechanistic analyses toward understanding chemical and physicochemical reactions occurring in polymerization systems and better routes for synthesis of well-defined polymers. In particular, new well-defined polymers or polymer assemblies are prepared by living polymerization techniques, and their structure–properties relationships are precisely analyzed. Projects in progress include: 1) kinetics and mechanisms of living radical polymerization (LRP); 2) synthesis of new polymeric materials by living polymerizations and their structure/properties studies; and 3) synthesis, properties, and applications of concentrated polymer brushes (CPB).



KEYWORDS

Precision Polymerization Living Radical Polymerization Polymer Brush Hybrid Materials Biointerface

Selected Publications

Tsujii, Y.; Nomura, Y.; Okayasu, K.; Gao, W.; Ohno, K.; Fukuda, T., AFM Studies on Microtribology of Concentrated Polymer Brushes in Solvents, *J. Phys.: Conf. Ser.*, **184**, 012031 (2009).

Arita, T.; Kayama, Y.; Ohno, K.; Tsujii, Y.; Fukuda, T., High-Pressure Atom Transfer Radical Polymerization of Methyl Methacrylate for Well-Defined Ultrahigh Molecular-Weight Polymers, *Polymer*, *49*, 2426-2429 (2008).

Tsujii, Y.; Ohno, K.; Yamamoto, S.; Goto, A.; Fukuda, T., Structure and Properties of High-Density Polymer Brushes Prepared by Surface-Initiated Living Radical Polymerization, *Adv. Polym. Sci.*, **197**, 1-45 (2006).

Ohno, K.; Morinaga, T.; Takeno, S.; Tsujii, Y.; Fukuda, T., Suspension of Silica Particles Grafted with Concentrated Polymer Brush: Effects of Graft Chain Length on Brush Layer Thickness and Colloidal Crystallization, *Macromolecules*, **40**, 9143-9150 (2007).

Ohno, K.; Morinaga, T.; Koh, K.; Tsujii, Y.; Fukuda, T., Synthesis of Monodisperse Silica Particles Coated with Well-Defined, High-Density Polymer Brushes by Surface-Initiated Atom Transfer Radical Polymerization, *Macromolecules*, **38**, 2137-2147 (2005).

Structure Analysis of Composite Membranes of Polymer-brush-afforded Nanoparticles and Ionic Liquid

Previously, we succeeded in fabricating a solid membrane of high ionic conductivity using self-assembled nanoparticles with CPBs in an ionic liquid (IL) and in developing a novel bipolar-type lithium-ion rechargeable batteries, where CPB represents the concentrated polymer brush, successfully and systematically synthesized by living radical polymerization techniques. This success was caused by the CPB effects (including super lubrication) originating from the highly stretched-chain conformation and high-dense segmental density in a solvent. In this study, we achieved in analyzing the higher-order structure of the composite membranes using concentrated ionic liquid-type polymer brush-modified nanoparticles (PSiP) and IL by ultra-small angle X-ray scattering (USAXS). The results showed that the higher-order structure of PSiP/IL composites changed from disorder to random hexagonal-closed-packing (rhcp), and face-centered-cubic (fcc) as the concentration of PSiP increased. As concentrated polymer brushes have unique properties such as high elasticity in a good solvent, it can be concluded that these results are applied to the model for particle assemblies with high-elasticity.




Figure 1. USAXS and FE-SEM images of composite membranes of concentrated ionic liquid-type polymer brush-modified nanoparticles and ionic liquid.

Surface Engineering of Cellulose Nanofiber by Adsorption of Diblock Copolymer Dispersant for Green Nanocomposite Materials

An effective approach for the dispersion of hydrophilic cellulose nanofiber (CNF) in hydrophobic high-density polyethylene (HDPE) is presented using adsorption of a diblock copolymer dispersant. The dispersant consists of both resin compatible poly(lauryl methacrylate) (PLMA) and cellulose interactive poly(2-hydroxyethyl methacrylate) blocks. The PLMA-adsorbed CNFs are characterized by FT-IR and contact angle measurement, revealing successful hydrophobization. X-ray CT imaging shows there are apparently less CNF aggregates in the nanocomposites if adding amount of the dispersant was enough. The good dispersion results in a high mechanical reinforcement, corresponding to 140 % higher Young's modulus and 84 % higher tensile strength than the neat HDPE. This study represents a potential approach to mediate CNF-CNF and CNF-resin interactions and to influence the CNF dispersion significantly without chemical functionalization, which may be compatible with a low-cost and environmentally green process and therefore an industrial application. Based on this concept, simultaneous nano-fibrillation compounding (SFC) process has been developed.




Figure 2. X-ray CT image and strain-stress curve of CNF-reinforced HDPE nanocomposite materials.

Division of Materials ChemistryPolymer Controlled Synthesis

http://os.kuicr.kyoto-u.ac.jp/index.html



Prof YAMAGO, Shigeru (D Sc)



Assoc Prof TOSAKA, Masatoshi (D Eng)



Program-Specific Assoc Prof NAKAMURA, Yasuyuki (D Sc)



Assist Prof KAYAHARA, Eiichi (D Eng)




Assist Prof HASHIMOTO, Sigma (D Eng)



PD PATEL, Vijay Kumar (D Sc)



PD LI, Wenjuan (D Sc)



Guest Scholar
HONORE, Jean-Claude
Aix-Marseille University, France, 4 March–18 March

Students

LU, Yangtian (D2)
SUN, Liansheng (D2)
KONO, Taro (D1)
FAN, Weijia (D1)
OII Rui (D1)

KOBAYASHI, Go (D1) PARK, Beomsu (D1) NOGUCHI, Naoki (M2) HUA SHEN, Zhiren (M2) SEKI, Toyomitsu (M2) SHITARA, Daichi (M1) HAYASHI, Toshiki (M1) YANO, Yuko (M1) LI, Xiaopei (M1) JEONG, Jaeyoung (RS) LIN, Yiming (RS)
YAO, Cheng (RS)
MIYOSHI, Shun-ichiro (UG)
OUE, Shohei (UG)
MURATA, Masahiro (UG)

Scope of Research

Our research focuses on creation of new organic molecules with potential as key reagents and materials for future science and technologies. Furthermore, we have been developing new organic and polymeric materials based on our tailor-made molecules. For example, we are developing a new living radical polymerization method using heavier heteroatom compounds as controlling agents. Another topic is the synthesis of cycloparaphenylenes, hoop-shaped π -conjugated molecules, based on new synthetic strategies. We also study various condensed states of polymers by both static and dynamic methods to understand the relationship between structure and physical properties.

KEYWORDS

Organic Synthesis Polymer Synthesis Living Radical Polymerization Polymer Properties Curved π -Conjugated Molecules

Selected Publications

Fan, W.; Nakamura, Y.; Yamago, S., Synthesis of Multivalent Organotellurium Chain Transfer Agents by Post-modification and Their Applications in Living Radical Polymerization, *Chem. Eur. J.*, 22, 17006-17010 (2016).

Nakamura, Y.; Lee, R.; Coote, M.; Yamago, S., Termination Mechanism of the Radical Polymerization of Acrylates: Elucidation of the Reactivity of Polymer-end and Mid-chain Radicals, *Macromol. Rapid Commun.*, 37, 506-513 (2016).

Kayahara, E.; Patel, V. K.; Mercier, A.; Kündig, E. P.; Yamago, S., Regioselective Synthesis and Characterization of Multinuclear Convex-bounded Ruthenium-[n]Cycloparaphenylene (n = 5 and 6) Complexes, *Angew. Chem. Int. Ed.*, **55**, 302-306 (2016).

Kayahara, E.; Fukayama, K.; Nishinaga, T.; Yamago, S., Size Dependence of [N]Cycloparaphenylenes (N = 5-12) in Electrochemical Oxidation, *Chem. Asian J.*, **11**, 1793-1797 (2016).

Kayahara, E.; Kouyama, T.; Kato, T.; Yamago, S., Synthesis and Characterization of [n]CPP (n = 5, 6, 8, 10, and 12) Radical Cation and Dications: Size-Dependent Absorption, Spin, and Charge Delocalization, J. Am. Chem. Soc., 138, 338-344 (2016).

Practical Synthesis of [n]Cycloparaphenylenes (n = 5, 7–12) by H₂SnCl₄-mediated Aromatization of 1,4-Dihydroxycyclo-2,5-diene Precursors

Cyclic precursors of cycloparaphenylenes (CPPs) containing 1,4-dihydroxy-2,5-cyclohexadien-1,4-diyl units were prepared by modifying the synthesis method of corresponding 1,4-dimethoxy derivatives, which was developed by Jasti and co-workers. Reductive aromatization of the diyl moieties by SnCl₂/2HCl took place under mild conditions and afforded CPPs incorporating 5 or 7–12 phenylene units with good yield. Highly strained [5]CPP was synthesized at more than 0.3 g. 119Sn NMR spectroscopy clarified the in situ formation of an ate complex, H₂SnCl₄, upon mixing a 2:1 ratio of HCl and SnCl₂, which serves as a highly active reducing agent under nearly neutral conditions. When more than two equivalents of HCl, in relation to SnCl₂, were used, acid-catalyzed decomposition of the CPP precursors took place. The stoichiometry of HCl and SnCl₂ was found to be critical in achieving the desired aromatization reaction of highly strained CPP precursors.





Figure 1. Synthesis of [n]Cycloparaphenylenes (n = 5, 7-12).

Synthesis of Multivalent Organotellurium Chain-Transfer Agents by Post-modification and Their Applications in Living Radical Polymerization

Multivalent or functionalized organotellurium chaintransfer agents (CTAs) for living radical polymerization were synthesized by post-modification. The synthesis involving the condensation between a carboxylic-acid-functionalized CTA and various amines proceeded in excellent yields without affecting the reactive tellurium moiety. The CTAs exhibited high synthetic versatility for radical polymerization and gave structurally well-controlled polymers, such as multiarmed polymers, from various monomers. Because all new CTAs are easily available on a large scale by simple purification, the current method significantly facilitates macromolecular engineering based on organotellurium-mediated radical polymerization (TERP).



Division of Materials ChemistryInorganic Photonics Materials

http://mizuochilab.kuicr.kyoto-u.ac.jp/indexE.html



Prof MIZUOCHI, Norikazu (D Sc)



Assist Prof MASAI, Hirokazu (D Eng)



Assist Prof MORISHITA, Hiroki (D Eng)



Assist Prof FUJIWARA, Masanori (D Sc)



Program-Specific Res SIMO-OKA, Takaaki (D Sc)

Lect (pt)

TOKUDA, Youmei (D Eng)

Faculty of Education, Shiga University

Students

KOBAYASHI, Satoshi (D3)* HAYASHI, Kan (D2)* KANEKO, Syunichi (M2) TAKAHASHI, Yuuya (M2) TORIMOTO, Aya (M2) MARUYAMA, Yuuichi (M2)* MORI, Teppei (M2)* MURAI, Takuya (M2)* DANJYO, Takuya (M1) FUJIMOTO, Syou (M1) MIMA, Daisuke (M1) ASHIDA, Takaki (UG) ISHIMOTO, Shintaro (UG)

*Special research students

Scope of Research

NV center in diamond has been extensively interested because the single spin of it can be manipulated and detected at room temperature. Furthermore, spin coherence time of the NV center is very long. The coherence time is the time to retain coherence (superposition state) and directly relates to the sensitivity of sensor of magnetic field, electric field and temperature. Therefore, the unique and excellent properties of the NV center are expected to be applied for quantum computing, quantum communication, bio-imaging, and high-sensitive sensor with nano-scale resolution.



KEYWORDS

Diamond Quantum Materials NV Center

Quantum Sensing Quantum Information Science

Selected Publications

Fukui, T.; Doi, Y.; Miyazaki, T.; Miyamoto, R.; Kato, H.; Matsumoto, T.; Makino, T.; Yamasaki, S.; Morimoto, R.; Tokuda, N.; Hatano, M.; Sakagawa, Y.; Morishita, H.; Tashima, T.; Miwa, S.; Suzuki, Y.; Mizuochi, N., Perfect Selective Alignment of Nitrogen-vacancy Center in Diamond, *Appl. Phys. Express*, 7, 055201 (2014).

Doi, Y.; Makino, T.; Kato, H.; Takeuchi, D.; Ogura, M.; Okushi, H.; Morishita, H.; Tashima, T.; Miwa, S.; Yamasaki, S.; Wrachtrup, J.; Suzuki, Y.; Mizuochi, N., Deterministic Electrical Charge State Initialization of Single Nitrogen-vacancy Center in Diamond, *Phys. Rev. X*, **4**, 011057 (2014).

Mizuochi, N.; Makino, T.; Kato, H.; Takeuchi, D.; Ogura, M.; Okushi, H.; Nothaft, M.; Neumann, P.; Gali, A.; Jelezko, F.; Wrachtrup, J.; Yamasaki, S., Electrically Driven Single Photon Source at Room Temperature in Diamond, *Nat. Photonics*, **6**, 299-303 (2012).

Zhu, X.; Saito, S.; Kemp, A.; Kakuyanagi, K.; Karimoto, S.; Nakano, H.; Munro, W. J.; Tokura, Y.; Everitt, M. S.; Nemoto, K.; Kasu, M.; Mizuochi, N.; Semba, K., Coherent Coupling of a Superconducting Flux-qubit to an Electron Spin Ensemble in Diamond, *Nature*, **478**, 221-224 (2011).

Neumann, P.; Mizuochi, N.; Rempp, F.; Hemmer, P.; Watanabe, H.; Yamasaki, S.; Jacques, V.; Gaebel, T.; Jelezko, F.; Wrachtrup, J., Multipartite Entanglement Among Single Spins in Diamond, *Science*, **320**, 1326 (2008).

Perfect Selective Alignment of Nitrogen-vacancy Center in Diamond

Nitrogen-vacancy (NV) centers in diamond have attracted significant interest because of their excellent spin and optical characteristics for quantum information and metrology. In the diamond crystal structure, the orientations of NV centers are classified according to the alignment along one of four possible crystallographic axes: [111], [111], [111] or [111] (Fig. 1). In most diamond samples, NV centers equally occupy these four orientations. To take advantage of the characteristics, the precise control of the orientation of the N-V axis in the lattice is essential. It is because improvement in readout contrast and a magnetic field sensitivity can be expected when compared to with those of standard samples with equal population of all NV orientations. Furthermore, spin and optical characteristics strongly depend on this orientation. In cases where photoluminescence (PL) is detected from the [111] direction, the PL intensity from N-V centers in which the N-V axis is parallel to [111] (NV | | [111]) is higher than others because electric dipole transitions are allowed for dipoles in the plane perpendicular to the N-V axis. With respect to spin, it is expected to play a key role at the quantum interface with photon and superconducting flux qubits.




Figure 1. Four possible orientations of NV centers in diamond crystal.

We experimentally showed that the orientation of more than 99 % of the NV centers can be aligned along the [111]-axis by CVD homoepitaxial growth on (111)-substrates. We also discuss about mechanisms of the alignment. We examined the atomistic generation mechanism for the NV defect aligned in the [111] direction of C(111) substrates with first-principles electronic structure calculations. Our result enables a fourfold improvement in magnetic-field sensitivity and opens new avenues to the optimum design of NV center devices.

Pure Negatively Charged State of NV Center in n-type Diamond

For the applications of NV center, controlling the charge state is crucial, because optical initialization and readout of the spin state of the NV centers are only possible in negatively charged one (NV⁻). However, upon illumination, the NV centers undergo stochastic charge-state transitions between NV- and a dark state of a neutral charge state of the NV center (NV⁰) as shown in Fig. 2. For example, upon excitation around 580 nm, where NV⁻ has the highest absorption, NV⁻ easily turns into the dark state of NV⁰ and the steady-state-population of NV⁻ decreases to about 10%, which could be revealed from single-shot charge-state measurements. Therefore, illumination at 532 nm is usually used in the experiment of NV⁻. This charge-state interconversion occurs upon illumination at any wavelength, so the steadystate NV⁻ population is always less than 75%–80%. We experimentally showed Fermi level control by phosphorus doping generates $99.4 \pm 0.1\%$ NV⁻ under 1 μ W and 593 nm excitation which is close to maximum absorption of NV-. The pure NV⁻ shows a five-fold increase of luminescence and a four-fold enhancement of an optically detected magnetic resonance under 593 nm excitation compared with those in intrinsic diamond. These results are expected to significantly enhance the versatile potential of NV centers.




Figure 2. Stochastic charge-state transitions between NV^- and a dark state of a neutral charge state of the NV center.

Division of Materials Chemistry - Nanospintronics -

http://www.scl.kyoto-u.ac.jp/~onoweb/



Prof ONO, Teruo (DSc)



Assoc Prof MORIYAMA, Takahiro KUSUDA, Toshiyuki (Ph D)



Techn Staff*



PD KIM, Sanghoon (Ph D)



FUJIMOTO, Junji (D Sc)

*Re-employed Staff



PD KIM, Duck-Ho (Ph D)

Students

KAKIZAKAI, Haruka (D3) YAMADA, Kihiro (D3) TANAKA, Kensho (D2) TANIGUCHI, Takuya (D2) KAMIYA, Michinari (D2) MIZUNO, Hayato (M2)


ANDO, Fuyuki (M2) OKUNO, Takaya (M2) ODA, Kento (M2) ISHIBASHI, Mio (M1) HIRATA, Yushou (M1) LI, Tian (M1)


Scope of Research


Conventional electronics uses only the charge of electrons, while traditional magnetic devices use only the spin degree of freedom of electrons. Aiming at complete control of both charge and spin in single solid-state devices, an emerging field called spintronics is rapidly developing and having an impact on information technologies. By combining the atomic-layer deposition with nanofabrication, we focus on the development of spin properties of various materials and the control of quantum effects in mesoscopic systems for novel spintronics devices.


KEYWORDS


Spintronics Quantum Transport Nano-fabrication Artificial Materials











Selected Publications

Moriyama, T.; Takei, S.; Nagata, M.; Yoshimura, Y.; Matsuzaki, N.; Terashima, T.; Tserkovnyak, Y.; Ono, T., Anti-damping Spin Transfer Torque through Epitaxial Nickel Oxide, Appl. Phys. Lett., 16, 162406 (2015).

Yoshimura, Y.; Kim, K.-J.; Taniguchi, T.; Tono, T.; Ueda, K.; Hiramatsu, R.; Moriyama, T.; Yamada, K.; Nakatani, Y.; Ono, T., Soliton-like Magnetic Domain Wall Motion Induced by the Interfacial Dzyaloshinskii-Moriya Interaction, Nat. Phys., 12, 157-161 (2016).

Matsuo, S.; Takeshita, S.; Tanaka, T.; Nakaharai, S.; Tsukagoshi, K.; Moriyama, T.; Ono, T.; Kobayashi, K., Edge Mixing Dynamics in Graphene p-n Junctions in the Quantum Hall Regime, Nat. Commun., 6, 8066 (2015).

Stigloher, J.; Decker, M.; Korner, H. S.; Tanabe, K.; Moriyama, T.; Taniguchi, T.; Hata, H.; Madami, M.; Gubbiotti, G.; Kobayashi, K.; Ono, T.; Back, C. H., Snell's Law for Spin Waves, Phys. Rev. Lett., 117, 037204 (2016).

Ando, F.; Kakizakai, H.; Koyama, T.; Yamada, K.; Kawaguchi, M.; Kim, S.; Kim, K.-J.; Moriyama, T.; Chiba, D.; Ono, T., Modulation of the Magnetic Domain Size Induced by an Electric Field, Appl. Phys. Lett., 109, 022401 (2016).

Modulation of the Magnetic Domain Size Induced by an Electric Field

The electric field (EF) control of magnetism has intensively investigated because of its potential importance for the reduction of power consumption in magnetic storage devices. In the past few years, we have been focusing on the electric field modulation of the magnetic anisotropy and the Curie temperature in magnetic thin films. However, the microscopic mechanism of why those magnetic properties change with an electric field were not clear in spite of several theoretical suggestions. In this work, we particularly focused on the configuration of the magnetic domains upon the application of the electric field (see Fig. 1a for the detail measurement setup). With the electric field of ± 10 V, we observed the significant change in the domain size (Fig. 1b). Detail analyses on the modification of the magnetic domain size revealed that it is the exchange interaction that is modulated with the electric field and is changing about 50% with ± 10 V. Our results suggest that the EF control of the magnetism is mainly driven by the modification of the exchange interaction which is a fundamental measure determining the magnetic interaction between microscopic spins.




Figure 1. (a) Schematic illustration of the experimental setup (b) Modulation of the magnetic domain size with the electric field of ± 10 V.

Anti-damping Spin Transfer Torque through Antiferromagnetic Material

Spin transfer torque (STT) has been an efficient and promising technique to control magnetization of ferromagnetic materials in modern spintronic devices. This novel technique is based on an interaction between electron spin and local magnetic moments. The same interaction should be conserved in antiferromagnets in which there are microscopic local magnetic moments that compensate each other to exhibit no net magnetization. In this work, we prepared MgO(001) substrate / Pt 5 nm / NiO 10 nm / FeNi 3 nm / SiO₂ 5 nm multilayers, in which the films are epitaxially grown until the NiO layer, and performed a spin torque ferromagnetic resonance (ST-FMR) measurement to quantify the anti-damping spin torque transported between the Pt and the FeNi through the antiferromagnetic NiO layer. A pure spin current is created by the spin Hall effect of the Pt and injected into the NiO. As shown in Figure 2(a), we found that the FMR linewidth monotonously varies with the spin current injection. As the ST-FMR measurement is only sensitive to the linewidth (i.e., magnetic damping) of the FeNi layer, this change in the linewidth in Pt/NiO/FeNi can be interpreted in a way that the spin current is transferred through the NiO and interacts with the FeNi. This intriguing spin current transport can be explained by the angular-momentum transfer mediated by the antiferromagnetic magnons as shown in Figure 2(b). Our results assure that the spin current exerts a spin torque on the NiO magnetic moments and excites their dynamics. The results open up a new field of antiferromagnetic spintronics.




Figure 2. (a) FMR linewidth δH as a function of the dc current. Due to the spin Hall effect of Pt, the spin current injection is proportional to the dc current. (b) Schematic illustration of the N (Pt) / AF (NiO) / F (FeNi) structure in which the anti-damping spin transfer torque takes place. N (Pt) layer injects the spin current \mathcal{F}_{stt} toward AF (NiO). The antiferromagnetic magnon in AF (NiO) carries the spin current \mathcal{F}_{ex} toward F (FeNi).

Division of Biochemistry – Biofunctional Design-Chemistry –

http://www.scl.kyoto-u.ac.jp/~bfdc/index.html



Prof FUTAKI, Shiroh (D Pharm Sc)



Senior Lect IMANISHI, Miki (D Pharm Sc)



Assist Prof KAWANO, Kenichi (D Pharm Sc)

Students

MURAYAMA, Tomo (D3) TSUJI, Shogo (D3) AKISHIBA, Misao (D1) SHINODA, Kouki (D1) ARAFILES, Jan Vincent (M2) IINO, Masatomo (M2) ISE, Shoko (M2)
MASUDA, Toshihiro (M2)
SASAKI, Yuichiro (M2)
WATANABE, Kensuke (M2)
NOMURA, Kayo (M1)
SAKAGAMI, Koki (M1)

SAKAMOTO, Kentarou (M1) HAYASHI, Shuhei (UG) OGUSHI, Masaya (UG) TAMEMOTO, Naoki (UG)

Scope of Research

The ultimate goal of our research is the regulation of cellular functions using designed peptides and proteins. Current research subjects include 1) development of novel intracellular delivery systems aiming at elucidation and control of cellular functions using designed membrane-permeable peptide vectors, 2) elucidation of the DNA binding modes of zinc finger proteins and TALEs, and design of artificial transcription factors with various DNA binding specificities, 3) elucidation and control of membrane curvature, and 4) design of stimulation-responsible artificial peptides and proteins.



KEYWORDS

Membrane-Permeable Peptides Peptide Design Membrane Curvature Intracellular Delivery DNA/RNA Binding Protein

Selected Publications

Kawaguchi, Y.; Takeuchi, T.; Kuwata, K.; Chiba, J.; Hatanaka, Y.; Nakase, I.; Futaki, S., Syndecan-4 Is a Receptor for Clathrin-Mediated Endocytosis of Arginine-Rich Cell-Penetrating Peptides, *Bioconjug. Chem.*, 27, 1119-1130 (2016).

Murayama, T.; Pujals, S.; Hirose, H.; Nakase, I.; Futaki, S., Effect of Amino Acid Substitution in the Hydrophobic Face of Amphiphilic Peptides on Membrane Curvature and Perturbation: N-Terminal Helix Derived from Adenovirus Internal Protein VI as a Model, *Biopolymers*, **106**, 430-439 (2016).

Backlund, C. M.; Takeuchi, T.; Futaki, S.; Tew, G. N., Relating Structure and Internalization for ROMP-Based Protein Mimics, *Biochim. Biophys. Acta*, **1558**, 1443-1450 (2016).

Nakase, I.; Noguchi, K.; Fujii, I.; Futaki, S., Vectorization of Biomacromolecules into Cells Using Extracellular Vesicles with Enhanced Internalization Induced by Macropinocytosis, *Sci. Rep.*, **6**, 34397 (2016).

Azuma, Y.; Kükenshöner, T.; Ma, G.; Yasunaga, J.; Imanishi, M.; Tanaka, G.; Nakase, I.; Maruno, T.; Kobayashi, Y.; Arndt, K. M.; Matsuoka, M.; Futaki, S., Controlling Leucine-Zipper Partner Recognition in Cells through Modification of a-g Interactions, *Chem. Commun.*, **50**, 6364-6367 (2014).

Use of Calmodulin EF-Hand Peptides as Ca²⁺-Switchable Recognition Tags

Peptide motifs specifically recognizing each other are useful molecular tags to detect specific proteins and to control protein interactions. Calmodulin is a representative calcium-binding protein composed of four Ca2+-binding motifs with a helix-loop-helix structure called EF-hands, in which two helices pack together at an angle of approximately 90 degrees. The two helices are separated by a calcium-binding loop region. Here, we revealed the potential of peptide segments derived from the third and fourth EF-hands (EF3 and EF4) to act as recognition tags. An analysis of the disulfide formation mode among cysteines inserted at the N- or C-terminus of these peptide segments suggested that EF3 and EF4 peptides form a heterodimer with a topology similar to that in the wild-type protein. Heterodimer formation was shown to be a function of the Ca²⁺ concentration, indicating that these structures can be used as Ca²⁺ -switchable recognition tags. Using this EF-tag system, membrane fusion of liposomes decorated with EF3 and EF4 peptides was successfully induced. Because the EF-tags have different modes of mutual recognition and dimer formation from conventional peptide tags, this system would be used as an orthogonal recognition system to conventional ones.




Figure 1. (A) A possible structure of EF3 and EF4 dimers in the presence of Ca^{2+} . (B) Schematic representation of complex formation by EF3 with EF4 dependent on Ca^{2+} .

Identifying Membrane Proteins Responsible for Cellular Uptake of Octaarginine by Photocrosslinking

Arginine-rich cell-penetrating peptides (CPPs) have been widely used as carriers for intracellular delivery of bioactive molecules. In spite of increasing evidence for involvement of endocytosis in the cellular uptake of arginine-rich CPPs, the primary cell-surface receptors for these peptide carriers that would initiate endocytic processes have remained unknown. Our previous effort to identify membrane receptors for octa-arginine (R8) peptide using the photo-cross-linking probe bearing a photoreactive diazirine was unsuccessful due to substantial amounts of cellular proteins nonspecifically bound to the affinity beads. To address this issue, here a photo-cross-linking probe was developed, in which a cleavable linker of a diazobenzene moiety was employed to selectively elute cross-linked proteins by reducing agent-mediated cleavage. It was demonstrated that introduction of the diazobenzene moiety into the photoaffinity probe results in efficient purification of crosslinked proteins with remarkable reduction of nonspecific binding proteins, leading to successful identification of 17 membrane-associated proteins that would interact with R8 peptide. Among the proteins identified, syndecan-4, one of the heparan sulfate proteoglycans, was revealed to be an endogenous membrane-associated receptor for the cellular uptake of R8 peptide via clathrin-mediated endocytosis. It was also clarified that the intracellular delivery of bioactive proteins mediated by R8 peptide this syndecan-4-dependent pathway was also involved in. These results reveal that syndecan-4 is a primary cell-surface target for R8 peptide that allows intracellular delivery of bioactive cargo molecules via clathrin-mediated endocytosis.




Figure 2. Identification of cell-surface receptors bound to R8 using photo-cross-linking reaction.

Division of Biochemistry - Chemistry of Molecular Biocatalysts -

http://www.scl.kyoto-u.ac.jp/~bunta/index-j.html



Prof HIRATAKE, Jun (D Agr)



Assist Prof WATANABE, Bunta (D Agr)

Students


ITO, Shota (M2) SAITO, Shingo (M2) NAESHIRO, Kazuki (M2) LI, Xian (M2) MAKINO, Katsunori (M1)

Scope of Research

Life is the integration of numerous chemical interactions conducted by low molecular weight compounds and proteins. Our interest is understanding the mechanisms of these interactions from the viewpoint of organic chemistry. Our current research focuses on the following: a) Design and synthesis of a series of chemical probes in order to reveal not only ligand—protein and protein—protein interactions, but also behavior of low molecular weight compounds *per se*. b) Unraveling biosynthetic pathways of bioactive natural products at enzyme level. c) Application of our chemical probes in development of novel practical bioactive compounds.

KEYWORDS

Enzyme Inhibitors γ-Glutamyl Transpeptidase Fructosyl Peptide Oxidase Diabetes Diagnosis Protein–Protein Interaction Inhibitors



Selected Publications

Kamiyama, A.; Nakajima, M.; Han, L.; Wada, K.; Mizutani, M.; Tabuchi, Y.; Kojima-Yuasa, A.; Matsui-Yuasa, I.; Suzuki, H.; Fukuyama, K.; Watanabe, B.; Hiratake, J., Phosphonate-Based Irreversible Inhibitors of Human γ-Glutamyl Transpeptidase (GGT). GGsTop is a Non-Toxic and Highly Selective Inhibitor with Critical Electrostatic Interaction with an Active-Site Residue Lys562 for Enhanced Inhibitory Activity, *Bioorg. Med. Chem.*, **24**, 5340-5352 (2016).

Watanabe, B.; Minami, S.; Ishida, H.; Yoshioka, R.; Nakagawa, Y.; Morita, T.; Hayashi, K., Stereospecific Inhibitory Effects of CCG-1423 on the Cellular Events Mediated by Myocardin-Related Transcription Factor A, *PLOS ONE*, **10**, [e0136242-1]-[e0136242-16] (2015).

Watanabe, B.; Ichiyanagi, A.; Hirokawa, K.; Gomi, K.; Nakatsu, T.; Kato, H.; Kajiyama, N., Synthesis and Inhibitory Activity of Substrate-Analog Fructosyl Peptide Oxidase Inhibitors, *Bioorg. Med. Chem. Lett.*, **25**, 3910-3913 (2015).

Nakajima, M.; Watanabe, B.; Han, L.; Shimizu, B.; Wada, K.; Fukuyama, K.; Suzuki, H.; Hiratake, J., Glutathione-Analogous Peptidyl Phosphorus Esters as Mechanism-Based Inhibitors of γ-Glutamyl Transpeptidase for Probing Cysteinyl-Glycine Binding Site, *Bioorg. Med. Chem.*, **22**, 1176-1194 (2014).

Hayashi, K.; Watanabe, B.; Nakagawa, Y.; Minami, S.; Morita, T., RPEL Proteins are the Molecular Targets for CCG-1423, an Inhibitor of Rho Signaling, *PLOS ONE*, **9**, [e89016-1]-[e89016-12] (2014).

Determination of a Key Residue of γ-Glutamyl Transpeptidase for Substrate Recognition

γ-Glutamyl transpeptidase (GGT) plays a central role in homeostasis of antioxidant tripeptide glutathione, and has been implicated in a vast array of physiological disorders. In this study, we synthesized a series of mechanism-based GGT inhibitors to probe electrostatic interactions between the acceptor site residues of GGT and substrates. Our chemical, enzymological, and molecular biological approaches revealed that 3-hydroxyphenylacetic acid is an excellent mimic of the cysteinylglycine moiety of glutathione, and Lys562 of human GGT strongly recognizes their negative charge on the carboxy group (Figure 1). We demonstrated that this interaction considerably enhances the human GGT specificity of our inhibitor named GGsTop. GGsTop exhibited no inhibitory activity at 10 mM on a representative member of glutamine-dependent amidotransferases essential for a wide range of biosynthetic pathway, and showed no cytotoxicity toward human fibroblasts and hepatic stellate cells up to 1 mM.

Substrate-Analog Fructosyl Peptide Oxidase Inhibitors

Fructosyl peptide oxidase (FPOX) is widely used in the

area of diabetes diagnosis today. In this study, we designed and synthesized its substrate-analog inhibitors in order to unveil the substrate recognition mechanism of FPOX by X-ray diffraction analysis of enzyme-inhibitor co-crystals. Kinetic study revealed that our substrate analogs act as competitive inhibitors with K_i values ranging from 11.1 to 782 μ M (Figure 2). Co-crystallization of the enzyme with our inhibitors in order to determine the three-dimensional structure of FPOX is now in progress.

Molecular Mechanism of Myocardin-Related Transcription Factor A Inhibitors

Myocardin-related transcription factor A (MRTF-A) plays a pivotal role in epidermal-msenchymal transition. Inhibition of its nuclear transport is regarded as one of the attractive therapeutic targets since MRTF-A is closely associated with cancer and tissue fibrosis. In this study, we revealed that CCG-1423, originally developed as a Rho inhibitor, binds to the nuclear localization signal of MRTF-A and inhibits its nuclear transport mediated by importin-α/β1 (Figure 3). We also demonstrated that CCG-1423 inhibits a migration of melanoma cells triggered by MRTF-A activation, and the potency is affected by the stereochemistry of CCG-1423. The difference is elucidated by the binding manner of each stereoisomer to MRTF-A that speculated by a molecular modeling approach.




Figure 1. Proposed binding mode of GGsTop, glutathione, and acceptor substrate (Gly-Gly) to Lys562.

Figure 2. Chemical structures and inhibitory activity of FPOX inhibitors.




Figure 3. Molecular mechanism of MRTF-A inhibitor CCG-1423.

Division of Biochemistry – Molecular Biology –

http://seitaibunshijyoho-e.weebly.com



Prof AOYAMA, Takashi (D Sc)



Assoc Prof TSUGE, Tomohiko (D Sc)



Assist Prof FUJIWARA-KATO, Mariko (D Agr)



Techn Staff YASUDA, Keiko



PD FUJIWARA, Takashi (D Agr)

Students

GOTO, Kakeru (D3) ZHANG, Xiao-Juan (D2) SHIMAMURA, Ryota (D2) WATARI, Machiko (D1)

KURODA, Ryo (M2)

Guest Res Assoc

LIN, Xiaoya Peking University, China, P.R., 16 September–15 December

Scope of Research

This laboratory aims at clarifying molecular bases of regulatory mechanisms for plant development, especially plant morphogenesis, with techniques of forward and reverse genetics, molecular biology, and biochemistry. Current major subjects are: 1) phospholipid signaling in cell morphogenesis, 2) the transcriptional network for cytokinin responses,

3) COP9 signalosome modulating signal transduction in the nuclei, and 4) the endoreduplication cell cycle in cell differentiation.

KEYWORDS

Morphogenesis Phospholipid Signaling RNA Signal Transduction COP9 Signalosome



Selected Publications

Wu, Z.; Zhu, D.; Lin, X.; Miao, J.; Gu, L.; Deng, X.; Yang, Q.; Zhu, D.; Cao, X.; Tsuge, T.; Dean, C.; Aoyama, T.; Gu, H.; Qu, L.-J., RNA Binding Proteins RZ-1B and RZ-1C Play Critical Roles in Regulating Pre-mRNA Splicing and Gene Expression during Development in *Arabidopsis, Plant Cell*, 28, 55-73 (2016).

Lin, Q.; Ohashi, Y.; Kato, M.; Tsuge, T.; Gu, H.; Qu, L.-J.; Aoyama, T., GLABRA2 Directly Suppresses Basic Helix-loop-helix Transcription Factor Genes with Diverse Functions in Root Hair Development, *Plant Cell*, **27**, 2894-2906 (2015).

Wada, Y.; Kusano, H.; Tsuge, T.; Aoyama, T., Phosphatidylinositol Phosphate 5-kinase Genes Respond to Phosphate Deficiency for Root Hair Elongation in *Arabidopsis thaliana*, *Plant J.*, **81**, 426-437 (2015).

Hayashi, K.; Nakamura, S.; Fukunaga, S.; Nishimura, T.; Jenness, M. K.; Murphy, A. S.; Motose, H.; Nozaki, H.; Furutani, M.; Aoyama, T., Auxin Transport Sites are Visualized in Planta using Fluorescent Auxin Analogs, *Proc. Natl. Acad. Sci. USA*, **111**, 11557-11562 (2014). Kato, M.; Aoyama, T.; Maeshima, M., The Ca²+-binding Protein PCaP2 Located on the Plasma Membrane is Involved in Root Hair Development

as a Possible Signal Transducer, Plant J., 74, 690-700 (2013).

Genetic Analysis of RNA-Binding Protein Genes in *Arabidopsis thaliana*

Some classes of RNA binding proteins (RBPs) play pivotal roles in regulating gene expression, both co-transcriptionally and post-transcriptionally. The Arabidopsis thaliana genome encodes more than 600 putative RBPs, but the detailed functions of the vast majority of these RBPs remain unclear. Serine/arginine-rich (SR) proteins belong to a group of RBPs. Among the 19 SR proteins in Arabidopsis, only a few have been studied functionally. Gain-of-function analyses of SR protein genes RS2Z33 and SR30, as well as loss-of-function analyses of SR45, SCL33, SCL33a, RS40, and RS41, have revealed that the proteins encoded by these genes regulate alternative splicing of their own pre-mRNA and that of other SR genes. Glycine-rich RNA binding proteins (GRPs), a small group of plant RBPs, contain an N-terminal RRM domain and a C-terminal glycine-rich stretch. GRPs in Arabidopsis include GRP1-8 and RZ-1A-C (Fig 1). GRP7 and GRP8 are important regulators of circadian oscillations, flowering time, responses to plant pathogens, and cold stress. RZ-1A-C has a zinc finger motif between the RRM domain and C terminus. RZ-1, which was first reported in the tobacco, belongs to a subgroup of GRPs and binds to a large ribonucleoprotein particle localized in the nucleus. Homologs of RZ-1 in Arabidopsis were shown to have RNA chaperone activity in Escherichia coli. Overexpression of RZ-1A, but not homologous genes RZ-1B and RZ-1C, confers freezing tolerance in transgenic Arabidopsis. However, there is no direct genetic evidence to support the apparently important roles of these GRPs in plant growth and development and in responses to environmental stimuli.

In this study, we provide functional evidence that Arabidopsis RZ-1B and RZ-1C represent a unique group of GRPs. rz-1b rz-1c double mutants displayed a wide variety of defects, including delayed germination, retarded development of root and shoot meristems, late flowering, reduced stature, and serrated leaves (Fig 2). RZ-1B and RZ-1C interact with SR proteins through their C-terminal domains, which are also essential for nuclear speckle localization. High-throughput RNA-seq analysis of rz-1b rz-1c double mutants revealed perturbation of the splicing of many genes. Defective splicing was also observed in plants overexpressing the C-terminal domain of RZ-1C, confirming that RZ-1B/1C regulates splicing via interaction with SR proteins. RZ-1B and RZ-1C were found to be required for maintaining the optimal expression levels of more than 3000 genes, including many developmental regulators. Taken together, these findings reveal the essential roles of RZ-1B and RZ-1C as regulators of plant development, splicing, and general gene expression via interaction with




Figure 2. Pleotropic Defects of the rz-1b rz-1c Double Mutant. (A) Image of seedlings grown vertically for 10 d after germination. Bar = 0.5 cm. (B) Statistics of the primary root length from day 4 to day 11 after germination. Error bars represent SD (n = 20). (C) Confocal image of root meristems from different genotypes. The image shows the representative root meristems from seedlings at 4 d after germination. The roots were stained with propidium iodide. Asterisks indicate the dividing cells in the endodermis. Bar = 50 mm. (D) Statistics of endodermal cells from different genotypes within the meristematic regions. Error bars represent SD (n = 15). (E) rz-1b rz-1c generates multiple SAMs during vegetative growth. The images show wild-type and rz-1b rz-1c seedlings 3 weeks after germination. The lower panels show magnified images of the upper panels. Arrows indicate the positions of SAMs. Bars = 5 mm (upper panel) and 1 mm (lower panel). (F) Scanning electron microscopy images of the SAMs of the wild type and rz-1b rz-1c. Arrows indicate the SAM in the wild type and the corresponding region in rz-1b rz-1c. Bar = 50 mm. (G) Histochemical staining of CLAVATA3:GUS reporter in wild-type and rz-1b rz-1c backgrounds. Bar = 80 mm.

Division of Biochemistry - Chemical Biology -

http://www.scl.kyoto-u.ac.jp/~uesugi/e/index.php



Prof UESUGI, Motonari (D Pharm Sc)



Assoc Prof OHKANDA, Junko (D Eng)




Senior Lect PERRON, Amelie (Ph D)



Assist Prof TAKEMOTO, Yasushi (D Eng)



Proj Res* SATO, Shinichi (D Eng)



Proj Res**
KATSUDA, Yousuke
(D Sc)



Proj Res**
TAKASHIMA, Ippei
(D Pharm Sc)



Proj Res**
TAKENOYA, Mihoko
(D Eng)

**PD of iCeMS

Researchers

FURUTA, Tomoyuki****
OKUDA, Satoshi****

Proj Res

GOETZE, Sebastian Andreas (Ph D) ***
TANTIPANJAPORN, Ajcharapan***

Researchers(pt)

AKIZUKI, Maika QIN, Ying****

Assist Techn Staff

MAO, Di****
TAKAYA, Junichiro (D Med Sc) ****

*Assoc Prof of iCeMS

Assist Res Staff

NAKASHIMA, Mitsue****

***Guest Res Assoc of iCeMS

Students

ASANO, Risa (D3) LIAO, Youqi (D3) PUNZALAN, Louvy Lynn (D2) YATSUZUKA, Kenji (D2) MENDOZA, Aileen de-Leon (D1) JUNG, Yejin (D1) CHUNG, Watson Xie Khim (M2) KOTAKE, Kenjiro (M1) TAJIRI, Ken (M1) HAKARIYA, Hayase (M1) YOSHIDA, Hiroki (M1)
PE, Kathleen Beverly Alog (M1)
VU, Hue Thi (M1)
LE, Toan Khanh (RS)

Scope of Research

Chemical biology is an interdisciplinary field of study that is often defined as "chemistry-initiated biology." As biological processes all stem from chemical events, it should be possible to understand or manipulate biological events using chemistry. Our laboratory has been discovering or designing unique organic molecules that modulate fundamental processes in human cells. Such synthetic organic molecules often serve as tools for basic cell biology. Discovery or design of small molecules with unique biological activities permits small-molecule-initiated exploration of complex cellular events. Our mission is to create a new world of bioactive syn-

KEYWORDS

****iCeMS

Cell Therapy Chemical Biology Small Molecules Chemical Library Chemical Genetics

thetic molecules: new modes of activity, new shapes, and new sizes. We hope to open new avenues for small-molecule applications in a range of fields, including future concepts in drug discovery and use of small molecules for cell therapy.

Selected Publications

Katsuda, Y.; Sato, S.; Asano, L.; Morimura, Y.; Furuta, T.; Sugiyama, H.; Hagihara, M.; Uesugi, M., A Small Molecule That Represses Translation of G-quadruplex-Containing mRNA, *J. Am. Chem. Soc.*, **138**, 9037-9040 (2016).


Takaya, J.; Mio, K.; Shiraishi, T.; Kurokawa, T.; Otsuka, S.; Mori, Y.; Uesugi, M., A Potent and Site-Selective Agonist of TRPA1, *J. Am. Chem. Soc.*, 137, 15859-15864 (2015).

Parvatkar, P.; Kato, N.; Uesugi, M.; Sato, S.; Ohkanda, J., Intracellular Generation of a Diterpene-Peptide Conjugate that Inhibits 14-3-3-Mediated Interactions, *J. Am. Chem. Soc.*, **137**, 15624-15627 (2015).

Sato, S.; Watanabe, M.; Katsuda, Y.; Murata, A.; Wang, D. O.; Uesugi, M., Live-cell Imaging of Endogenous mRNAs with a Small Molecule, *Angew. Chem. Int. Ed.*, **54**, 1855-1858 (2015).


A Small Molecule That Represses Translation of G-Quadruplex-Containing mRNA


The G-quadruplexes form highly stable nucleic acid structures, which are implicated in various biological processes in both DNA and RNA. Although DNA G-quadruplexes have been studied in great detail, biological roles of RNA G-quadruplexes have received less attention. In this study, a screening of a chemical library permitted identification of a small-molecule tool that binds selectively to RNA G-quadruplex structures. The polyaromatic molecule, RGB-1, stabilizes RNA G-quadruplex, but not DNA versions or other RNA structures. RGB-1 intensified the G-quadruplex-mediated inhibition of RNA translation in mammalian cells, decreased expression of the NRAS proto-oncogene in breast cancer cells, and permitted identification of a novel sequence that forms G-quadruplex in NRAS mRNA. RGB-1 may serve as a unique tool for understanding cellular roles of RNA G-quadruplex structures.



A Potent and Site-Selective Agonist of TRPA1

TRPA1 is a member of the transient receptor potential (TRP) cation channel family that is expressed primarily on sensory neurons. This chemo-sensor is activated through covalent modification of multiple cysteine residues with a wide range of reactive compounds including allyl isothiocyanate (AITC), a spicy component of wasabi. The present study reports on potent and selective agonists of TRPA1, discovered through screening 1,657 electrophilic molecules. In an effort to validate the mode of action of hit molecules, we noted a new TRPA1-selective agonist, JT010 (molecule 1), which opens the TRPA1 channel by covalently and site-selectively binding to Cys621 (EC50 = 0.65 nM). The results suggest that a single modification of Cys621 is sufficient to open the TRPA1 channel. The TRPA1-selective probe described herein might be useful for further mechanistic studies of TRPA1 activation.





Division of Environmental Chemistry - Molecular Materials Chemistry -

http://molmat.kuicr.kyoto-u.ac.jp/index-e.html



KAJI, Hironori (D Eng)



Assist Prof FUKUSHIMA, Tatsuya (D Eng)

Students

KUBO, Shosei (D1)

OGATA, Isamu (M2)

MIURA, Tsutomu (M2)



Assist Prof SHIZU, Katsuyuki (D Eng)



Techn Staff OHMINE, Kyoko

Assist Techn Staff MIYAKE, Harumi



Techn Staff MAENO, Ayaka



PD SUZUKI, Katsuaki (D Human & Environtl. Studies)

Researcher NAKAGAWA, Hiromichi (D Eng)

WADA, Yoshimasa (M2) KIDO, Tsuyoshi (M1) HABUKA, Yume (M1)

TSUJI, Tomohiko (UG) HIGAKI, Sotaro (UG)

Scope of Research

Our research goal is to develop high-performance organic electroluminescence devices, organic solar cells, and polymer materials. Toward this, we carry out syntheses, device fabrications, precise structure characterizations, and quantum chemical calculations for high functional organic materials. Along with exploring

novel synthetic routes and novel devices, we perform detailed analyses of structures and dynamics, mainly by sophisticated solid-state NMR spectroscopy, in order to obtain structure-dynamics-property relationships.

KEYWORDS

Organic Light-Emitting Diodes Organic Solar Cells Solid-State NMR Quantum Chemical Calculation Amorphous Materials



Selected Publications

Uratani, H.; Kubo, S.; Shizu, K.; Suzuki, F.; Fukushima, T.; Kaji, H., Detailed Analysis of Charge Transport in Amorphous Organic Thin Layer by Multiscale Simulation without Any Adjustable Parameters, Sci. Rep., 6, 39128 (2016).

Shizu, K.; Kaji, H., Organic Electroluminescent Materials Realizing Efficient Conversion from Electricity to Light, J. Photopolym. Sci. Technol., 29, 305-310 (2016).

Nishimura, H.; Fukushima, T.; Wakamiya, A.; Murata, Y.; Kaji, H., The Influence of Quasiplanar Structures of Partially Oxygen-Bridged Triphenylamine Dimers on the Properties of Their Bulk Films, Bull. Chem. Soc. Jpn., 89, 726-732 (2016).

Hayase, G.; Nonomura, K.; Kanamori, K.; Maeno, A.; Kaji, H.; Nakanishi, K., Boehmite Nanofiber-Polymethylsilsesquioxane Core-Shell Porous Monoliths for a Thermal Insulator under Low Vacuum Conditions, Chem. Mater., 28, 3237-3240 (2016).

Wada, Y.; Shizu, K.; Kubo, S.; Fukushima, T.; Miwa, T.; Tanaka, H.; Adachi, C.; Kaji, H., Highly Efficient Solution-Processed Host-Free Organic Light-Emitting Diodes Showing an External Quantum Efficiency of Nearly 18% with a Thermally Activated Delayed Fluorescence Emitter, Appl. Phys. Express, 9, 032102-1-032102-3 (2016).

Hayase, G.; Kanamori, K.; Maeno, A.; Kaji, H.; Nakanishi, K., Dynamic Spring-Back Behavior in Evaporative Drying of Polymethylsilsesquioxane Monolithic Gels for Low-Density Transparent Thermal Superinsulators, J. Non-Cryst. Solids, 434, 115-119 (2016).

Highly Efficient Solution-processed Host-free Organic Light-emitting Diodes using a Thermally Activated Delayed Fluorescence Emitter

Organic light-emitting diodes (OLEDs) have attracted great interest as a promising technique for developing solid-state lighting source and flexible flat-panel displays. To date, vacuum vapor deposition has been widely used to fabricate OLEDs. However, it has drawbacks, including high production cost and difficulty in fabricating multi-dopant OLEDs. Solution processing is an alternative method to vacuum vapor deposition for fabricating OLEDs. It has an advantage of lowering the production cost and realizing large-area high-resolution displays. An emitting layer of an OLED is generally a doped film composed of an emitting dopant and a host material. To realize a high-performance OLED, the host material is required to have high excited-state energies, ambipolar transport properties, and suitable HOMO and LUMO energy levels. Host-free OLEDs contain a neat film as an emissive layer (that is,




Figure 1. (a) Energy diagram of solution-processed host-free OLEDs containing DMAC-TRZ as an emitter. (b) Luminance–EQE characteristics of the solution-processed host-free OLEDs.

the emissive layer consists of an emitting material alone), which can remove the limitations of host materials. In this study, using a spin-coated neat film of a thermally activated delayed fluorescence emitter (DMAC-TRZ) as an emissive layer, we fabricated a solution-processed OLED. The OLED exhibited a maximum external quantum efficiency (EQE) of 17.6% (Figure 1), which is the highest value obtained for solution-processed host-free OLEDs, including phosphorescent systems. By further optimizing the device structure, we achieved an EQE of 16.8% at high luminance.

Charge Transport Simulations in Amorphous Thin Films for OLEDs

A fundamental understanding of charge transport properties of organic thin films is essential to improve OLED performance. Charge transport properties of organic molecules have been discussed in terms of their HOMO and LUMO energy levels and electronic couplings between molecules. The HOMO and LUMO can be calculated using quantum chemical calculations for isolated molecules, while the electronic couplings are considered to depend largely on an aggregated structure. We performed multiscale charge transport simulations for amorphous structures of N, N'-dicarbazole-3,5-benzene (mCP) and 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), which have been used as a host material for OLEDs. By explicitly considering organic molecules, we investigated the contribution of respective molecular pairs to the charge transport in amorphous thin films. The molecular-level analysis of the charge transport simulations showed that molecular pairs with large electronic couplings are not most important in charge transport processes. Charges were found to be transported effectively in the forward direction via other molecular pairs that do not have substantially large electronic couplings.




Figure 2. Image of charge transport in amorphous mCP films.

Division of Environmental Chemistry

- Hydrospheric Environment Analytical Chemistry -

http://inter3.kuicr.kyoto-u.ac.jp/scope_E.html



Prof SOHRIN, Yoshiki (D Sc)



Assoc Prof UMETANI, Shigeo (D Sc)



Assist Prof TAKANO, Shotaro (D Sc)



Techn Staff MINAMI, Tomoharu (D Eng)

Students

ZHENG, Linjie (D2) TSUJISAKA, Makoto (M2) UEHARA, Wataru (M2) OKITA, Shutaro (M1)

OTSUKA, Yuhei (M1)

Scope of Research

(i) Biogeochemistry of trace elements in the hydrosphere: Novel analytical methods are developed for trace metals and their isotopes. Distribution of trace elements in the hydrosphere and their effects on the ecosystem are investigated. The study also covers hydrothermal activity, deep biosphere, and paleocean.


(ii) Ion recognition: Novel ligands and ion recognition system are designed, synthesized, and characterized.





Marine Chemistry Analytical Chemistry
Trace Elements Stable Isotopes

Metal Ion Recognition



Selected Publications

Nakashima, Y.; Shimizu, A.; Maruo, M.; Sohrin, Y., Trace Elements Influenced by Environmental Changes in Lake Biwa: (I) Seasonal Variations under Suboxic Hypolimnion Conditions during 2007 and 2009, *Limnol.*, **17**, 151-162 (2016).

Sohrin, Y.; Nakashima, Y.; Maruo, M., Trace Elements Influenced by Environmental Changes in Lake Biwa: (II) Chemical Variations in the Hypolimnion over the Last Half-century, *Limnol.*, **17**, 163-173 (2016).

Minami, T.; Konagaya, W.; Zheng, L.; Takano, S.; Sasaki, M.; Murata, R.; Nakaguchi, Y.; Sohrin, Y., An Off-line Automated Preconcentration System with Ethylenediaminetriacetate Chelating Resin for the Determination of Trace Metals in Seawater by High-resolution Inductively Coupled Plasma Mass Spectrometry, *Anal. Chim. Acta*, **854**, 183-190 (2015).

Takano, S.; Tanimizu, M.; Hirata, T.; Sohrin, Y., Isotopic Constraints on Biogeochemical Cycling of Copper in the Ocean, *Nat. Commun.*, **5**, [5663-1]-[5663-7] (2015).

Vu, H. T. D.; Sohrin, Y., Diverse Stoichiometry of Dissolved Trace Metals in the Indian Ocean, Sci. Rep., 3, [1745-1]-[1745-5] (2013).

Firdaus, M. L.; Minami, T.; Norisuye, K.; Sohrin, Y., Strong Elemental Fractionation of Zr-Hf and Nb-Ta across the Pacific Ocean, *Nature Geosci.*, **4**, 227-230 (2011).

Stoichiometry of Al, Mn, Fe, Co, Ni, Cu, Zn, Cd, and Pb in Seawater around the Juan de Fuca Ridge

We have studied the distributions of Al, Mn, Fe, Co, Ni, Cu, Zn, Cd, and Pb around the Juan de Fuca Ridge (JdFR) in total dissolvable (td), dissolved (d), and labile particulate (lp) fractions, which represent a fraction in unfiltered samples, filtered samples through an AcroPak capsule filter, and the difference between td and d, respectively. Al and Fe were dominated by lp-species, while Ni, Zn, and Cd were dominated by d-species with undetectable amounts of lp-species. Major findings in this study are as follows: (i) The continental margin (CM) provided large sources of Al, Mn, Fe, and Co from the surface to ~2000 m in depth. The supply from CM caused high surface concentrations of dMn and dCo, a subsurface (100-300 m depth) maximum of dCo, and intermediate (500-2000 m depth) maxima of lpAl and lpFe. The supply of dFe from CM was ~10 times that from the high-temperature hydrothermal activity at station BD21, which is located at ~3 km from the Middle Valley venting site and ~200 km from Vancouver Island. (ii) DPb was maximum at the top layer of North Pacific Intermediate Water, probably owing to isopycnal transport of anthropogenic Pb via advection of subducted surface waters. Although dCo and dPb had different sources in the upper water, they showed a strong linearity below 300 m ($r^2 = 0.95$, n = 38), indicating concurrent scavenging. (iii) A high-temperature hydrothermal plume occurred at a depth of 2300 m at BD21, accounting for maxima of dAl, dMn, dFe, lpCu, and lpPb and a minimum of dCu. (iv) Strong bottom maxima of lpAl, lpMn, lpFe, lpCo, and lpPb occurred above the abyssal plain at the western foot of the JdFR, indicating resuspension of sediments. However, bottom maxima of d-species were apparent only for dAl and dCu.

A Simple and Rapid Method for Isotopic Analysis of Nickel, Copper, and Zinc in Seawater Using Chelating Extraction and Anion Exchange

Stable isotope ratios of Ni, Cu, and Zn are powerful tools to elucidate biogeochemical cycling of the trace metals in the ocean. However, analytical difficulties have retarded development of isotopic study of these metals. Especially, isotopic analysis of seawater Ni is troublesome, because Ni is not retained on anion exchange resin and passed through the column with seawater matrix. We have developed a simple and rapid method for simultaneous analyses of Ni, Cu, and Zn isotope ratio in seawater using NOBIAS Chelate-PA1 resin and anion exchange resin. A NOBIAS Chelate-PA1 resin column quantitatively collected Ni, Cu, and Zn from seawater and effectively removed seawater matrix. Subsequent anion exchange purified Ni, Cu, and Zn from each other. The blanks of this method (0.22 ng for Ni, 0.29 ng for Cu, and 0.53 ng for Zn) were sufficiently low to determine isotope ratios of Ni, Cu, and Zn in surface seawater. Using this method, we analyzed GEOTRACES reference seawater samples, i.e., SAFe D1 and SAFe D2, and seawater samples collected from some depths in the subarctic North Pacific. The results were consistent with reported values.



Figure 1. Members of the Hakuho-Maru KH-12-4 cruise, during which samples for this work were collected.




Figure 2. Outline of the analytical method.

Takano et al., (in preparation)

Division of Environmental Chemistry- Solution and Interface Chemistry -

http://www.scl.kyoto-u.ac.jp/~yoeki/



Prof HASEGAWA, Takeshi (Ph D)



Assist Prof WAKAI, Chihiro (D Sc)



Assist Prof SHIMOAKA, Takafumi (D Sc)

KISE, Ryuma (M1)

Lect (pt)

OSAWA, Masatoshi (D Eng) Hokkaido University

Students

SHIOYA, Nobutaka (D2) UKAI, Hironori (M2)

IZUMI, Naoki (M2) HADA, Miyako (M2)

Scope of Research

To understand the chemical structure and properties of a molecular aggregated system, the keywords of molecular interactions and orientation are focused on, and the molecular aggregates are investigated by using originally developed spectroscopic techniques. The current major topics are: (1) perfluoroalkyl-specific properties in a condensed system and a single molecule; (2) controlling factors of a molecular packing and orientation in a thin film of an organic semiconductor compound; (3) the optimal experimental condition and the correction factor of pMAIRS based on electrodynamics for estimating an accurate quantitative molecular orientation in a thin film having a surface roughness.

KEYWORDS

Infrared and Raman Spectroscopy Perfluoroalkyl Compounds pMAIRS Surface and Interface Chemistry Organic Semiconductor



Selected Publications

Wakai, C.; Shimoaka, T.; Hasegawa, T., Characterization of Adsorbed Molecular Water on the Surface of a Stretched Polytetrafluoroethylene Tape Analyzed by ¹H NMR, *J. Phys. Chem. B*, **120**, 2538-2543 (2016).

Shimoaka, T.; Tanaka, Y.; Shioya, N.; Morita, K.; Sonoyama, M.; Amii, H.; Takagi, T.; Kanamori, T.; Hasegawa, T., Surface Properties of a Single Perfluoroalkyl Group on Water Surfaces Studied by Surface Potential Measurements, *J. Colloid Interf. Sci.*, **483**, 353-359 (2016).

Hada, M.; Shioya, N.; Shimoaka, T.; Eda, K.; Hada, M.; Hasegawa, T., Comprehensive Understanding of Structure-Controlling Factors of a Zinc Tetraphenylporphyrin Thin Film Using pMAIRS and GIXD Techniques, *Chem. Eur. J.*, **22**, 16539-16546 (2016).

Shimoaka, T.; Hasegawa, T., Molecular Structural Analysis of Hydrated Ethylene Glycol Accounting for the Antifreeze Effect by Using Infrared Attenuated Total Reflection Spectroscopy, *J. Mol. Liquids*, **223**, 621-627 (2016).

Iwasa, J.; Kumazawa, K.; Aoyama, K.; Suzuki, H.; Norimoto, S.; Shimoaka, T.; Hasegawa, T., In Situ Observation of a Self-Assembled Monolayer Formation of Octadecyltrimethoxysilane on a Silicon Oxide Surface Using a High-Speed Atomic Force Microscope, *J. Phys. Chem. C*, **120**, 2807-2813 (2016).

Characterization of Adsorbed Molecular Water on the Surface of a Stretched Polytetrafluoroethylene Tape Analyzed by ¹H NMR

A single molecule often exhibits a largely different material character from a bulk matter. Although a perfluoroalkyl (Rf) compound is a representative one, many interests have mostly been devoted to the bulk character only thus far, leaving the single molecular character unclear. Recently, a new theoretical framework, stratified dipole-arrays (SDA) theory, has appeared for comprehensive understanding of Rf compounds in terms of both single and bulk systems. On this theory, a mechanically stretched polytetrafluoroethylene (PTFE) is expected to exhibit a single-molecular character having dipole-driven properties, which should attract molecular water. In the present study, a stretched PTFE tape is revealed to attract molecular water (not water droplet) in fact, and the adsorbed water molecules are highly restricted in motion by the dipole-dipole interaction studied by using ¹H NMR, which agrees with the prediction by the SDA theory.




Figure 1. ¹H NMR spectra of strongly adsorbed molecular water on a stretched PTFE tape.

Surface Properties of a Single Perfluoroalkyl Group on Water Surfaces Studied by Surface Potential Measurements

A discriminative study of a single perfluoroalkyl (Rf) group from a bulk material is recently recognized to be necessary toward the total understanding of Rf compounds based on a primary chemical structure. The single molecule and the bulk matter have an interrelationship via an intrinsic two-dimensional (2D) aggregation property of an Rf group, which is theorized by the stratified dipole-arrays (SDA) theory. Since an Rf group has dipole moments along many C–F bonds, a single Rf group would possess a hydrophilic-like character on the surface. To reveal the hydration character of a single Rf group, in the present study, surface potential (ΔV) measurements are performed for Langmuir monolayers of Rf-containing compounds. From a comparative study with a monolayer of a normal hydrocarbon compound, the hydration/dehydration dynamics of a lying

Rf group on water has first been monitored by ΔV measurements, through which a single Rf group has been revealed to have a unique "dipole-interactive" character, which enables the Rf group interacted with the water 'surface.' In addition, the SDA theory proves to be useful to predict the 2D aggregation property across the phase transition temperature of 19 °C by use of the ΔV measurements.




Figure 2. Schematic image of a dipole-interactive Rf group on the water surface and a fairly large negative surface potential.

Comprehensive Understanding of Structure-Controlling Factors of a Zinc Tetraphenylporphyrin Thin Film Using pMAIRS and GIXD Techniques

The performance of an organic electronic device is significantly influenced by the anisotropic molecular structure in the film, which has long been difficult to predict especially for a solution process. In the present study, a zinc tetraphenylporphyrin (ZnTPP) thin film prepared by a solution process was chosen to comprehensively explore the molecular-arrangement mechanism as a function of representative film-preparation parameters: solvent, film-preparation technique, and thermal annealing. The anisotropic structure was first analyzed by using a combination of infrared p-polarized multiple-angle incidence resolution spectrometry (pMAIRS) and grazing incidence X-ray diffraction (GIXD), which readily revealed the molecular orientation and crystal structure, respectively. As a result, the real dominant factor was found to be the evaporation time of the solvent that determines the initial two different molecular arrangements, types-I and -II, while the thermal annealing was found to play an additional role of improving the molecular order. The correlation between the molecular orientation and the crystal structure was also revealed through the individual orientation analysis of the porphyrin and phenyl rings.




Figure 3. pMAIRS spectra of a ZnTPP thin film and a schematic image of the two conjugated rings on the surface.

Division of Environmental Chemistry- Molecular Microbial Science -

http://www.scl.kyoto-u.ac.jp/~mmsicr/mmstojp/Top_en.html



Prof KURIHARA, Tatsuo (D Eng)



Assist Prof KAWAMOTO, Jun (D Agr)



Assist Prof OGAWA, Takuya (D Agr)

Assist Res Staff KITAYAMA, Kaori

Students

SUGIURA, Miwa (D3) KAWAI, Soichiro (D3) OHKE, Yoshie (D3) CHEN, Chen (D2) TOKUNAGA, Tomohisa (D2) TOYOTAKE, Yosuke (D1) OTA, Masaki (M2) KUMAGAI, Fumihito (M2) SUGIMOTO, Saki (M2) FURUKAWA, Takahiro (M2) YOKOYAMA, Fumiaki (M2) CHEN, Tsai-Min (M2) KIMURA, Syunsaku (M1) TANAKA, Asako (M1) HATSUHARA, Hikari (M1) YAGURA, Kazuki (M1) YAMAGUCHI, Toshiaki (M1) SUWANAWAT, Nittikarn (M1)

Scope of Research

Microorganisms are found almost everywhere on Earth. They have great diversity of capacities to adapt to various environments, including chemically and physically unusual environments. Our main subject is to clarify the molecular basis of environmental adaptations of microorganisms and their application. Specific functions of proteins and lipids with essential roles in environmental adaptation of extremophilic microorganisms are of our particular interest. We also undertake mechanistic analysis of microbial enzymes, in particular those involved in unique metabolic pathways, and their application.

KEYWORDS

Extremophiles
Bacterial Cold-adaptation Mechanism
Polyunsaturated Fatty Acid
Phospholipid Acyltransferase
Membrane Vesicle



Selected Publications

Ito, T.; Gong, C.; Kawamoto, J.; Kurihara, T., Development of a Versatile Method for Targeted Gene Deletion and Insertion by Using the *pyrF* Gene in the Psychrotrophic Bacterium, *Shwanella livingstonensis* Ac10, *J. Biosci. Bioeng.*, **122**, 645-651 (2016).

Ohke, Y.; Maruyama, S.; Tarui, J.; Wang, Y.; Kawamoto, J.; Kurihara, T., A Phosphoporin Homolog Plays a Critical Role in the Dissimilatory Iron-Respiration Linked to Iron (III) Reduction by a Cold-adapted Bacterium, *Shewanella livingstonensis* Ac10, *Trace Nutr. Res.*, **33**, 35-42 (2016)

Sugiura, M.; Park, J.; Kawamoto, J.; Esaki, N.; Kurihara, T., Regulatory Mechanism of Membrane Protein Production in an EPA-Producing Bacterium, *Shewanella livingstonensis* Ac10, *Trace Nutr. Res.*, **33**, 63-72 (2016).

Purification and Enzymatic Characterization of Bacterial 1-Acyl-sn-glycerol-3-phosphate Acyltransferase PlsC

A cold-adapted bacterium Shewanella livingstonensis Ac10 produces phospholipids esterified at the sn-2 position with eicosapentaenoic acid (EPA), which are important biomembrane components to survive in a cold environment. It is known that, in a bacterial de novo phospholipid synthesis, acylation at the sn-2 position is catalyzed by 1-acyl-sn-glycerol-3-phosphate acyltransferase PlsC. We previously carried out gene deletion experiments to find that the microbe has five PlsC homologs (PlsC1-5), among which PlsC1 is exclusively responsible for the production of the EPA-containing phospholipids. To gain further insights into the enzymatic properties of PlsC1, we attempted to purify and characterize the enzyme. After careful investigation of purification conditions, we finally succeeded in purifying PlsC1 in an active form and revealed its enzymatic features such as optimal reaction conditions and substrate specificity. It is notable that PlsC1 showed a higher activity toward unsaturated fatty acids including EPA than saturated ones. These findings will enable us to investigate a protein interaction network underlying the cold adaptation of S. livingstonensis Ac10 and to analyze as yet unidentified reaction mechanism of PlsC.

Characterization of Membrane Vesicles Produced by *Shewanella* Species

It is known that most bacteria secrete membrane vesicles (MVs), which are composed of lipid membrane and contain nucleic acids, proteins and periplasmic solutes, to outer milieu. Though much attention has been paid to their physiological roles as well as the application to biotechnology, the molecular mechanism for MV secretion remains largely unknown. To better understand the mechanism, we characterized MVs from two cold-adapted Shewanella species. By using cryo-electron microscopy techniques, an EPAproducing bacterium S. livingstonensis Ac10 was shown to secrete MVs that mainly have a spherical single-bilayered structure with diameter of approximately 100 nm. Fatty acid profiling showed that specific fatty acids, e.g. hexadecanoic acid, are predominantly loaded onto MVs. Moreover, it was notable that the deletion of EPA biosynthetic genes altered the size and productivity of MVs. On the other hand, we found that Shewanella sp. HM13 secretes a much larger amount of MVs than S. livingstonensis Ac10. The MVs from this novel strain were found to include a single major protein. These features will be suitable for developing a new low-temperature protein production system where a protein of interest is concentrated in secreted MVs.




Figure 1. Schematic view of PlsC1-catalyzed reaction. PlsC1 showed a higher activity toward unsaturated fatty acids including EPA (20:5) and hexadecenoic acid (16:1) than saturated ones such as hexadecanoic acid (16:0). G3P, *sn*-glycerol 3-phosphate; AGP, 1-acyl-*sn*-glycerol 3-phosphate; PA, phosphatidic acid; GPAT, G3P acyltransferase.




Figure 2. Electron microscopic image of MVs from *Shewanella* sp. HM13. The bar indicates 100 nm.

Division of Multidisciplinary ChemistryPolymer Materials Science

http://www.scl.kyoto-u.ac.jp/~polymat/index.html



Prof TAKENAKA, Mikihito (D Eng)



Assoc Prof NISHIDA, Koji (D Eng)



Assist Prof OGAWA, Hiroki (D Eng)

Students

HIGASHINO, Naruhisa (M2) MIYATAKE, Yuki (M2) NAGANO, Yoshiki (M2) OKAMOTO, Masanori (M1) FUJITA, Shuuhei (M1) TSUJIOKA, Kouta (M1) KUROSAKI, Satoshi (M1) OMOSU, Takuya (UG) MATSUNAGA, Tsubasa (UG) FUWA, Takuto (UG) NISHIMURA, Yu (UG)

Scope of Research

The structure and molecular motion of polymer substances are studied, mainly using scattering methods such as X-ray, neutron, and light with intent to solve fundamentally important problems in polymer science. The main project are studied on 1) the morphologies and the dynamics of self-assembling processes in block copolymers, 2) the hierarchical structures in crystalline polymers, 3) the viscoelastic effects in glassy materials, 4) formation pro-

cesses and structure of polymer gels, and 5) the structure ad molecular motion of polyelectrolyte solutions.

KEYWORDS

Polymer Physics Polymer Properties Self Assembly Softmatter Hierarchical Structure



Selected Publications

Wang, Y. C.; Kim, M. I.; Akasaka, S.; Saijo, K.; Hasegawa, H.; Hikima, T.; Takenaka, M., Fddd Structure in Polystyrene-block-polyisoprene Diblock Copolymer/Polystyrene Homopolymer Blends, *Macromolecules*, **49** (6), 2257-2261 (2016).

Matsumoto, K.; Terashima, T.; Sugita, T.; Takenaka, M.; Sawamoto, M., Amphiphilic Random Copolymers with Hydrophobic/Hydrogen-Bonding Urea Pendants: Self-Folding Polymers in Aqueous and Organic Media, *Macromolecules*, **49** (**20**), 7917-7927 (2016).

Ogawa, H.; Takenaka, M.; Miyazaki, T.; Fujiwara, A.; Lee, B.; Shimokita, K.; Nishibori, E.; Takata, M., Direct Observation on Spin-Coating Process of PS-*b*-P2VP Thin Films, *Macromolecules*, **49**, 3471-3477 (2016).

Ishige, R.; Ohta, N.; Ogawa, H.; Tokita, M.; Takahara, A., Fully Liquid-Crystalline ABA Triblock Copolymer of Fluorinated Side-Chain Liquid-Crystalline A Block and Main-Chain Liquid-Crystalline, B Block: Higher Order Structure in Bulk and Thin Film States, *Macromolecules*, **49**, 6061-6074 (2016).

Formation of OBDD Structure in PS-PI-PDMS

Ordered bicontinuous double-diamond morphology (OBDD) was formed in cast film of polystyrene-block-polyisoprene-block-polydimethylsiloxiane (SID) triblock copolymer from toluene solution. The following two step phase separation was found to occurs during solvent cast: In the early stage of the cast process, hexagonally-packed polydimethylsiloxiane (PDMS) cylinders were self-assembled in polystyrene (PS)/polyisoprene (PI) matrix order-order transition, then order-order transition was induced by microphase separation between PS and PI association with the increase of concentration.




Figure 1. Formation of OBDD structure in PS-PI-PDMS

Quasi-Equilibrium Gelation Temperature of Aqueous Methylcellulose

Aqueous methylcellulose (MC) forms hydrogel upon heating and the gel dissolves upon cooling i.e. thermo-reversible gelation. However, the values of the gelation temperature differ depending on the sources of MC samples and on the methods of determination as well. In this study, we have obtained a lower limit of the gelation temperature for a specific MC below which aqueous MC does not form the gel, by dually extrapolating the holding time for gelation to infinite and the mechanical probing force to zero. The gelation temperature obtained in this manner decreased with increasing concentration of the MC; to cite a case, it was located at ca. 21 °C for 10 wt%, which is ca. 10 °C lower than a conditional value estimated after holding 1 h without these dual extrapolations.

Direct Observation on Spin-coating Process of PS-b-P2VP Thin Films

During the spin-coating process of symmetric poly(sty-rene-*b*-2-vinylpyridine) (PS-*b*-P2VP) block copolymers, after the formation of the micelles in dilute solution, the




Figure 2. Lower limit of gelation temperature (diamond marks), lower limit of cloud point (open-circle marks) and conditional gelation temperature estimated by employing a stainless-steel ball after holding 1 h (cross marks). Lines are guide for eyes.

selective solvent induced two kinds of the morphological transition. Firstly, the disordered spherical micelles were transformed into a BCC lattice of spheres of which the (110) plane was oriented perpendicularly to the substrate surface. Secondly, further evaporation induced a transition from spheres on the BCC lattice into cylindrical structures. The orientation of the cylinders perpendicular to the substrate surface was induced by solvent convection perpendicular to the substrate, which occurs during rapid solvent evaporation. After this transition, vitrification of PS and P2VP prevented any further transition from cylinders to the more thermodynamically stable lamellar structures, as are generally observed as the bulk equilibrium state.




Figure 3. Spin-coating process of PS-P2VP thin film.

Division of Multidisciplinary Chemistry – Molecular Rheology –

http://rheology.minority.jp/en/



Prof WATANABE, Hiroshi (D Sc)



Assoc Prof MATSUMIYA, Yumi (D Eng)



Program-Specific Assist Prof DOI, Yuya (D Eng)

Students

ISHIBASHI, Takuma (M2) KUBO, Kosei (M2) OHKITA, Yuya (M2) NISHINO, Takuma (M1) ISHIMOTO, Ryushin (UG) MOTOMURA, Yoshihiro (UG) SATO, Ohji (UG)

Guest Res Assoc


RAKKAPAO, Natthida (Ph D) KWON, Oh Min Prince of Songkla University, Thailand, 11 July–13 August Seoul National University, Korea, R., 14 April–31 August

Scope of Research

Research focus is placed on the molecular origin of rheological properties of various materials. Depending on time and temperature, homogeneous polymeric materials exhibit typical features of glass, rubber, and viscous fluid while heterogeneous polymeric systems exhibit plasticity in addition to these features. For basic understanding of these features, the molecular motion and structures at various scales are studied for polymeric systems in deformed state. Rheological measurements are performed with various rheometers. Auto-correlation of the molecular orientation is also investigated with dynamic dielectric spectroscopy. Analysis of rheological and dielectric behavior elucidates new aspect of softmatter physics.

KEYWORDS

Rheology Dielectric Spectroscopy Softmatter



Selected Publications

Ankiewicz, S.; Orbey, H.; Watanabe, H.; Lentzakis, H.; Dealy, J., On the Use of Continuous Relaxation Spectra to Characterize Model Polymers, *J. Rheol.*, **60**, 1115-1120 (2016).

Matsumiya, Y.; Watanabe, H.; Urakawa, O.; Inoue, T., Experimental Test for Viscoelastic Relaxation of Polyisoprene Undergoing Monofunctional Head-to-Head Association and Dissociation, *Macromolecules*, **49(18)**, 7088-7095 (2016).


Matsumiya, Y.; Watanabe, H., Nonlinear Stress Relaxation of Miscible Polyisoprene/Poly(p-tert-butylstyrene) Blends in Pseudomonodisperse State, *Macromolecules*, **49(12)**, 4544-4556 (2016).

Kwon, Y.; Matsumiya, Y.; Watanabe, H., Viscoelastic and Orientational Relaxation of Linear and Ring Rouse Chains Undergoing Reversible End-Association and Dissociation, *Macromolecules*, **49(9)**, 3593-3607 (2016).

Doi, Y.; Iwasa, Y.; Watanabe, K.; Nakamura, M.; Takano, A.; Takahashi, Y.; Matsushita, Y., Synthesis and Characterization of Comb-Shaped Ring Polystyrenes, *Macromolecules*, **49**, 3109-3115 (2016).


Experimental Test for Viscoelastic Relaxation of Polyisoprene Undergoing Monofunctional Head-to-Head Association and Dissociation

A viscoelastic test was made for end-carboxylated polyisoprene (PI-COOH) of the molecular weight $M = 30.5 \times 10^{-5}$ 103 that underwent the interchain association and dissociation through hydrogen bonding of the COOH groups at the chain end. As a reference, the test was made also for neat PI unimer (with no COOH group at the chain end) and for PI₂ dimer (with $M = 61.0 \times 10^3$), the latter being synthesized through end-coupling of PI- anions (precursor of the PI-COOH sample). The PI-COOH, neat unimer, and dimer samples were diluted in oligomeric butadiene (oB) to a concentration of 10 wt %. The neat unimer and dimer exhibited nonentangled Rouse behavior at this concentration, as expected from their molecular weights. At low temperatures ($T \le 0$ °C) the PI-COOH sample relaxed slower than the reference unimer but faster than the dimer, whereas the relaxation of PI-COOH approached that of the unimer with increasing T > 0 °C, and this change of the relaxation time of PI-COOH was associated with changes in the angular frequency (\omega) dependence of the dynamic modulus. This behavior of PI-COOH was well described by a recently proposed theory considering motional coupling between the end-associating unimer and its dimer at chemical equilibrium. On the basis of this result, an effect of the polymeric character of PI-COOH chain on the viscoelastically detected association/dissociation of the hydrogen bonding of the COOH groups was discussed.



Viscoelastic and Orientational Relaxation of Linear and Ring Rouse Chains Undergoing Reversible End-Association and Dissociation

For dilute telechelic linear and ring Rouse chains undergoing reversible end-association and dissociation, the time (t) evolution equation was analytically formulated for the bond vector of the subchain (or segment), $\mathbf{u}^{[c]}(n,t)$ with n being the subchain index and the superscript c specifying the chain (c = L and R for the linear and ring chains). The end-association of the linear chain (i.e., ring formation) occurs only when the ends of the linear chain come into close proximity. Because of this constraint for the ring formation, the time evolution equation for $\mathbf{u}^{[L]}(n,t)$ of the linear chain was formulated with a conceptually new, twostep expansion method: $\mathbf{u}^{[L]}(n,t)$ was first expanded with respect to its sinusoidal Rouse eigenfunction, $\sin(p\pi n/N)$ with p = integer and N being the number of subchains per chain, and then the series of odd sine modes is re-expanded with respect to cosine eigenfunctions of the ring chain, $\cos(2\alpha\pi n/N)$ with α = integer, so as to account for that constraint. This formulation allowed analytical calculation of the orientational correlation function, $S^{[c]}(n,m,t) = b^{-2} \langle u_x^{[c]} \rangle$ $(n,t)u_{\nu}^{[c]}(m,t)$ (c = L, R) with b being the subchain step length, and the viscoelastic relaxation function, $g^{[c]}(t) \propto$ $\int_0^N S^{[c]}(n,n,t) dn$. It turned out that the terminal relaxation of $g^{[R]}(t)$ and $g^{[L]}(t)$ of the ring and linear chains is retarded and accelerated, respectively, due to the motional coupling of those chains occurring through the reaction. This coupling breaks the ring symmetry (equivalence of all subchains of the ring chain in the absence of reaction), thereby leading to oscillation of the orientational anisotropy $S^{[R]}(n,n,t)$ of the ring chain at long t with the subchain index n. The coupling also reduces a difference of the anisotropy $S^{[L]}(n,n,t)$ of the linear chain at the middle $(n \sim N/2)$ and end $(n \sim 0)$.



Division of Multidisciplinary Chemistry – Molecular Aggregation Analysis –

http://www.em2.kuicr.kyoto-u.ac.jp/index.html



Assist Prof MURDEY, Richard James (Ph D)


Scope of Research

The laboratory explores the electronic and photo-electronic properties of organic semiconductor thin films, examining how the physics of charge generation and transport reflects the precise and intricate way molecules assemble, orient, and order in the solid state. Characterization of film growth and studies on the influence of environmental variables on the film structure are combined with *in situ* electrical conductance and photoconductivity measurements. The experimental conditions are carefully controlled to discover new details about charge injection and transport, charge generation, and intermolecular electronic interactions.


KEYWORDS

Organic Semiconductors
Molecular Electronics
Thin Films
Electrical Conductivity and Photoconductivity
Charge Generation and Transport


 $1 \, \mu m \times 1 \, \mu m$ AFM images of thin organic films grown on silicon wafers.




10 nm pentacene



10 nm pentacene + 10 nm C₆₀



10 nm C₆₀



10 nm C₆₀ + 10 nm pentacene

Selected Publications

Murdey, R.; Sato, N., Photocurrent Action Spectra of Organic Semiconductors, *Advances in Organic Crystal Chemistry Comprehensive Reviews 2015*, 627-652 (2015).

Murdey, R.; Sato, N., Interpretation of the Thermal Activation Energy of Conduction for Molecular Semiconductor Thin Films with Blocking Contacts, *Jpn. J. Appl. Phys.*, **53**, 05FY04 (2014).

Murdey, R.; Sato, N., Voltage Stress Induced Reversible Diode Behavior in Pentacene Thin Films, J. Chem. Phys., 137, 234703 (2012).

Murdey, R.; Sato, N., *In Situ* Conductance Measurements of Copper Phthalocyanine Thin Film Growth on Sapphire [0001], *J. Chem. Phys.*, **134**, 234702 (2011).

Nonlinear Photocurrent Spectroscopy

As photocurrent is proportional to the quantity of absorbed radiation it is possible in principle to extract an optical absorption spectrum using the photocurrent recorded at different wavelengths. Photocurrent in this instance can be considered as a measure of the absorbed light.

If the relationship between photocurrent, I, and the absorbed light intensity, F, is not linear, however, the incident light intensity for each data point must be attenuated such that the photocurrent remains constant – unless a mathematical function can be found to transform the photocurrent into a linear function of absorbance. We have determined that this linearization function, called the current response function, can be expressed as,

$$A = -\log\left(1 - \frac{1}{\gamma} \frac{I}{eF} \left(\frac{I}{I_d} + 1\right)^{\frac{1-\gamma}{\gamma}}\right) .$$

Y is a scaling parameter, while I_d and γ may be determined by a simple fitting procedure. The expression appears to be generally valid for most organic semiconductor films.

This new technique is called Nonlinear Photocurrent Spectroscopy (NPS). Absorption spectra can be quickly and easily obtained for thin and irregular shaped films, even on nontransparent substrates. There is no need to adjust the incident light intensity for each wavelength. Since NPS has very high sensitivity, the method is well-suited for examining gap states and impurity states in organic semi-conductors, as well as for studying very thin films down to a single monolayer. A demonstration of this sensitivity is shown in Figure 1.




Figure 1. Absorbance spectra of 20 nm and 1 nm free base phthalocyanine thin films derived from photocurrent measurements after application of the nonlinear current response function and scaled to an optical reference. The reference optical spectra for the 20 nm film measured with the transmitted beam intensity (dotted blue line) is also shown.

Conductance in Organic Radical Solids

Organic radical molecules are distinctive for their singly occupied molecular orbital (SOMO) which causes them to behave as narrow gap intrinsic semiconductors in the solid state.




Figure 2. The relation between the molecular orbitals and the solid state energy gap $E_{\rm g}$ in a) typical closed shell organic semiconductors, and b) open shell organic radicals. Charge held in gap states is said to be "trapped".

Radical bisphthalocyaninato complexes are open shell organic radical compounds which combine high electrical conductivity with chemical and thermal stability, making them promising materials for studying the charge transport of organic semiconductors. The small energy gap effectively removes the influence of deep traps, greatly enhancing the response time and facilitating reliable measurements.




Figure 3. Molecular structure of the phthalocyanine sandwich complex bis(phthalocyanininato)terbium(III) (TbPc₂). A single unpaired electron is shared between the two phthalocyanine ligands.

In recent work we have shown that the conductance equation of TbPc₂ thin films may be separated into exponential and linear terms in temperature. The energy gap is found by this method to be 0.316 eV.

Advanced Research Center for Beam Science – Particle Beam Science –

http://wwwal.kuicr.kyoto-u.ac.jp/www/index-e.htmlx



Assoc Prof IWASHITA, Yoshihisa (D Sc)



Techn Staff TONGU, Hiromu

Researcher(pt)

HOKONOHARA, Hisao

Lect (pt)

SHIMIZU, Hirohiko M. (D Sc)

Laboratory for Particle Properties, Department of Physics, Nagoya University

SAEKI, Takayuki (D Sc)

High Energy Accelerator Research Organization

Students

IMAJO, Sohei (D3) KITAHARA, Ryunosuke (D3) MIYAWAKI, Eisuke (M1) FUWA, Yasuhiro (D3) IKEDA, Hideki (M2)

Guest Scholar

JAMESON, Robert A. (Ph D) Goethe University, Frankfurt Germany, 3-29 November

Scope of Research


We are studying particle beam science which includes particle beam generation, acceleration and manipulation for fundamental sciences as well as for practical applications, such as new materials and cancer therapy. We also concentrate on electromagnetics design such as Neutron Optics, including neutron beam focusing to highly enhance their efficiency for advanced measurements. We are the first in the world to demonstrate active neutron acceleration in order to seek the neu-

tron Electric Dipole Moment. In addition, we contribute to advanced fault detection techniques for the International Linear Collider project superconducting accelerating cavities.



Beam Physics Accelerator Physics Neutron Optics

Phase Rotation International Linear Collider



Selected Publications

Imajo, S.; Mishima, K.; Kitaguchi, M.; Iwashita, Y.; Yamada, N. L.; Hino, M.; Oda, T.; Ino, T.; Shimizu, H. M.; Yamashita, S.; Katayama, R., Pulsed Ultra-cold Neutron Production Using a Doppler Shifter at J-PARC, *Prog. Theor. Exp. Phys.*, **2016-1**, 013C02-1-22 (2016).

Iwashita, Y.; Tongu, H.; Fuwa, Y.; Ichikawa, M., Compact Permanent Magnet H⁺ ECR Ion Source with Pulse Gas Valve, *Rev. Sci. Instrum.*, **87-2**, 02A718-1-02A718-3 (2016).

Fuwa, Y.; Iwashita, Y.; Tongu, H.; Inoue, S.; Hashida, M.; Sakabe, S.; Okamura, M.; Yamazaki, A., RF Synchronized Short Pulse Laser Ion Source, *Rev. Sci. Instrum.*, 87, 02A911-1-02A911-4 (2016).

Arimoto, Y.; Higashi, N.; Igarashi, Y.; Iwashita, Y.; Ino, T.; Katayama, R.; Kitaguchi, M.; Kitahara, R.; Matsumura, H.; Mishima, K.; Nagakura, N.; Oide, H.; Otono, H.; Sakakibara, R.; Shima, T.; Shimizu, H. M.; Sugino, T.; Sumi, N.; Sumino, H.; Taketani, K.; Tanaka, G.; Tanaka, M.; Tauchi, K.; Toyoda, A.; Tomita, T.; Yamada, T.; Yamashita, S.; Yokoyama, H.; Yoshioka, T., Development of Time Projection Chamber for Precise Neutron Lifetime Measurement Using Pulsed Cold Neutron Beams, *Nucl. Instr. Meth. Phys. Res. A*, **799**, 187-196 (2015).

Yamada, M.; Iwashita, Y.; Ichikawa, M.; Fuwa, Y.; Tongu, H.; Shimizu, H. M.; Mishima, K.; Yamada, N. L.; Hirota, K.; Otake, Y.; Seki, Y.; Yamagata, Y.; Hino, M.; Kitaguchi, M.; Kennedy, S. J.; Lee, W. T.; Andersen, K. H.; Guerard, B.; Manzin, G.; Geltenbort, P., Pulsed Neutronbeam Focusing by Modulating a Permanent-magnet Sextupole Lens, *Prog. Theor. Exp. Phys.*, **2015**, 043G01 (2015).

Kubo, T.; Iwashita, Y.; Saeki, T., Radio-frequency Electromagnetic Field and Vortex Penetration in Multilayered Superconductors, *Appl. Phys. Lett.*, **104**, 032603 (2014).

Measurement of Thermal Cross Section of ¹⁴N(n,p)¹⁴C Using Cold Neutron Beam at J-PARC

The cross section of the $^{14}N(n,p)^{14}C$ reaction is one of the key parameters in the slow-neutron capture process of stellar nucleosynthesis occurring in the helium burning shells whose typical temperature is ~ 25 keV. In late 1980s, the $^{14}N(n,p)^{14}C$ reaction cross section at around 30 keV have been measured by two groups, and their results deviated from each other by a factor of more than two. One of them has been supported by later measurements, but the extrapolated cross section of their results assuming the 1/v law differs from the recent evaluated cross section with the deviation of more than $1.6 \, \sigma$.

To improve the reliability of the evaluated cross section data, we measured the thermal cross section of the ¹⁴N(n,p)¹⁴C reaction using a pulsed cold neutron beam at J-PARC (Japan Proton Accelerator Research Complex). Our experiment has a feature that the ¹⁴N(n,p)¹⁴C and ³He(n,p)³H reactions are measured simultaneously by Time Projection Chamber (TPC), where the cross section of the latter reaction has been measured within the error of 0.13% as 5333(7) barn. The TPC, which was originally developed for a neutron lifetime measurement by the NOP (Neutron Optics and Physics) collaborators at J-PARC (Figure 1), consisted of a drift cage with a multi-wire proportional chamber (MWPC) inside the seal-off vacuum vessel (Figure 2). An operational gas was the mixture of 80 kPa ⁴He, 20 kPa N₂, and a few Pa ³He, where the partial pressures were measured by the gas injection system.

Pulsed cold neutron beams in the energy range of 1~20 meV are produced by the nuclear spallation reaction of Hg target using a 3 GeV pulsed proton beam at MLF (the Materials and Life Science Facility). Since the length of a neutron bunch is 40 cm which is shorter than the length of the TPC sensitive volume, background events caused by neutron capture reactions in materials of inner walls or




Figure 1. Experimental set up at J-PARC/MLF Beam Line 05.

frames of the TPC could be reduced by selecting only the events whose particle tracks were fully contained in the sensitive volume. Since the Q-values of the ¹⁴N(n,p)¹⁴C and ³He(n,p)³H reactions are 626 keV and 764 keV, respectively, they were able to be separately detected by the deposit energy difference of the reactions (Figure 3). Thus we could deduce the thermal cross section of ¹⁴N(n,p)¹⁴C relative to the thermal cross section of ³He(n,p)³H by using the known ratios of the target densities of ¹⁴N and ³He and the detection efficiencies for (n,p) reactions on those nuclei.

To check the quality of the separation between the $^{14}N(n,p)^{14}C$ and $^{3}He(n,p)^{3}H$ reactions in the energy spectrum, we made measurements using the operational gases with three different partial pressures of ^{3}He , and obtained a preliminary result on the thermal cross section of the $^{14}N(n,p)^{14}C$ reaction as 1.864(3) barn with the world best accuracy of 0.3%.

The method of our experiment is in principle applicable to the measurements of other (n,p) and (n,α) reaction cross sections, for example, ${}^{10}B(n,p){}^{10}Be$, which is also one of the important reactions in astrophysics. We are considering a measurement of the thermal cross section of the ${}^{10}B(n,p){}^{10}Be$ reaction in a near future.




Figure 2. Schematic drawing of the Time Projection Chamber (TPC). Electrons ionized by a charged particle in the operational gas drift to the multi-wire proportional counter (MWPC). Deposit energy in the TPC is determined by measuring an amount of the electrons detected by MWPC.




Figure 3. Deposit energy distribution in the measurement with the ³He partial pressure of 9 Pa. The peaks corresponding to the ¹⁴N(n,p)¹⁴C and ³He(n,p)³H reactions are found in the energy regions of 520~700 keV and 700~880 keV, respectively, and are well separated with the threshold level of 700 keV (blue line).

Advanced Research Center for Beam Science – Laser Matter Interaction Science –

http://laser.kuicr.kyoto-u.ac.jp/e-index.html



Prof SAKABE, Shuji (D Eng)



Assoc Prof HASHIDA, Masaki (D Eng)



Assist Prof INOUE, Shunsuke (D Sc)



PD NAKAMIYA, Yoshihide (D Sc)

Lect (pt)

TOKITA, Shigeki (D Eng) Osaka University

Researcher (pt)

MASUNO, Shinichiro

Students

MORI, Kazuaki (D3) FURUKAWA, Yuki (M1) TERAMOTO, Kensuke (D2)


Scope of Research

The interaction of femtosecond laser pulses with matter involves interesting physics not seen with nanosecond laser pulses. Through investigation of the interaction physics, the potential of intense femtosecond lasers for new applications is being developed (e.g., laser-produced radiation and laser processing). Ultra-intense lasers can produce intense radiations (e.g., electrons, ions, and THz), which are promising as next-generation radiation sources. Ultra-short lasers can process any matter without thermal dissociation. Femtosecond laser processing is also the next-generation of laser processing.

Our laboratory is equipped with an ultra-intense femtosecond laser named T6, to study the physics of intense laser–matter interactions and its applications.

KEYWORDS

Intense Laser Science
Laser Plasma Radiations (electrons, ions, and THz)
Ultrafast Electron Diffraction (UED)
Laser Nano-ablation Physics
Femtosecond Laser Processing



Selected Publications

Gouda, A. M.; Sakagami, H.; Ogata, T.; Hashida, M.; Sakabe, S., The Formation Mechanism of the Periodic Nanograting Structure by the Weibel Instability, *Appl. Phys. A*, **122**, 454-1-454-6 (2016).

Hashida, M.; Nishii, T.; Miyasaka, Y.; Sakagami, H.; Shimizu, M.; Inoue, S.; Sakabe, S., Orientation of Periodic Grating Structures Controlled by Double-pulse Irradiation, *Appl. Phys. A*, **122**, 484-1-484-5 (2016).

Furukawa, Y.; Sakata, R.; Konishi, K.; Ono, K.; Matsuoka, S.; Watanabe, K.; Inoue, S.; Hashida, M.; Sakabe, S., Demonstration of Periodic Nanostructure Formation with Less Ablation by Double-pulse Laser Irradiation on Titanium, *Appl. Phys. Lett.*, **108**, 264101(2016).

Hashida, M.; Nishii, T.; Miyasaka, Y.; Sakagami, H.; Shimizu, M.; Inoue, S.; Sakabe, S., Threshold Fluence for Femtosecond Laser Nanoablation for Metals, *Electr. Communi. Jpn*, **99**, 88-95 (2016).

Inoue, S.; Maeda, K.; Tokita, S.; Mori, K.; Teramoto, K.; Hashida, M.; Sakabe, S., Single Plasma Mirror Providing 104 Contrast Enhacement and 70% Reflectivity for Intense Femtosecond Lasers, *Appl. Opt.*, **55**, 5647-5651 (2016).

Highly Intensified Emission of the Laser Accelerated Fast Electrons by Using Solid-plasma Composite Target

Intense ultrashort electrons pulses are driven by the interaction of intense short laser pulses with solid targets. These laser-accelerated fast electrons have many possible applications such as fast ignition for inertial confinement fusion, ultrafast electron diffraction measurement, and ultrafast transient field measurement. For these applications using fast electrons as probe pulses with high temporal resolution, it is desirable for a greater number of electrons to be emitted from the laser plasma. However, most laser-accelerated electrons cannot escape from the laser plasma because they are trapped by a strong quasi-static electric field, called the sheath field, produced around the steep density gradient boundary between the solid/plasma and the vacuum. Almost all electrons are bound within the solid target, and only about 1% of the hot electrons can escape. Therefore, only a small fraction of electrons accelerated by intense short-pulse laser escapes from the laser plasma, and the most of electrons expend their energy heating the target or producing other types of radiation.

We demonstrate the intensification of electrons escaping from an intense laser plasma by using double femtosecond laser pulses. An intense pulse from a chirped pulse amplification laser (CPA1) for driving fast electrons is used to irradiate a foil target, the rear of which is pre-irradiated with another laser pulse (CPA2). Pre-irradiation with CPA2 controls the electron density distributions in the target to suppress sheath field growth and expand the target plasma into which the fast electrons are released. The number of escaping electrons increases greatly when the target is irradiated with CPA2 540 ps prior to CPA1. The number of electrons released is 7 times that for single pulse irradiation. These results are supported by two-dimensional (2D) particle-in-cell (PIC) simulations of plasma produced by CPA2 and analytical evaluation considering the expansion of the plasma.




Figure 1. (a) schematic of the experimental setup. (b) Images of fast electrons emitted from laser plasma.

Threshold Fluence of Femtosecond Laser Nano-Ablation for Metals

Femtosecond laser nano-ablation of Ti by short-pulse

laser irradiation (800 nm/40 fs) is studied in the laser fluence range of 0.07 – 0.5 J/cm². To determine the ablation threshold, the ablation rate dependence on laser fluence is precisely measured. Multi-shot ablation threshold of titanium is found to be 0.074 J/cm² in which value is good agreement with that reported previously by other group. To discuss the ablation mechanism, all the ablation thresholds for metals previously published are plotted as a function of work function and melting temperature. We found that the ablation thresholds correlated with work function of metals. Experimental data suggested that the femtosecond laser ablation is mainly due to multi-photon absorption and optical field ionization.

Reduction of Ablation Rate by Double-Pulse Laser Irradiation on Titanium

Reduction of ablation rate is demonstrated on a titanium surface irradiated by a double-pulse beam parallel-polarized at time delays from $\Delta t = 0.16 - 1280$ ps. The first-pulse fluence F_1 and the delayed pulse fluence F_2 are kept below or above the ablation threshold $F_{\rm TH} \sim 100 \ {\rm mJ/cm^2}$ of Ti. A pair of laser pulses with fluences of 70 and 140 mJ/cm² is used for ablation rate experiment. Figures 1 show the dependence of ablation rate on interval Δt for the fluences $(F_1, F_2) = (70, 140)$. The maximum and minimum ablation rates are 15.5 nm for $\Delta t = 0.16$ ps and 4.4 nm for $\Delta t = 80$ ps, respectively. The maximum is as much as that (17.1 nm) for a single pulse of 210 mJ/cm²; the minimum (4.4 nm) is half of that for a 140 mJ/cm² single pulse. Thus, the first pulse modifies the surface to suppress ablation even if its fluence is below the ablation threshold. The experimental results suggest that the first pulse influences the surface, suppressing ablation by the following pulse even though no laser plasma is produced on the surface, and that this influence continues for rather a long time (several hundred picoseconds). Double-pulse irradiation certainly has significant potential for future applications of laser nano-processing.




Figure 2. Dependence of ablation rate on time interval for fluences $(F_1, F_2) = (70, 140)$

Advanced Research Center for Beam Science – Electron Microscopy and Crystal Chemistry –

http://eels.kuicr.kyoto-u.ac.jp/Root/English



Prof KURATA, Hiroki (D Sc)



Assist Prof NEMOTO, Takashi (D Sc)



Assist Prof HARUTA, Mitsutaka (D Sc)



Program-specific Res* OGAWA, Tetsuya (D Sc)



Program-specific Res* KIYOMURA, Tsutomu

*Nanotechnology Platform

Students

FUJIYOSHI, Yoshifumi (D3) YAMAGUCHI, Atsushi (D3) LAI, Ming Wei (D2) INOUE, Shota (M2)

HIGUCHI, Hiroki (M1)

Guest Scholar

SCHAPER, Andreas Karl (Ph D) Phillips University of Marburg, Germany, 26 September–17 October

Scope of Research

We study crystallographic and electronic structures of materials and their transformations through direct imaging of atoms or molecules by high-resolution electron spectromicroscopy, which realizes energy-filtered imaging and electron energy-loss spectroscopy as well as high-resolution imaging. By combining this with scanning probe microscopy, we cover the following subjects: 1) direct structure analysis, electron crystallographic analysis, 2) epitaxial growth of molecules, 3) structure formation in solutions, and 4) fabrication of low-dimensional functional assemblies.

KEYWORDS

EELS STEM Elemental Ratio Surface Plasmon Electronic Structure





Selected Publications

Haruta, M.; Kurata, H., Direct Observation of Crystal Defects in an Organic Molecular Crystals of Copper Hexachlorophthalocyanine by STEM-EELS, Sci. Rep., 2, [252-1]-[252-4] (2012).

Aso, R.; Kan, D.; Shimakawa, Y.; Kurata, H., Atomic Level Observation of Octahedral Distortions at the Perovskite Oxide Heterointerface, Sci. Rep., 3, [2214-1]-[2214-6] (2013).

Saito, H.; Namura, K.; Suzuki, M.; Kurata, H., Dispersion Relations for Coupled Surface Plasmon-polariton Modes Excited in Multilayer Structures, *Microscopy*, **63**, 85-93 (2014).

Saito, H.; Kurata, H., Formation of a Hybrid Plasmonic Waveguide Mode Probed by Dispersion Measurement, *J. Appl. Phys.*, **117**, [133107-1]-[133107-7] (2015).

Haruta, M.; Hosaka, Y.; Ichikawa, N.; Saito, T.; Shimakawa, Y.; Kurata, H., Determination of Elemental Ratio in an Atomic Column by Electron Energy-Loss Spectroscopy, *ACS Nano*, **10**, 6680-6684 (2016).

Determination of Elemental Ratio in an Atomic Column by Electron Energy Loss Spectroscopy

Elemental mapping with atomic-column resolution can be obtained using electron energy loss spectroscopy (EELS) combined with scanning transmission electron microscopy (STEM). One step in STEM-EELS is quantitative determination of elemental ratio in an atomic column. However, during elemental mapping using STEM-EELS, the elemental signals do not necessarily localize at atomic-column positions. The spatial resolution of an EELS signal is constrained by the delocalization of inelastic scattering. Careful attention to the mixing of signals from neighboring atomic-columns is required to analyze a spectrum at a single point. In the present study, the elemental ratio of Fe to Mn at octahedral and tetrahedral sites in brownmillerite Ca₂Fe_{1.07}Mn_{0.93}O₅ was determined quantitatively with atomic resolution using STEM-EELS. This study presents the experimental criteria for semiquantitative analysis of the L_{2,3}-edges of Fe and Mn in a perovskite-related structure.

A HAADF image along the [101] direction and an EELS data cube for the Fe and Mn $L_{2,3}$ -edges were simultaneously collected using an aberration-corrected JEM ARM-200F operating at 80 kV with a probe-forming aperture of 21.3 mrad. Spectra were recorded as line scans along the (010) direction. The scan step was 0.19 Å/pixel, which corresponded to 20 segment divisions in the B–B' cation distance. In the present research, the line-profiling method was used instead of two-dimensional mapping to avoid some practical problems (damage and stability) due to oversampling conditions.

Figure 1 shows a HAADF image with the projected structure overlaid. The cation L_{2,3}-edge intensities after background-subtraction are shown in Figure 2b. The individual spectra extracted from the octahedral and tetrahedral sites contain a weak intensity from the nonpreferred cation at both polyhedral sites, as shown in Figure 2d. The delocalization factors of the Fe and Mn L23-edge excitations were approximately 1.2 and 1.3 Å, respectively, which are small enough to resolve the atomic distance (3.8 Å) between B and B' cations. The relative composition profiles obtained by quantifying the intensity at each spatial increment in the elemental profiles of Figure 2b are shown in Figure 2c. The relative composition profiles resembled wide isosceles trapezoids. The elemental profile reflects the one-dimensional distribution of excitation probabilities for a specific energy loss; a maximal excitation should occur just above each atomic position. In contrast, the relative composition profile should be constant near each atomic position if the spectra contain only the contribution from illuminated atomic columns. The flat region of the relative composition profile (Figure 2c) indicates very little contribution from nearest neighbor columns and the quantitative analysis is achieved near-atomic resolution. Such a profile can be obtained only by oversampling of spectral data. The present experiment found that the relative composition profile was nearly constant within a 1.2 Å radius around the atomic position of each polyhedral site. The average ratio of Fe to Mn in this flat region (Figure 2c) was (17.5:82.5) \pm 5.9 in octahedral sites and (81.6:18.4) \pm 6.0 in tetrahedral sites. The result at octahedral sites roughly agreed with previous results from neutron diffraction, $(14.4:85.6) \pm 0.2$. Simple multislice calculations were used to estimate the error due to electron dechanneling combined with a delocalization factor. The Mn atomic fraction profile derived from EELS intensity ratio was calculated by assuming "true fractions" of Mn at octahedral site, x=0.87. The simulated profile accurately reproduced the flat region near an octahedral site, as expected, with a radius of 0.9 Å, which agreed well with the experimental value. For a thin sample (t < 10 nm), the estimated error was negligibly small.

Results showed that relative composition profiles with an isosceles trapezoid-shape are useful for quantifying the resolution of measurements. An area with a radius of about 1.2 Å surrounded both atomic sites and provided information at near-atomic resolution. This study demonstrates that the experimental spectra at the L_{2,3}-edge, obtained from octahedral atomic columns of 3d transition metals in perovskite-like structures, can be quantitatively interpreted with an uncertainty of 10% (accuracy of calculated partial cross sections of L-shell using Hartree-Slater model) without full quantum mechanical simulation.





Figure 1. HAADF image of the [101] direction in $Ca_2Fe_{1.07}Mn_{0.93}O_5$. Inset shows the corresponding structure. Spectral imaging was performed along the dashed line.

Figure 2. Line profiles and spectra for $Ca_2Fe_{1.07}Mn_{0.93}O_5$: (a) HAADF profiles; (b) EELS elemental (010) line profiles for the $L_{2.3}$ -edges of Fe and Mn; (c) profile of the relative compositions of Fe and Mn; (d) spectra measured above the octahedral and tetrahedral cations.



Advanced Research Center for Beam Science — Structural Molecular Biology —

http://www.scl.kyoto-u.ac.jp/~fujii/indexE.html



Assoc Prof ITO, Yoshiaki (D Sc)



Assist Prof FUJII, Tomomi (D Sc)


Scope of Research


Our laboratory analyzes the electronic state of atomic or molecular structure in materials, which are obtained from diffraction images and spectra observed by X-ray irradiation of these substances, respectively, to elucidate the relationships between the structure and functions or physical properties. Our main themes are (1) high-res-

KEYWORDS

Crystal X-ray Crystallographic Analysis Structural Biology Protein Crystallography Structure and Function

olution experimental and theoretical studies on the natural width of elements in materials, (2) the development in the measurements of the diagram lines in soft X-rays region, (3) structural determination of novel protein molecules and their complexes, and (4) studies on the structural basis of functions, physical properties, and intermolecular interactions.







Selected Publications

Fujii, T.; Sato, A.; Okamoto, Y.; Yamauchi, T.; Kato, S.; Yoshida, M.; Oikawa, T.; Hata, Y., The Crystal Structure of Maleylacetate Reductase from *Rhizobium* sp. Strain MTP-10005 Provides Insights into the Reaction Mechanism of Enzymes in Its Original Family, *Proteins: Structure, Function, and Bioinformatics*, **84**, 1029-1042 (2016).

Fujii, T.; Yamauchi, T.; Ishiyama, M.; Gogami, Y.; Oikawa, T.; Hata, Y., Crystallographic Studies of Aspartate Racemase from *Lactobacillus sakei* NBRC 15893, *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.*, **71**, 1012-1016 (2015).

Fujii, T.; Goda, Y.; Yoshida, M.; Oikawa, T.; Hata, Y., Crystallization and Preliminary X-ray Diffraction Studies of Maleylacetate Reductase from *Rhizobium* sp. Strain MTP-10005, *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.*, **64**, 737-739 (2008).

Fujii, T.; Oikawa, T.; Muraoka, I.; Soda, K.; Hata, Y., Crystallization and Preliminary X-ray Diffraction Studies of Tetrameric Malate Dehydrogenase from the Novel Antarctic Psychrophile *Flavobacterium frigidimaris* KUC-1, *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.*, **63**, 983-986 (2007).

Fujii, T.; Sakai, H.; Kawata, Y.; Hata, Y., Crystal Structure of Thermostable Aspartase from *Bacillus* sp. YM55-1: Structure-based Exploration of Functional Sites in the Aspartase Family, *J. Mol. Biol.*, **328**, 635-654 (2003).

Structural Features and Low-temperature Adaptation of Aspartate Racemase

Amino-acid racemases are responsible for the racemization of amino acids and can be mainly grouped into two families, pyridoxal 5'-phosphate (PLP) dependent and PLP independent. Aspartate racemase (AspR) catalyzes the interconversion between L- and D-aspartate and belongs to the PLP-independent racemase group. The enzyme is thought to employ a two-base mechanism to catalyze both the directions of racemization and utilize two cysteine residues as the conjugated catalytic acid and base in the catalytic reaction. The only crystal structure of the PLP-independent amino-acid racemase is now available from a hyperthermophilic archaeon. To elucidate the structure and the low-temperature adaptation of the racemase group, we determined the crystal structures of AspR from Lactobacillus sakei NBRC 15893 (LsAspR), which works in the low-to-medium temperature range, and for comparison AspR from Thermococcus litoralis DSM 5473 (TlAspR), which has the maximum activity at 95 °C.

LsAspR and TlAspR were crystallized at 20 °C by the sitting-drop vapour-diffusion method using a precipitant solution of 25% (v/v) PEG-MME 550, 5% (v/v) 2-propanol and 0.1 M sodium acetate pH 4.8 and a precipitant solution of 24% (w/v) PEG1500, 0.2 M L-proline and 0.1 M HEPES pH 7.5, respectively. Diffraction experiments were performed at beamlines NE-3A of Photon Factory AR, and BL-5A of Photon Factory, Tsukuba, Japan. The crystals were flash-cooled in a nitrogen stream at 100 K. Diffraction data were collected at a wavelength of 1.000 Å using Quantum 270 and 315r CCD detectors. The crystals belonged to space groups P3₁21 with unit cell parameters of a = b = 105.5 Å and c = 96.5 Å, and $P2_12_11$ with unit cell parameters of a = 90.26, b = 125.78, c = 40.64 Å, respectively. The structures of LsAspR and TlAspR were determined by molecular replacement and refined at 2.6 Å resolution (R = 23.8%, $R_{\text{free}} = 31.6\%$) and 2.0 Å resolution $(R = 18.7\%, R_{\text{free}} = 25.0\%)$, respectively.

Both LsAspR and TlAspR molecules are homodimers with molecular two-fold axis. The subunit of each enzyme molecule comprises the N-terminal and C-terminal domains. In each domain, a central four-stranded parallel β -sheet is flanked by six α -helices. The molecules are formed mainly by intersubunit interactions between the N-terminal α -helices and intersubunit hydrogen-bonds between the N-terminal β -sheets in the dimer interface. The active-site cleft exists between both the domains. The spatial arrangement of the strictly conserved cysteine residues in the cleft reveals the Cys residues involved in the enzymatic catalysis: Cys84 and Cys196 of LsAspR and Cys83 and Cys194 of TlAspR.

A structural comparison of LsAspR and TlAspR reveals structural factors probably involved in thermostability of AspR. The molecular volume, intersubunit interaction, and the number of ion pairs suggest that the LsAspR molecule is more loose than that of TlAspR (Fig. 1). Most characteristic difference at the dimer interface between LsAspR and TlAspR is the interactions between α 1 helices (Fig. 2). Moreover, π - π interaction between indole rings of Trp74 is found at the dimer interface of TlAspR (Fig. 3). In LsAspR, Leu75 corresponds to Trp74 in TlAspR and does not make an intersubunit interaction.



Figure 1. Superposition of LsAspR and TlAspR dimers. Cyan, LsAspR; Red, TlAspR.




Figure 2. Interactions between $\alpha 1$ helices in dimer interface. (Left panel) LsAspR. (Right panel) TlAspR.




Figure 3. π - π Interaction between indole rings of Trp74 at dimer interface of TlAspR.

International Research Center for Elements Science - Synthetic Organotransformation -

http://es.kuicr.kyoto-u.ac.jp/index_en.html



Prof NAKAMURA, Masaharu (D Sc)



Assoc Prof TAKAYA, Hikaru (D Eng)



Assist Prof ISOZAKI, Katsuhiro (D Eng)



Program-Specific Assist Prof Program-Specific Assist Prof ADAK, Laksmikanta (D Eng)



IWAMOTO, Takahiro (D Eng)



PD PINCELLA, Francesca (Ph D)



PD FAN, Lulu (Ph D)

Researcher (pt) MATSUDA, Hiroshi

Students

NAKAJIMA, Sho (D3) YOSHIDA, Ryota (D3) AGATA, Ryosuke (D2) AOKI, Yuma (D2)

OKUZONO, Chiemi (M1) UENO, Ryo (M1) HOSOKAWA, Atsushi (UG) SHANOH, Takafumi (UG)

Guest Res Assoc

O'BRIEN, Harry University of Bristol, UK, 21 June-29 August


Scope of Research

Our research activity focuses on the development of new molecular transformations, which can contribute to better or ideal synthesis of functional molecules as well as to exploitation of new chemical (metal and carbon) resources. The present research subjects are (1) metal-catalyzed carbon-carbon and carbon-heteroatom bond forming reactions by using

universal metals such as iron (2) development of smart materials based on synergistic effect of various metals on artificial peptides (3) development of smart metallic nanoparticle catalysts based on supramolecular approaches (4) understanding of reaction mechanism of these catalytic reactions with the help of quantum chemical methods and synchrotron X-ray absorption spectroscopy.

KEYWORDS

Selective Organic Reaction Metalated Peptide Catalyst Solution-Phase XAS Analysis Iron Catalyst Metallic Nanoparticle Catalyst



Selected Publications

Yoshida, R.; Isozaki, K.; Yokoi, T.; Yasuda, N.; Sadakane, K.; Iwamoto, T.; Takaya, H.; Nakamura, M., ONO-pincer Ruthenium Complexbound Norvaline for Efficient Catalytic Oxidation of Methoxybenzenes with Hydrogen Peroxide, Org. Biomol. Chem., 14, 7468-7479 (2016). Yamaguchi, A.; Isozaki, K.; Nakamura, M.; Takaya, H.; Watanabe, T., Discovery of 12-mer Peptides that Bind to Wood Lignin, Scientific Reports, 6, 21833 (2016).

Isozaki, K.; Yokoi, T.; Yoshida, R.; Ogata, K.; Hashizume, D.; Yasuda, N.; Sadakane, K.; Takaya, H.; Nakamura, M., Synthesis and Functions of ONO-Pincer Ruthenium-Complex-Bound Norvalines, Chemistry an Asian Journal, 11, 1076-1091 (2016).

Iron-Catalyzed Cross-Coupling Reactions

Transition-metal-catalyzed carbon-carbon and carbonheteroatom bond forming reactions are powerful tools for the synthesis of functional molecules. Our group focuses on use of iron as a catalyst, and so far developed a variety of iron-catalyzed cross-coupling reactions, some of which can't be achieved by conventional transition-metal catalysts such as palladium. For example, we recently developed iron-catalyzed enantioselective cross-coupling reaction of α-chloroesters with aryl Grignard reagents, which is the first iron-catalyzed enantioselective coupling reaction. In addition, we found a novel intermolecular and intramolecular C-H amination of diarylamines to provide dihydrophenazine derivatives. Importantly, these rections enable facile access to several important pharmaceutical and functional molecules. Furthermore we clarified the mechanisms of these reactions by using synchrotron X-ray absorption spectroscopy combined with theoretical calculations.



Figure 1. (a) Iron-catalyzed enantioselective cross-coupling reaction of α -chloroesters with aryl Grignard reagents, and (b) intermolecular and intramolecular C-H amination of diarylamines.

Novel Transition-Metal Catalyst Bound with Functional Amino Acid or Peptide

This project focuses on development of smart materials based on synergistic effect of various metals on artificial peptides. Based on this concept, novel ruthenium catalysts bound with norvaline have been developed, achieving efficient and selective oxidation of several methoxy-benzene analogues to quinones. In this reaction, norvaline act as a mediator inducing micelle-like aggregates, which is the origin of high efficiency and selectivity. Now we are trying further modification of the catalysts to oxidize much challenging aromatic substrate toward investigation of future chemical resources.

Figure 2. Efficient catalytic oxidation of methoxybenzenes with ruthenium catalyst bound with norvaline.

Supramolecular Approach for Creating Enhanced Catalysis of Metallic Nanoparticles

This project focuses on the development of highly active and selective metallic nanoparticle and cluster catalysts. To achieve this purpose, four key methodologies have been developed: 1) creation of reaction field with advanced self-assembly, 2) design of selective molecular transformation with multi-points intermolecular interaction, 3) size selective synthesis of metallic nanoparticles and clusters, and 4) utilization of plasmonic resonance with light.



International Research Center for Elements Science - Advanced Solid State Chemistry –

http://www.scl.kyoto-u.ac.jp/~shimakgr/indexE.html



Prof SHIMAKAWA, Yuichi (D Sc)



Assoc Prof KAN, Daisuke (D Sc)



Assist Prof SAITO, Takashi (D Sc)



Techn Staff ICHIKAWA, Noriya (D Eng)



PD (JSPS) FABIO, Denis Romero (Ph D)

Students


HOSAKA, Yoshiteru (D3) GUO, Haichuan (D3) XIONG, Peng (D2) OZAKI, Yusuke (M2) NISHIMURA, Tomoe (M2) WADA, Tamaki (M2) TAN, Zhenhong (M1) NIWA, Yasuyuki (M1)

Scope of Research

Transition metal oxides have a wide variety of interesting and useful functional properties, including electronic conduction, superconductivity, ferroelectricity, and ferromagnetism. In fact, some of these oxides are used in current electronic devices. Our research mainly focuses on perovskite-structured transition metal oxides with novel functional properties due to complex couplings between their lattices, charges and spins. We are currently exploring such functional oxides with advanced oxide-synthesis techniques such as high-pressure synthesis and epitaxial thin film growth.

KEYWORDS

Solid State Chemistry Functional Transition Metal Oxides Epitaxial Thin Film Growth High Pressure Synthesis Perovskite Structured Oxides



Selected Publications

Kan, D.; Aso, R.; Sato, R.; Haruta, M.; Kurata, H.; Shimakawa, Y., Tuning Magnetic Anisotropy by Interfacially Engineering the Oxygen Coordination Environment in a Transition-metal Oxide, *Nat. Mater.*, **15**, 432-437 (2016).

Chen, W.-T.; Mizumaki, M.; Seki, H.; Senn, M.; Saito, T.; Kan, D.; Attfield, J. P.; Shimakawa, Y., A Half-metallic A- and B-Site-ordered Quadruple Perovskite Oxide CaCu₃Fe₂Re₂O₁₂ with Large Magnetization and a High Transition Temperature, *Nat. Comm.*, **5**, [3909-1]-[3909-7] (2014).

Matsumoto, K.; Haruta, M.; Kawai, M.; Sakaiguchi, A.; Ichikawa, N.; Kurata, H.; Shimakawa, Y., Selective Reduction of Layers at Low Temperature in Artificial Superlattice Thin Films, *Sci. Rep.*, 1, [27-1]-[27-14] (2011).

Inoue, S.; Kawai, M.; Ichikawa, N.; Kageyama, H.; Paulus, W.; Shimakawa, Y., Anisotropic Oxygen Diffusion at Low Temperature in Perovskite-structure Iron Oxides, *Nat. Chem.*, **2**, 213-217 (2010).

Long, Y. W.; Hayashi, N.; Saito, T.; Azuma, M.; Muranaka, S.; Shimakawa, Y., Temperature-induced A-B Intersite Charge Transfer in an A-Site-ordered LaCu₃Fe₄O₁₂ Perovskite, *Nature*, **458**, 60-63 (2009).

Ferromagnetism Induced by Substitution of the Iron(IV) Ion by an Unusual High-Valence Nickel(IV) Ion in Antiferromagnetic SrFeO₃

In addition to the usual valence states of 3d transition-metal cations in oxides, such as Fe^{II}/Fe^{III} for Fe and Ni^{II} for Ni, unusual high-valence states, such as Fe^{IV} and Ni^{IV}, can be stabilized in a strongly oxidizing atmosphere. Iron(IV) is stabilized in the cubic perovskite SrFeO₃, which is synthesized at high pressure and temperature. The iron(IV) ion in the perovskite, having a nominal d⁴ electron configuration in the corner-sharing FeO₆ octahedron, has a spin of S=2, and at low temperatures the compound shows a helical magnetism as a result of competing ferromagnetic and antiferromagnetic interactions between the spins. Ni^{IV}, on the other hand, is stabilized in hexagonal perovskites such as SrNiO₃ and BaNiO₃ and is expected to have a low spin state (S=0). Indeed, BaNiO₃ is nonmagnetic.

In the course of preparing a solid solution of $SrFeO_3$ and $SrNiO_3$, we found that Ni^{IV} is stabilized in the $SrFeO_3$ -type cubic perovskite $Sr(Fe^{IV}_{I-x}Ni^{IV}_x)O_3$ with x up to 0.5. Interestingly, the substitution of Fe^{IV} ions by Ni^{IV} ions in the helical antiferromagnetic $SrFeO_3$ induces ferromagnetism with a transition temperature (T_c) higher than room temperature. X-ray absorption spectroscopy and x-ray magnetic circular dichroism spectroscopy imply that substantial magnetic moments of the Ni^{IV} ion were induced by the large magnetic moments of Fe^{IV} through orbital hybridization with oxygen and as a result of this orbital hybridization, ferromagnetism, with the transition temperature above room temperature, is achieved.




Figure 1. Temperature dependence of magnetic susceptibility for Sr(Fe_{1-x}Ni_x)O₃ measured under 10 kOe magnetic field upon heating after zero-field cooling

Interface Engineering of Oxygen Coordination Environment as a New Route for Exploring Functional Properties of Transition Mental Oxides

The oxygen coordination environments of the transition metals in transition metal oxides often underpin a wide variety of functional properties through crystal field splitting and magnetic interactions. It has been demonstrated that artificial heterostructures and superlattices consisting of dissimilar oxides provide a good platform on which structural distortions associated with changes in metal-oxygen bonds, namely, the oxygen coordination environments, can be engineered. We show that the oxygen coordination environment (or the Ru-O-Ru bond angle) in SRO can be interfacially controlled by inserting a very thin CSTO layer (0-4 monolayers thick) into the SRO/GSO heterointerface. Our aberration-corrected scanning transmission electron microscopy (STEM) observations reveal that the Ru-O-Ru bond angle in the entire SRO layer is controlled through the Ru-O-Ti bond angle, which is determined by the oxygen position at the SRO/CSTO interface. More importantly, the Ru-O-Ti bond angle (or the interfacial oxygen position) can be engineered by layer-by-layer control of the thickness of the CSTO buffer layer that stabilizes the oxygen coordination environment, a Ru-O-Ru bond angle not seen in bulk SRO. Our magneto-transport characterization of the SRO/ CSTO/GSO heterostructures also shows that the Ru-O-Ti bond angle plays a decisive role in magnetic anisotropy with the uniaxial magnetic easy axis of the SRO layer, highlighting the significant effect of the interfacially engineered oxygen coordination environment on the spin-orbit interaction of the SRO layer.




Figure 2. Interface engineering of oxygen coordination environments of perovskite SrRuO₃.

International Research Center for Elements Science - Organometallic Chemistry -

http://om.kuicr.kyoto-u.ac.jp



Prof OZAWA, Fumiyuki (D Eng)



Assist Prof (D Eng)



Assist Prof WAKIOKA, Masayuki TAKEUCHI, Katsuhiko (D Sc)

Researcher(pt)

ICHIHARA, Nobuko

Students

TAGUCHI, Hiro-omi (D1) SASAKI, Daichi (M2) TAKAHASHI, Rina (M2)

TANAKA, Yuto (M1) YAMASHITA, Natsumi (M1) XU, Kai (M1)

MORITA, Hazuki (UG) WAKAYAMA, Rin (UG) HARADA, Kazuki (UG)

Scope of Research

This laboratory aims to establish new synthetic methodologies and new functional materials by designing well-defined catalysts based on transition metal chemistry. New concepts and ideas of molecular-based catalysts are accumulated by mechanistic investigations using experimental methods such as spectroscopy and kinetic techniques, as well as theoretical methods. The research subjects include: 1) development of novel organotransition metal systems for catalysis based on precise ligand design, and 2) preparation of π -conjugated polymers using direct arylation.

KEYWORDS

Transition Metal Complex Homogeneous Catalyst Reaction Mechanism Low-coordinate Phosphorus Ligand π-Conjugated Polymer



Selected Publications

Takeuchi, K.; Taguchi, H.; Tanigawa, I.; Tsujimoto, S.; Matsuo, T.; Tanaka, H.; Yoshizawa, K.; Ozawa, F., A Square Planar Complex of Platinum(0), Angew. Chem. Int. Ed., 55, 15347-15350 (2016).

Ozawa, F.; Nakajima, Y., PNP-Pincer Type Phosphaalkene Complexes of Late Transition Metals, Chem. Rec., 16, 2314-2323 (2016).

Taguchi, H.; Sasaki, D.; Takeuchi, K.; Tsujimoto, S.; Matsuo, T.; Tanaka, H.; Yoshizawa, K.; Ozawa, F., Unsymmetrical PNP-Pincer Type Phosphaalkene Ligands Protected by a Fused-Ring Bulky Eind Group: Synthesis and Applications to Rh(I) and Ir(I) Complexes, Organometallics, 35, 1526-1533 (2016).

Iizuka, E.; Wakioka, M.; Ozawa, F., Mixed-Ligand Approach to Palladium-Catalyzed Direct Arylation Polymerization: Effective Prevention of Structural Defects Using Diamines, *Macromolecules*, 49, 3310-3317 (2016).

Nakajima, Y.; Tsuchimoto, T.; Chang, Y.-H.; Takeuchi, K.; Ozawa, F., Reactions of [Cu(X)(BPEP-Ph)] (X = PF₆, SbF₆) with Silyl Compounds. Cooperative Bond Activation Involving Non-coordinating Anions, *Dalton Trans.*, **45**, 2079-2084 (2016).

A Square Planar Complex of Platinum(0)

The Pt(0) complex [Pt(PPh₃)(Eind₂-BPEP)] with a pyridine-based PNP-pincer-type phosphaalkene ligand (Eind₂-BPEP) has a highly planar geometry around Pt with $\Sigma(Pt) = 358.6^{\circ}$. This coordination geometry is very uncommon for formal d¹⁰ complexes, and the Pd and Ni homologues with the same ligands adopt distorted tetrahedral geometries. DFT calculations reveal that both the Pt and Pd complexes are M(0) species with nearly ten valence electrons on the metals whereas their atomic orbital occupancies are evidently different from one another. The Pt complex has a higher occupancy of the atomic 6s orbital because of strong s—d hybridization due to relativistic effects, thereby adopting a highly planar geometry reflecting the shape and orientation of the partially unoccupied d_{x2-y2} orbital.




Figure 1. Molecular Structure of [Pt(PPh₃)(Eind₂-BPEP)].

Unsymmetrical PNP-Pincer Type Phosphaalkene Ligands Protected by a Fused-Ring Bulky Eind Group: Synthesis and Applications to Rh(I) and Ir(I) Complexes

Unsymmetrical PNP-pincer type phosphaalkene ligand 2-(Phospholanylmethyl)-6-(2-phosphaethenyl)pyridine (Eind-PPEP) protected by an octaethyl-s-hydrindacen-4-yl (Eind) group was synthesized. The fused-ring bulky Eind





Figure 2. Non-innocent PNP-pincer type phosphaalkene ligand protected by a fused-bulky Eind group.

group successfully prevents the loss of the P=C bond and enables us to compare the reactivity of Rh(I) and Ir(I) complexes toward ammonia. The complex K[RhCl(Eind-PPEP*)], bearing a dearomatized Eind-PPEP* ligand, undergoes simple ligand displacement to give [Rh(NH₃) (Eind-PPEP*)], whereas the iridium analogue K[IrCl(Eind-PPEP*)] causes N-H bond cleavage to form [Ir(NH₂) (Eind-PPEP)]. DFT calculations indicate a thermodynamic cause of the metal-dependent product change.

Mixed-Ligand Approach to Palladium-Catalyzed Direct Arylation Polymerization: Effective Prevention of Structural Defects Using Diamines

We found a novel mixed ligand catalyst for palladium-catalyzed direct arylation polymerization (DArP) of 2,6-diiododithienosilole (DTS- I_2) and thienopyrroledione (TPD- H_2) to give poly(DTS-alt-TPD). It has been documented that this monomer combination has a marked tendency to form homocoupling and branching defects in polymer chains, and the latter defects eventually lead to the formation of insoluble materials. In this study, we demonstrated that the combined use of $P(o\text{-MeOC}_6H_4)_3$ and TMEDA ligands effectively prevents the defect formation. The side reactions that afford structural defects constitute a sequential process triggered by the reduction of DTS-I units. TMEDA as a simple diamine effectively inhibits the reduction of DTS-I units, giving poly(DTS-alt-TPD) ($M_n^{GPC} = 20,000$) in high cross-coupling selectivity (99%).



Scheme 1.

International Research Center for Elements Science - Nanophotonics -

http://www.scl.kyoto-u.ac.jp/~opt-nano/index-e.html



Prof KANEMITSU, Yoshihiko (D Eng)



Assist Prof IHARA, Toshiyuki (D Sc)



Assist Prof TAHARA, Hirokazu (D Sc)



PD TEX, David Michael (D Eng)



PD LE, Phuong Quang (D Sc)

Students


YAMADA, Takumi (D1) HIROSHIGE, Nao (M2) HANDA, Taketo (M2) YARITA, Naoki (M1)

Scope of Research

Our research interest is to understand optical and quantum properties of nanometer-structured materials and to establish opto-nanoscience for creation of innovative functional materials. Space- and time-resolved laser spectroscopy is used to study optical properties of semiconductor quantum nanostructures and strongly correlated electron systems in low-dimensional materials. The main subjects are as follows: 1) investigation of optical properties of single nanostructures through the development of a high-resolution optical microscope, 2) development of nanoparticle assemblies with new optical functionalities, and 3) ultrafast optical spectroscopy of excited states of semiconductor nanostructures.

KEYWORDS

Femtosecond Laser Spectroscopy Single Photon Spectroscopy Semiconductor Nanoparticles Solar Cells Perovskites



Selected Publications

Yamashita, D.; Handa, T.; Ihara, T.; Tahara, H.; Shimazaki, A.; Wakamiya, A.; Kanemitsu, Y., Charge Injection at the Heterointerface in Perovskite CH₃NH₃PbI₃ Solar Cells Studied by Simultaneous Microscopic Photoluminescence and Photocurrent Imaging Spectroscopy, *J. Phys. Chem. Lett.*, 7, 3186-3191 (2016).

Yamada, T.; Yamada, Y.; Nishimura, H.; Nakaike, Y.; Wakamiya, A.; Murata, Y.; Kanemitsu, Y., Fast Free-Carrier Diffusion in CH₃NH₃PbBr₃ Single Crystals Revealed by Time-Resolved One- and Two-Photon Excitation Photoluminescence Spectroscopy, *Adv. Electron. Mater.*, **2**, [1500290-1]-[1500290-5] (2016).

Ihara, T.; Kanemitsu, Y., Absorption Cross-section Spectrum of Single CdSe/ZnS Nanocrystals Revealed through Photoluminescence Excitation Spectroscopy, *Phys. Rev. B*, **92**, [155311/1]-[155311/5] (2015).

Yamada, Y.; Yamada, T.; Le, P. Q.; Maruyama, N.; Nishimura, H.; Wakamiya, A.; Murata, Y.; Kanemitsu, Y., Dynamic Optical Properties of CH₃NH₃PbI₃ Single Crystals As Revealed by One- and Two-photon Excited Photoluminescence Measurements, *J. Am. Chem. Soc.*, **137**, 10456-10459 (2015).

Yamada, Y.; Nakamura, T.; Endo, M.; Wakamiya, A.; Kanemitsu, Y., Photocarrier Recombination Dynamics in Perovskite CH₃NH₃PbI₃ for Solar Cell Applications, *J. Am. Chem. Soc.*, **136**, 11610-11613 (2014).

Biexciton Cascade Emission and Absorption Cross Section of Single Semiconductor Nanocrystals

The sequential two-photon emission process known as biexciton cascade emission is a characteristic phenomenon that occurs in photoexcited semiconductor nanocrystals (NCs). This process occurs when a biexciton state is created in the NCs; thus, the occurrence of the process is related to the photoabsorption properties of the NCs. We presented a simple equation that connects the photoabsorption of single NCs and the biexciton cascade emission. The equation is found to be independent of the quantum yields of photoluminescence (PL). With this equation and using an analysis of second-order photon correlation, the absolute absorption cross section of the single NCs could be evaluated, obtaining values on the order of $10^{-14} \, \mathrm{cm}^2$.




Figure 1. Schematic of the biexciton cascade emission process.

Microscopic Photoluminescence and Photocurrent Imaging of CH₃NH₃PbI₃ Perovskite Solar Cells

Metal halide perovskite semiconductors are attracting much attention owing to their high photovoltaic properties in solution-processed solar cells. Detailed understanding of carrier recombination and transport processes is needed to further improve the power conversion efficiency. We analyzed these processes in perovskite solar cells by microscopic imaging of time-resolved photoluminescence (TR-PL) and photocurrent (PC). Simultaneous measurements of TR-PL and PC show a positive correlation




Figure 2. Microscopic image of photoluminescence and photocurrent intensities in perovskite solar cell.

between PL intensity and PL lifetime, and a negative correlation between PL and PC intensities. We found that these correlations are caused by the competition between the photocarrier injection from the perovskite layer to the charge transport layer and the photocarrier recombination in the perovskite layer.

Free Carriers Versus Excitons in CH₃NH₃PbI₃ Perovskite Thin Films at Low Temperatures

Organometal halide perovskites are an excellent class of materials for developing cost-effective electronic devices including solar cells, light-emitting diodes, and lasers. Therefore, insightful knowledge of photocarrier recombination dynamics in perovskites, which determines fundamentally performance of perovskite-based devices, is important. We investigated the dynamic responses of photocarriers in CH₃NH₃PbI₃ (MAPbI₃) thin films at low temperatures using various time-resolved laser spectroscopy. The spectroscopic data consistently indicate that the high-temperature tetragonal phase remains in MAPbI₃ films. We observed a fast charge transfer from the major orthorhombic phase to the minor tetragonal phase, which likely prevents the formation of excitons in the orthorhombic phase. Thus, the photocarrier recombination dynamics in two phases could be described by a free-carrier model, rather than an exciton model, even at low temperatures.




Figure 3. Two-dimensional contour image of a time-resolved photoluminescence spectrum of MAPbI₃ thin films at 15 K.

Bioinformatics Center Chemical Life Science –

http://cls.kuicr.kyoto-u.ac.jp



Prof OGATA, Hiroyuki (D Sc)



Assoc Prof GOTO, Susumu (D Eng)



Assist Prof (D Sc)



Program-Specific Res BLANC-MATHIEU, Romain YOSHIZAWA, Akiyasu (D Sc)

Researchers(pt)

SHIMIZU, Yugo YAMAMOTO, Rumiko

Students

YOSHIKAWA, Genki (M2) NISHIYAMA, Hiroki (M2) LI, Yanze (M1) PRODINGER, Florian (RS) ARAMAKI, Takuya (UG)

KURONISHI, Megumi (UG) YAMASHITA, Shohei (UG) NURSHAHIRA, Binti Yusuf (UG) TUAN, Watie Binti Tuan Mat (UG)

Scope of Research

We are interested in understanding the functioning and evolution of biological systems at varying scales from tiny microbes up to the Earth's environment, by leveraging rapidly accumulating big data in life science and bioinformatics approaches. We currently focus on 1) the evolution of viruses and their links to the origin of life, 2) microbial ecology

in different ecosystems, and 3) the development of bioinformatics methods and biological knowledge resources for biomedical and industrial applications. To fuel these research activities, we take part in environmental sampling campaigns such as Tara Oceans. Our resources and developed tools are accessible through GenomeNet (www.genome.jp) to scientific communities and the public.



Bioinformatics GenomeNet **Environmental Genomics** Virology Molecular Evolution



Selected Publications

Blanc-Mathieu, R.; Ogata, H., DNA Repair Genes in the Megavirales Pangenome, Curr. Opin. Microbiol., 31, 94-100 (2016). Mihara, T. et al., Linking Virus Genomes with Host Taxonomy, Viruses, 8, 66 (2016).

Moriya, Y. et al., Identification of Enzyme Genes Using Chemical Structure Alignments of Substrate-product Pairs, J. Chem. Inf. Model., 56, 510-516 (2016).

Guidi, L. et al., Plankton Networks Driving Carbon Export in the Oligotrophic Ocean, Nature, 532, 465-470 (2016).

Diversity of the Giant Virus Family Megaviridae

Megaviridae is a proposed family of eukaryotic viruses classified in the group of nucleocytoplasmic large double-stranded DNA viruses (NCLDVs). They possess large genomes and atypical gene contents as compared to other viruses. Genomic data suggest that the diversity of Megaviridae is vast and perhaps greater than that of cellular organisms. In the present work, we aimed at characterizing the diversity of Megaviridae at a single site using high-throughput sequencing analysis of PCR-amplified DNA polymerase family B (PolB) gene.

PolB was chosen as a marker gene for PCR amplification. PolB is encoded in all available Megaviridae genome sequences. Its high level of divergence makes it impossible to design a single PCR primer pair that would amplify all Megaviridae PolB sequences observed in current metagenomic data. To overcome this limitation we designed 82 degenerate primer pairs that target a conserved domain of PolB based on 923 PolB sequences from isolated Megaviridae and environmental Megaviridae. *In silico* tests demonstrated that this set of primer pairs specifically covers 97.4% of the 923 PolB sequences.

We first experimentally tested a subset of 34 primer pairs on environmental DNA extracted from water samples collected at the sea surface of Osaka bay. PCR-products (*n*=34) were sequenced with Illumina MiSeq to produce paired-end reads. The reads were quality trimmed, merged and clustered by 100% identity threshold. Representative sequences of the clusters were then searched for homology against a reference database containing PolB sequences of cellular organisms and viruses. Then, PolB sequences from all primer pairs were pooled together and aligned. The alignment was trimmed so that the amplicon sequences were well aligned at their both ends. As a result, we obtained 3,400,247 high quality merged reads. All the PolB fragments were predicted to be of Megaviridae origin by a phylogenetic analysis using pplacer.

We clustered all PolB amplicon sequences into operational taxonomic units (OTUs) at 97% identity cutoff. The




Figure 1. Rarefaction curve for the number of OTUs. Singleton OTUs are excluded.

number of OTUs reached near saturation at 5,000 OTUs (Figure 1), suggesting that even a single sea water sample contains diverse viruses of Megaviridae.

A Characterization of the Genome of XacN1, a Jumbo Phage Infecting the Citrus Canker Agent Xanthomonas Citri

There is an increasing interest in the use of phages for controlling plant-pathogenic bacteria in agricultural activities. *Xanthomonas citri* is the bacterial pathogen of citrus canker disease, which is one of the serious citrus plant diseases and leads to significant economic damages worldwide. XacN1 is a bacteriophage that infects *X. citri*. Electron microscopic analysis revealed XacN1 is a large myovirus composed of an icosahedral head of 140 nm in diameter with a 145 nm contractile tail. XacN1 has the second largest double stranded DNA genome (384 kb) among sequenced phages. Its interesting structural feature is the presence of 65 kb direct repeats at the extremities of the linear genome, likely serving for genome replication.

We predicted 592 open reading frames (ORFs) in the XacN1 genome. Homology search revealed that the ORFs include important genes such as genes involved in replication, translation system, and those encoding structural proteins. The genome was found to encode a surprisingly large number of tRNA genes, i.e., 58 tRNAs corresponding to all the twenty amino acids and suppressor tRNAs. The tRNAs showed a significant tendency: codons highly used by the phage were less frequently used by the host bacterium and vice versa. Phylogenetic tree reconstructions of structural proteins indicate that four myoviruses and XacN1 with genome size >300 kb form a clade (Figure 2). Further analysis of the XacN1 genome will lead to uncover yet unidentified unique evolutionary and functional properties of giant phages.




Figure 2. Maximum likelihood phylogenetic tree of tail sheath proteins. Blue and green circles indicate phages with a genome greater than 300 kb and 200 kb, respectively.

Bioinformatics Center - Mathematical Bioinformatics -

http://www.bic.kyoto-u.ac.jp/takutsu/index.html



Prof AKUTSU, Tatsuya (D Eng)



Assist Prof HAYASHIDA, Morihiro (D Inf)



Assist Prof TAMURA, Takeyuki (D Inf)



SPD (JSPS) SIMONE, Marini (Ph D)



PD
JIRA, Jindalertudomdee
(D Inf)

Students

BAO, Yu (D3) FUKUSAKO, Yuta (M2) PI, Wenya (RS) LIU, Pengyu (D1) LI, Ruiming (M1)

CAO, Yu (RS)

Guest Scholars

KAWAKAMI, Yuko (M2)

VINCHENT, Alexandra University of Montpellier, France, 20 April–16 July HOU, Wenpin (Ph D) The University of Hong Kong, China, P.R., 23 May–22 August

Guest Res Assoc


YANG, Jinn-Moon (Ph D) National Chiao Tung University, Taiwan, 4 February–4 May CHENG, Xiaoqing (Ph D) The University of Hong Kong, China, P.R., 6 March–3 April

Scope of Research

Due to rapid progress of genome sequencing technology, whole genome sequences of organisms ranging from bacteria to human have become available. In order to understand the meaning behind the genetic code, we have been developing algorithms and software tools for analyzing biological data based on advanced information technologies such as theory of algorithms, artificial intelligence, and machine learning. We are currently studying the following topics: systems biology, scale-free networks, protein structure prediction, inference of biological networks, chemo-informatics, and discrete and stochastic methods for bioinformatics.

KEYWORDS

Scale-free Networks Boolean Networks Chemical Graphs Grammar-based Compression Protein Complexes





Selected Publications

Hou, W.; Tamura, T.; Ching, W. K.; Akutsu, T., Finding and Analyzing the Minimum Set of Driver Nodes in Control of Boolean Networks, *Advances in Complex Systems*, **19**, 1650006 (2016).

Takemoto, K.; Akutsu, T., Analysis of the Effect of Degree Correlation on the Size of Minimum Dominating Sets in Complex Networks, *PLoS ONE*, **11**, e0157868 (2016).

Jindalertudomdee, J.; Hayashida, M.; Akutsu, T., Enumeration Method for Structural Isomers Containing User-defined Structures Based on Breadth-first Search Approach, *J. Comput. Biol.*, **23**, 625-640 (2016).

Ishitsuka, M.; Akutsu, T.; Nacher, J. C., Critical Controllability in Proteome-wide Protein Interaction Network Integrating Transcriptome, *Sci. Rep.*, **6**, 23541 (2016).

Enumeration Method for Structural Isomers Containing User-defined Structures Based on Breadth-first Search Approach

Enumeration of chemical structures is useful for discovering new compounds and drugs, and elucidating chemical structures from mass spectrometry. We previously developed efficient algorithms, BfsSimEnum, BfsMulEnum, BfsBenNaphEnum for enumerating tree-like chemical compounds without and with multiple bonds, and compounds containing rings such as benzene and naphthalene, respectively. For many instances, the algorithms were able to enumerate chemical structures faster than other existing methods.

In this study, we propose a novel efficient enumeration algorithm, BfsStructEnum, which allows users to define desired cyclic structures and enumerates all nonredundant chemical compounds containing only user-defined structures as cyclic structures from a given chemical formula.

For evaluating the performance, we confirmed that the number of enumerated structures of BfsStructEnum was the same as that of an existing method, MOLGEN for several instances. Furthermore, we also found that BfsStructEnum was significantly faster than MOLGEN. It implies that compression of cyclic structures into single nodes makes the enumeration more efficiently.




Figure 1. Example of the assignment process of atom position lists to pyridine nodes (denoted by p1 and p2).

Finding Influential Genes using Gene Expression Data and Boolean Models of Metabolic Networks

Selection of influential genes using gene expression data from normal and disease samples is an important topic in bioinformatics. In this research, we propose a novel computational method for the problem, which combines gene expression patterns from normal and disease samples with a mathematical model of metabolic networks. This method seeks a set of k genes knockout of which drives the state of the metabolic network towards that in the disease samples. We adopt a Boolean model of metabolic networks and formulate the problem as a maximization problem under an integer linear programming framework. We applied the proposed method to selection of influential genes using gene expression data from normal samples and disease (head and neck cancer) samples. The result suggests that the proposed method can select more biologically relevant genes than an existing P -value based ranking method can.




Figure 2. In Boolean metabolic networks, rectangles and circles represent reactions and metabolites, respectively. Each node is assigned either 0 or 1 to represent the status of the node.

Bioinformatics Center - Bio-knowledge Engineering -

http://www.bic.kyoto-u.ac.jp/pathway/index.html



Prof MAMITSUKA, Hiroshi NGUYEN, Canh Hao (D Sc)



Assist Prof (D Knowledge Science)



Assist Prof YAMADA, Makoto (D Statistical Science)



Program-Specific Res WIMALAWARNE, Kishan (D Eng)

Students

TOHZAKI, Yudai (M1)

KIRIHATA, Tatsuro (UG)


Scope of Research

We are interested in graphs and networks in biology, chemistry, and medical sciences, including metabolic networks, protein-protein interactions and chemical compounds. We have developed original techniques in machine learning and data mining for analyzing these graphs and networks, occasionally combining with table-format datasets, such as gene expression and chemical properties. We have applied the techniques developed to real data to demonstrate the performance of the methods and find new scientific insights.

KEYWORDS

Bioinformatics Computational Genomics Data Mining Machine Learning Systems Biology

Bio-knowledge Engineering



Selected Publications

Gao, J.; Yamada, M.; Kaski, S.; Mamitsuka, H.; Zhu, S., A Robust Convex Formulations for Ensemble Clustering, Proceedings of the 25th International Joint Conference on Artificial Intelligence (IJCAI 2016), 1476-1482 (2016).

Peng, S.; You, R.; Wang, H.; Zhai, C.; Mamitsuka, H.; Zhu, S., DeepMeSH: Deep Semantic Representation for Improving Large-scale MeSH Indexing, Bioinformatics 32 (12) (Proceedings of the 24th International Conference on Intelligent Systems for Molecular Biology (ISMB 2016)), i70-i79 (2016).

Yuan, Q.-J.; Gao, J.; Wu, D.; Zhang, S.; Mamitsuka, H.; Zhu, S., DrugE-Rank: Improving Drug-Target Interaction Prediction of New Candidate Drugs or Targets by Ensemble Learning to Rank, Bioinformatics 32 (12) (Proceedings of the 24th International Conference on Intelligent Systems for Molecular Biology (ISMB 2016)), i18-i27 (2016).

Mohamed, A.; Nguyen, C. H.; Mamitsuka, H., NMRPro: An Integrated Web Component for Interactive Processing and Visualization of NMR Spectra, Bioinformatics, 32, 2067-2068 (2016).

Shinkai-Ouchi, F.; Koyama, S.; Ono, Y.; Hata, S.; Ojima, K.; Shindo, M.; duVerle, D.; Kitamura, F.; Doi, N.; Takigawa, I.; Mamitsuka, H.; Sorimachi, H., Predictions of Cleavability of Calpain Proteolysis by Quantitative Structure-Activity Relationship Analysis Using Newly Determined Cleavage Sites and Catalytic Efficiencies of an Oligopeptide Array, Mol. Cell. Proteomics, 15, 1262-1280 (2016).

Nguyen, C. H.; Mamitsuka, H., New Resistance Distances with Global Information on Large Graphs, Proceedings of the Nineteenth International Conference on Artificial Intelligence and Statistics (AISTATS 2016) (JMLR Workshop and Conference Proceedings, 51), 639-647 (2016).

Ultra High-Dimensional Nonlinear Feature Selection for Big Biological Data

Life sciences are going through a revolution thanks to the possibility to collect and learn from massive biological data. Efficient processing of such "big data" is extremely important for many medical and biological applications, including disease classification, biomarker discovery, and drug development. The complexity of biological data is dramatically increasing due to improvements in measuring devices such as next-generation sequencers, microarrays and mass spectrometers. As a result, we must deal with data that includes many observations (hundreds to tens of thousands) and even larger numbers of features (thousands to millions). Machine learning algorithms are charged with learning patterns and extracting actionable information from biological data. These techniques have been used successfully in various analytical tasks, such as genome-wide association studies and gene selection.

However, the scale and complexity of big biological data pose new challenges to existing machine learning algorithms. There is a trade-off between scalability and complexity: linear methods scale better to large data, but cannot model complex patterns. Nonlinear models can handle complex relationships in the data but are not scalable to the size of current datasets. In particular, learning nonlinear models requires a number of observations that grows exponentially with the number of features. Biological data generated by modern technology has as many as millions of features, making the learning of nonlinear models




Figure 1. Results for the enzyme dataset. Circles indicate best accuracy/independence rate according to t-tests (\$p<0.05\$). Differences in accuracy are significant for m=5,20,30,40. (B)~Independence rate vs. m: all differences are significant.

unfeasible with existing techniques. To make matters worse, current nonlinear approaches cannot take advantage of distributed computing platforms.

A promising approach to make nonlinear analysis of big biological data computationally tractable is to reduce the number of features. This method is called feature selection. Biological data is often represented by matrices where rows denote features and columns denote observations. Feature selection aims to identify a subset of features (rows) to be preserved, while eliminating all others.

Here we propose a novel feature selection framework for big biological data that makes it possible for the first time to identify very few relevant, non-redundant features among millions. The proposed method is based on two components: Least Angle Regression (LARS), an efficient feature selection method, and the Hilbert-Schmidt Independence Criterion (HSIC), which enables the selection of features that are non-linearly related. These properties are combined to obtain a method that can exploit nonlinear feature dependencies efficiently, and furthermore enables distributed implementation on commodity cloud computing platforms. We name our algorithm Least Angle Nonlinear Distributed (LAND) feature selection. Through experiment on a large and high-dimensional dataset (million features with tens of thousand samples), we show that our approach can find a set of independent features without losing classification accuracy (Figure 1 A and B). Moreover, the result is achieved in hours of cluster computing time (Figure 1 C). The selected features are relevant and non-redundant, making it possible to obtain accurate and interpretable models that can be run on a laptop computer.




Figure 2. Computation time of feature selection algorithms. We ran the algorithms on reduced-dimensionality datasets. We estimated the mRMR runtime for d> 50,000, given that it scales linearly with d (dotted line).

HAKUBI RESEARCHERS' ACTIVITIES IN ICR

Hakubi Project: Fosterage and Support of Young Researchers, Kyoto University

Research Topic

Algorithmic Graph Theory with Applications to Bioinformatics



Program-Specific Assoc Prof JANSSON, Jesper (Ph D)

Host Laboratory Laboratory of Mathematical Bioinformatics Host Professor AKUTSU, Tatsuya

Outline of Research

This year, my research collaborators and I have continued our work on efficient algorithms for computing different types of consensus trees. Given an input consisting of k phylogenetic trees with n leaves each and with identical leaf label sets, the running times of our algorithms are:

- O(k n) time [majority rule consensus tree],
- O(k n) time [loose consensus tree],
- O(k n²) time [greedy consensus tree],
- $\min\{O(k n^2), O(k n (k + \log^{2} n))\}\$ time [frequency difference consensus tree],
- O(k n log n) [Adams consensus tree],
- $O(n^2)$ [R* consensus tree with k=2],
- $O(n^2 \log^{4/3} n)$ [R* consensus tree with k=3],
- $O(n^2 \log^{k+2} n) [R^*$ consensus tree with k>3],
- $O(k n^3 + 2.733^n)$ [minimally resolved local consensus tree], and
- $O(k n^3 + 4^n poly(n))$ [minimally rooted-triplet-inducing consensus tree].

ACTIVITIES OF JOINT USAGE/RESEARCH CENTER

JURC Cooperative Research Subjects 2016

 $(1 \text{ April } 2016 \sim 31 \text{ March } 2017)$

STARTING-UP SUBJECTS (IN SPECIFIC FIELDS CHOSEN BY JURC)

A Study of Magneto-optic and Electro-optic Effects by Laser Induced Pico-second Electromagnetic Pulses

YASUHARA, Ryo, National Institutes of Natural Sciences, National Institute for Fusion Science

Host in JURC SAKABE, Shuji

Diagnostics of Li-Ion Batteries with Laser-Accelerated Protons KATO, Yoshiaki, The Graduate School for the Cretion of New Photonics Industries

Host in JURC SAKABE, Shuji

Fundamental Study on Micro-fabrication of Metal with Controlling Laser Absorption

KUSABA, Mitsuhiro, Department of Electronics, Information and Communication Engineering, Osaka Sangyo University

Host in JURC HASHIDA, Masaki

Feasibility Study of Deuteron Beam for Compact Neutron Sources HIROTA, Katsuya, Graduate School of Science, Nagoya University **Host in JURC** IWASHITA, Yoshihisa

Development of Neutron Resonance Spin Flipper at BL06(VIN ROSE) at J-PARC/MLF

HINO, Masahiro, Research Reactor Institute, Kyoto University **Host in JURC** IWASHITA, Yoshihisa

Fundamental Research on Compact Neutron Source with Permanent Magnet

OHNUMA, Masato, Division of Quantum Science and Engineering Faculty of Engineering, Hokkaido University

Host in JURC IWASHITA, Yoshihisa

Synthesis of Transition Metal Complexes with a Multidentate Phosphaalkene Ligand and Their Application to Catalytic Reactions MATSUO, Tsukasa, Faculty of Science and Engineering / Graduate School of Science and Engineering Research, Kindai University Host in JURC TAKEUCHI, Katsuhiko

Development of a Novel 3d Metal Catalyst Bearing a M-E Bond as a Key Active Site

NAKAJIMA, Yumiko, Interdisciplinary Research Center for Catalytic Chemistry, National Institute of Advanced Industrial Science and Technology

Host in JURC OZAWA, Fumiyuki

Dielectric Function and Excitonic Effect of Dynamical Screening in Semiconducting Materials for Photoelectric Conversion

SUZUURA, Hidekatsu, Graduate School of Engineering, Hokkaido University

Host in JURC KANEMITSU, Yoshihiko

I : International Joint Research

F: Female PI

Characterization of Self-assembled Peptides on Two-dimensional Semiconductor Nanomaterials by Microscopic Photoluminescence Lifetime Measurements

HAYAMIZU, Yuhei, Graduate School of Engineering, Tokyo Institute of Technology

Host in JURC KANEMITSU, Yoshihiko

Investigation of Photocarrier Characteristics in Halide-perovskite-based Photodevices

YAMADA, Yasuhiro, Department of Physics, Graduate School of Science, Chiba University

Host in JURC KANEMITSU, Yoshihiko

Development of Synthetic Organic Reactions Mediated by Low-valent Fe Complexes of N-heterocyclic Carbenes

OHKI, Yasuhiro, Department of Chemistry, Graduate School of Science, Nagoya University

Host in JURC NAKAMURA, Masaharu

Development of Transition metal Complexes Based on Controlled Coordination Ability of NHC

YAMAGUCHI, Yoshitaka, Faculty of Engineering, Division of Materials Science and Chemical Engineering, Yokohama National University

Host in JURC NAKAMURA, Masaharu

Precise Separation and Characterization of Novel Thiolate Protected Metal Clusters by Using LC/MS

NEGISHI, Yuichi, Faculty of Science, Department of Applied Chemistry, Tokyo University of Science

Host in JURC TAKAYA, Hikaru

Analysis and Control of Complex Bipartite Networks JOSE, C. Nacher, Department of Science, Toho University **Host in JURC** AKUTSU, Tatsuya

Flexible and Efficient Search Algorithms for Structured Data in

TAKASU, Atsuhiro, Division of Digital Content and Media Sciences Research, National Institute of Informatics

Host in JURC AKUTSU, Tatsuya

Pathway Database for Human Gut Microbiome and the Aapplication Toward Metagenomics

YAMADA, Takuji, Graduate School of Bioscience and Biotechnology, Tokyo Institute of Technology

Host in JURC GOTO, Susumu

Biological Molecular Interaction Data Analysis Using Auxiliary Information

SHIGA, Motoki, Informatics Course, Department of Electrical, Electronic and Computer Engineering, Faculty of Engineering, Gifu University

Host in JURC MAMITSUKA, Hiroshi

Total Synthesis of Dendrochrysanene

KURAMOCHI, Koji, Graduate School of Life and Environmental Sciences, Kyoto Prefectural University

Host in JURC KAWABATA, Takeo

Polymer Nano-architectonics for Bridging a Gap between Macroscopic Interfacial Phenomena and Molecular Function NAKANISHI, Waka, MANA, National Institute for Materials

Host in JURC SAKAKIBARA, Keita

Precursor Approach towards Controlled Preparation of Organic Heterojunction Layers

SUZUKI, Mitsuharu, Graduate School of Materials Science, Nara Institute of Science and Technology (NAIST)

Host in JURC MURATA, Yasujiro

Development of Novel Nonalternant Heterocycles toward Electronic Materials

KUROTOBI, Kei, National Institute of Technology, Kurume College Host in JURC MURATA, Yasujiro

Study on Transportation of Metal Ions through a Liquid Membrane Aiming at Highly Efficient Mutual Separation of Metal Ions MUKAI, Hiroshi, Faculty of Education, Kyoto University of Education

Host in JURC SOHRIN, Yoshiki

Simulation of Photoacoustic Signals Generated by Gold Nanoparticles OKAWA, Shinpei, National Defense Medical College Host in JURC TERANISHI, Toshiharu

Kinetic Analysis of Hydrogen Absorption Process Observed on Shape-controlled Pd Nanoparticles

YAMAUCHI, Miho, International Institute for Carbon-Neutral Energy Research, Kyushu University

Host in JURC TERANISHI, Toshiharu

Preparation and Evaluation of Colorful Dve-Sensitized Solar Cell Devices Using Designed Anthocyanins and Their Theoretical Studies toward Improvement of the Efficiency

YOSHIDA, Kumi, Department of Complex Systems Science, Graduate School of Information Science, Nagoya University F

Host in JURC MURATA, Yasujiro; WAKAMIYA, Atsushi

Test of Soret Effect via Plasmon Heating of Periodic Metal Domain Structure

SHIMADA, Ryoko, Department of Mathematical and Physical Sciences, Faculty of Science, Japan Women's University F

Host in JURC WATANABE, Hiroshi

EXPANDING SUBJECTS (IN SPECIFIC FIELDS CHOSEN BY JURC)

Intense THz Emission for Nonlinear Interaction Physics NAGASHIMA, Takeshi, Faculty of Science and Engineering, Setsunan University

Host in JURC HASHIDA, Masaki

Study for High-repetition Electron/Ion Generation with High Power

ARIKAWA, Yasunobu, Institute of Laser Engineering, Osaka University

Host in JURC INOUE, Shunsuke

Research and Development on Future Accelerator toward ILC project HAYANO, Hitoshi, Accelerator Laboratory, High Energy Accelerator Research Organization

Host in JURC IWASHITA, Yoshihisa

Study on Magnification of the Pulsed-neutron Transmission Image Using the Sextupole Magnet, Aimed at Visualization of Charge and Discharge in the Electrode Materials of Li-ion Batteries

KINO, Koichi, Research Institute for Measurement and Analytical Instrumentation, National Institute of Advanced Industrial Science and Technology

Host in JURC IWASHITA, Yoshihisa

X-ray Structural Biology Study of Flavin Reductase Involved in Resorcinol Catabolism

OIKAWA, Tadao, Faculty of Chemistry, Materials and Bioengineering, Kansai University

Host in JURC FUJII, Tomomi

X-ray Crystallographic Studies on Thermostability and Substrate Specificity of L-Asparaginase

KATO, Shiro, International Institute of Rare Sugar Research and Education, Kagawa University

Host in JURC FUJII, Tomomi

Construction and Functionality of Phosphasilene and Silanethione at Transition-metals

OKAZAKI, Masaaki, Graduate School of Science and Technology, Hirosaki University

Host in JURC OZAWA, Fumiyuki

Mechanistic Studies of C-H Bond Functionalization Reactions Using Iron and Related Base Metal Catalysts

YOSHIKAI, Naohiko, Division of Chemistry and Biological Chemistry, School of Physical and Mathematical Sciences, Nanyang Technological University

Host in JURC NAKAMURA, Masaharu

Development of Cooperative Asymmetric Pincer-Fe Catalyst ARAI, Takayoshi, Department of Chemistry, Graduate School of Science, Chiba University

Host in JURC NAKAMURA, Masaharu

Elucidation of Electronic Structures of Cycloparaphenylenes and Their Application to Materials Science

UCHIYAMA, Masanobu, Graduate School of Pharmaceutical Science, The University of Tokyo

Host in JURC YAMAGO, Shigeru

Synthesis of Cyclic π -conjugated Molecules and Their Properties SUZUKI, Toshiyasu, Institute for Molecular Science, National Institute of Natural Sciences

Host in JURC YAMAGO, Shigeru

MiSeq Amplicon Analysis of Megaviridae DNA Polymerase YOSHIDA, Takashi, Graduate School of Agriculture, Kyoto University

Host in JURC OGATA, Hiroyuki

Comparative Population Genomics of Parmales KUWATA, Akira, Tohoku National Fisheries Research Institute Host in JURC OGATA, Hiroyuki

Ι

Molecular Network Analysis by Combinations of Machine Learning and Statistics

KAYANO, Mitsunori, Research Center for Global Agromedicine, Obihiro University of Agriculture and Veterinary Medicine

Host in JURC MAMITSUKA, Hiroshi

Colloidal Microparticles with Polymer Brushes for Laser Applications FURUMI, Seiichi, Faculty of Science Division 1, Tokyo University of Science

Host in JURC OHNO, Kohji

Organic Photovoltaic Devices Composed of Novel Fluorine-Containing Donor Polymer and Fullerene Derivatives

IE, Yutaka, The Institute of Scientific and Industrial Research, Osaka University

Host in JURC MURATA, Yasujiro

Exploration of Novel Charge Transport Materials for Perovskite Solar Cell via Direct Evaluation of Charge Transfer Process SAEKI, Akinori, Graduate School of Engineering, Osaka University **Host in JURC** WAKAMIYA, Atsushi

A Study on Open-shell Molecules in Cyclophenylene Structures ABE, Manabu, Graduate School of Science, Hiroshima University **Host in JURC** YAMAGO, Shigeru

A Study of Roll-off Mechanisms of External Quantum Efficiency of Organic Light-emitting Diodes Based on Thermally-assisted Delayed Fluorescent Materials

NAITO, Hiroyoshi, Department of Electronics, University of Osaka Prefecture

Host in JURC KAJI, Hironori

Making of the Vertical Cross Section of Bioactive Trace Metals in the East China Sea

NAKAGUCHI, Yuzuru, Faculty of Science and Engineering, Kindai University

Host in JURC SOHRIN, Yoshiki

Structural Analysis of Mixed Oxide Nanoparticles Protected by Organic Compounds and Their Magnetic Properties

HARADA, Masafumi, Faculty of Human Life and Environment, Nara Women's University

Host in JURC TERANISHI, Toshiharu

Examination of Randomness of Lithium Silicate Glasses using Lightscattering Spectroscopy

KOREEDA, Akitoshi, Department of Physical Science, College of Science and Engineering, Ritsumeikan University

Host in JURC MASAI, Hirokazu

Stress-Optical Law for Polymeric Liquids under High Shear MASUBUCHI, Yuichi, National Composite Center, Nagoya University

Host in JURC MATSUMIYA, Yumi

Dynamics of Entanglements between Polymer Networks by Contrast Variation Optical-rheometry

KATASHIMA, Takuya, Graduate School of Science, Osaka University

Host in JURC MATSUMIYA, Yumi

STARTING-UP SUBJECTS (ON-DEMAND FROM RELATED COMMUNITIES)

Study on Lipid Secretion Pathways of Plant Cells

YAZAKI, Kazufumi, Research Institute for Sustainable Humanosphere, Kyoto University

Host in JURC AOYAMA, Takashi

Study on the Regulatory Mechanism of Plant Epidermal Cell Differentiation

TOMINAGA, Rumi, Graduate School of Biosphere Science, Hiroshima University

F

Host in JURC AOYAMA, Takashi

Role of PIP5K and ROP-GTPase in Plant Cell Morphogenesis KUSANO, Hiroaki, Department of Technology and Science, Tokyo University of Science

Host in JURC AOYAMA, Takashi

Chemical Biology of Endogenous Vitamin D Derivatives KITTAKA, Atsushi, Faculty of Pharma Sciences, Teikyo University **Host in JURC** UESUGI, Motonari

Effect of Dzyaloshinskii-Moriya-Interaction on Domain Wall Motion NAKATANI, Yoshinobu, Department of Communication Engineering and Informatics, The University of Electro-Communications

Host in JURC ONO, Teruo

Study of Tunnel Magnetoresistance Effect of the Magnetic Insulator Films with Perpendicular Magnetic Anisotropy for Development of High Spin Polarized Current Source

TANAKA, Masaaki, Department of Engineering Physics, Electronics and Mechanics, Nagoya Institute of Technology

Host in JURC ONO, Teruo

Selective Chemical Modification of Biomolecules in Membrane by Functionalized Catalysts

KUNISHIMA, Munetaka, Faculty of Pharmaceutical Sciences, Institute of Medical, Pharmaceutical, and Health Sciences, Kanazawa University

Host in JURC KAWABATA, Takeo

Characteristics of Membrane Vesicles Produced by Intestinal Bacteria and Their Biogenesis

KURATA, Atsushi, Faculty of Agriculture, Kindai University **Host in JURC** KURIHARA, Tatsuo

Precise Analysis for Brill Transition of Polyurea

MATSUBA, Go, Graduate School of Science and Engineering, Yamagata University

Host in JURC NISHIDA, Koji

Physical and Structural Properties of Glass-ceramics Fabricated at High Temperatures and High Pressures

MASUNO, Atsunobu, Institute of Industrial Science, The University of Tokyo

Host in JURC SHIMAKAWA, Yuichi

Solvent Extraction for Metal Ion Using Synergistic Effect of Surfactants

KURAHASHI, Kensuke, Environmental and Materials Chemistry Course, Osaka Prefecture University College of Technology Host in JURC SOHRIN, Yoshiki Development of Functional Materials Highly Selective for the Separation of Rare Metals

YAMAZAKI, Shoko, Department of Chemistry, Nara University of Education

Host in JURC UMETANI, Shigeo

Au Nanoparticle Single-electron Transistors by Using Four-legs Sb-porphyrin and Three-legs Phenol as Ligand and Anchor and Vice Versa

MAJIMA, Yutaka, Materials and Structures Laboratory, Tokyo Institute of Technology

Host in JURC TERANISHI, Toshiharu

Molecular Structure Determination by Combination of Spectroscopy and Chemometrics

SHIMADA, Toru, Faculty of Education, Hirosaki University Host in JURC HASEGAWA, Takeshi

In-situ Observation of Interfacial Chemical Reaction Using Polarization Modulation Infrared Reflection Adsorption Spectroscopy HAMA, Tetsuya, Institute of Low Temperature Science, Hokkaido University

Host in JURC HASEGAWA, Takeshi

Development of a Prediction Method Using Cell Penetrating Peptides for Anticancer Effects on a New Anticancer Drug "PARP inhibitors" OKUDA, Akiko, Graduate School of Health Sciences, Niigata University

Host in JURC FUTAKI, Shiroh

Construction of Novel Cell-penetrating Peptides that Have Retentivity in Blood

KUWATA, Keiko, Institute of Transformative Bio-Molecules, Nagoya University

Host in JURC FUTAKI, Shiroh

Cell Penetrating Peptide (CPP)-based Material Delivery System into Colon Cancer Stem Cells

OHASHI, Wakana, Graduate School of Medicine and Pharmaceutical Sciences for Research, University of Toyama F

Host in JURC FUTAKI, Shiroh

Preparation of Sm-doped Amorphous Materials for Radiophotoluminescence Applications

OKADA, Go, Graduate School of Materials Science, Nara Institute of Science and Technology

Host in JURC MASAI, Hirokazu

Creation of Organic Functional Materials Based on Nitrogencontaining Macrocyclic Compounds

IWANAGA, Tetsuo, Department of Chemistry, Faculty of Science, Okayama University of Science

Host in JURC MURATA, Yasujiro

Viscoelastic Relaxation of Linear and Ring Rouse Chains at Association/Dissociation Equilibrium

KWON, Youngdon, School of Chemical Engineering, Sungkyunkwan University

Host in JURC MATSUMIYA, Yumi

Preparation and Mechanical Properties of Noncovalent-Bonded Soft Elastomers

NORO, Atsushi, Graduate School of Engineering, Nagoya University Host in JURC WATANABE, Hiroshi

Synthesis of Metal Complexes Bearing Chiral Multidentate Carboxylic Acid Ligands and Application to Catalytic Reactions SUGIURA, Masaharu, Faculty of Life Sciences, Kumamoto University

Host in JURC NAKAMURA, Masaharu

F

EXPANDING SUBJECTS (ON-DEMAND FROM RELATED COMMUNITIES)

Prevention of NASH-induced Cancer Progression by SREBP Inhibitors

NAKAGAWA, Hayato, The University of Tokyo Hospital Host in JURC UESUGI, Motonari

Developments of Novel Spintornics Materials with Ferrimagnetic Spinel Ultrathin Films

NAGAHAMA, Taro, Laboratory of Advanced Materials Chemistry, Graduate School of Engineering, Hokkaido University

Host in JURC ONO, Teruo

Electric Field Induced Magnetic Domain Wall Motion CHIBA, Daichi, School of Engineering, The University of Tokyo Host in JURC ONO, Teruo

Relationship between Molecular Motions and Triplet Lifetime in Amorphous Materials

HIRATA, Shuzo, Department of Materials Science and Engineering, Tokyo Institute of Technology

Host in JURC KAJI, Hironori

Theoretical Study on Chemoselective Acylation Catalyzed by 4-Pyrrolidinopyridine Derivatives

YAMANAKA, Masahiro, Department of Chemistry, College of Science, Rikkyo University

Host in JURC KAWABATA, Takeo

Functional Analysis of the Bound D-Amino Acids in Food Proteins OHMORI, Taketo, Department of Biomedical Engineering, Osaka Institute of Technology

Host in JURC KURIHARA, Tatsuo

Functional Studies on Proteins Involved in the Microbial Metal Respiration

MIHARA, Hisaaki, Department of Biotechnology, College of Life Sciences, Ritsumeikan University

Host in JURC KURIHARA, Tatsuo

Search for Four Wave-mixing in the Vacuum

HONMA, Kensuke, Graduate School of Science, Hiroshima University

Host in JURC SAKABE, Shuji

Exploration of Novel Functional Properties of Transition Metal Oxides Using Ionic Liquid Gating

HATANO, Takafumi, Department of Crystalline Materials Science, Nagoya University

Host in JURC KAN, Daisuke

Ι

Research on Physical Properties and Functions of Complex Iron Oxide Thin Films

FUJII, Tatsuo, Graduate School of Natural Science and Technology, Okayama University

Host in JURC SAITO, Takashi

Control of Chromophore Orientation by Means of Dipole-Arrays of Perfluoroalkyl Amphiphile

YAMADA, Norihiro, Graduate School of Education, Chiba University

Host in JURC HASEGAWA, Takeshi

Toward Long Spin Coherence Time of NV Center by High Quality Diamond Growth

TOKUDA, Norio, Faculty of Electrical and Computer Engineering, Institute of Science and Engineering, Kanazawa University

Host in JURC MIZUOCHI, Norikazu

Electrical Control of Spin of NV Center

YAMASAKI, Satoshi, Power Electronics Research Center, National Institute of Advanced Industrial Science and Technology

Host in JURC MIZUOCHI, Norikazu

Development of Nanoarchitectures Formed from Composites of Magnetic Metal Complexes with Diblock Copolypeptitde Amphiphiles and Their Application for Magnetic Probes KUROIWA, Keita, Department of Nanoscience, Sojo University Host in JURC SASAMORI, Takahiro

Structural Analysis of Chiral Lithium Binaphtholate Complex and Development of Novel Catalysts

NAKAJIMA, Makoto, Faculty of Life Sciences, Kumamoto University

Host in JURC KAWABATA, Takeo

Mechanistic Studies of Hydrogen Transfer Ability of Aliphatic Amines and Application to Related Reactions

KOTANI, Shunsuke, Priority Organization for Innovation and Excellence, Kumamoto University

Host in JURC FURUTA, Takumi

SUBJECTS FOCUSING OF JOINT USAGE OF JURC/ICR **FACILITIES**

Correlation between Melting Point and Crystalline Structure of Ionic Liquid-polymer Gel Electrolyte Dependent on Concentration of Lithium Salt

OSAKA, Noboru, Department of Chemistry, Graduate School of Science, Okayama University of Science

Host in JURC TOSAKA, Masatoshi

Synthesis and Elucidation of Properties of Unsymmetrically-Substituted Disilyne and Related π -Electron Systems

IWAMOTO, Takeaki, Graduate School of Science, Tohoku University Host in JURC TOKITOH, Norihiro

Experimental Electron Density Distribution Analysis of Organogermanium Compounds

HASHIZUME, Daisuke, Center for Emergent Matter Science, RIKEN

Host in JURC TOKITOH, Norihiro

Synthesis and Structural Elucidation of Unsaturated Compounds of Group 14 Elements

MATSUO, Tsukasa, Faculty of Science and Engineering, Kindai University

Host in JURC TOKITOH, Norihiro

Synthesis and Structure of Kinetically Stabilized Main Group Element Compounds using 9-Triptycyl Groups MINOURA, Mao, Faculty of Science, Rikkyo University

Host in JURC TOKITOH, Norihiro

Studies on Structure-property Relationship of Pyrylium Salts NAGAHORA, Noriyoshi, Department of Chemistry, Faculty of

Science, Fukuoka University

Host in JURC SASAMORI, Takahiro

STEM-EELS Measurement of a Plasmonic Full Bandgap and Cavity Modes

HATA, Satoshi, Interdisciplinary Graduate School of Engineering Sciences, Kyushu University

Host in JURC KURATA, Hiroki

Nano-electron Spectroscopic Study on Hydrogen and Helium Behavior in Divertor Materials for Nuclear Fusion Devices

MIYAMOTO, Mitsutaka, Interdisciplinary Faculty of Science and Engineering, Shimane University

Host in JURC KURATA, Hiroki

Fabrication of Metal Nanowires and Nanomeshes and Their Catalytic Activity

KAWAI, Takeshi, Faculty of Engineering, Tokyo University of Science

Host in JURC KURATA, Hiroki

Study on Novel Supramolecular Assemblies Using High-resolution Mass Spectrometry

YAMANAKA, Masamichi, Faculty of Science, Department of Chemistry, Shizuoka University

Host in JURC TAKAYA, Hikaru

Novel Synthesis of Metal Clusters Protected with Organic Ligands and Determination of Their Compositions

KOYASU, Kiichirou, Department of Chemistry, Graduate School of Science, The University of Tokyo

Host in JURC TAKAYA, Hikaru

Development of Direct Visualization Methods of Steroid Hormones by Imaging Mass Spectrometry

HATANO, Osamu, Department of Anatomy and Cell Biology, Nara Medical University

Host in JURC ISOZAKI, Katsuhiro

SUBJECTS ENCOURAGING JOINT PROGRAM

Research Collaborations Aiming for a New Paradigm in Oxide Electronics

HITOSUGI, Taro, School of Materials and Chemical Technology, Tokyo Institute of Technology

Host in JURC KAN, Daisuke

Modification of Cell-penetrating Peptides for the Effective Delivery of Molecules into the Cells

BÁNÓCZI, Zoltán, Department of Organic Chemistry, Eötvös Loránd University of Sciences, Hungary

I

Host in JURC FUTAKI, Shiroh

The Twelfth International Workshop for East Asian Young Rheologists INOUE, Tadashi, Graduate School of Science, Osaka University Ι Host in JURC WATANABE, Hiroshi

Clarification of the Relationship between Layered Oxide Structure and Its Magnetic Property

SUZUKI, Furitsu, School of Chemical Sciences, The University of Auckland

Ι Host in JURC KAJI, Hironori

JURC Publications (Selected Examples)

(until 31 May 2016)

Purely Organic Electroluminescent Material Realizing 100% Conversion from Electricity to Light

Kaji, H.; Suzuki, H.; Fukushima, T.; Shizu, K.; Suzuki, K.; Kubo, S.; Komino, T.; Oiwa, H.; Suzuki, F.; Wakamiya, A.; Murata, Y.; Adachi, C., *Nat. Commun.*, **6**, [8476-1]-[8476-8] (2015).

Abstract

Efficient organic light-emitting diodes have been developed using emitters containing rare metals, such as platinum and iridium complexes. However, there is an urgent need to develop emitters composed of more abundant materials. Here we show a thermally activated delayed fluorescence material for organic light-emitting diodes, which realizes both approximately 100% photoluminescence quantum yield and approximately 100% up-conversion of the triplet to singlet excited state. The material contains electron-donating diphenylaminocarbazole and electron-accepting triphenyltriazine moieties. The typical trade-off between effective emission and triplet-to-singlet up-conversion is overcome by fine-tuning the highest occupied molecular orbital and lowest unoccupied molecular orbital distributions. The nearly zero singlet-triplet energy gap, smaller than the thermal energy at room temperature, results in an organic light-emitting diode with external quantum efficiency of 29.6%. An external quantum efficiency of 41.5% is obtained when using an out-coupling sheet. The external quantum efficiency is 30.7% even at a high luminance of 3,000 cd m⁻².

Photoluminescence of Monovalent Indium Centres in Phosphate Glass

Masai, H.; Yamada, Y.; Okumura, S.; Yanagida, T.; Fujimoto, Y.; Kanemitsu, Y.; Ina, T., *Sci. Rep.*, **5**, 13646 (2015).

Abstract

Valence control of polyvalent cations is important for functionalization of various kinds of materials. Indium oxides have been used in various applications, such as indium tin oxide in transparent electrical conduction films. However, although metastable In+ (5 s2 configuration) species exhibit photoluminescence (PL), they have attracted little attention. Valence control of In+ cations in these materials will be important for further functionalization. Here, we describe In+ species using PL and X-ray absorption fine structure (XAFS) analysis. Three absorption bands in the UV region are attributed to the In+ centre: two weak forbidden bands (${}^{1}S_{0} \rightarrow {}^{3}P_{1}, {}^{1}S_{0} \rightarrow {}^{3}P_{2}$) and a strong allowed band (${}^{1}S_{0} \rightarrow {}^{1}P_{1}$). The strongest PL excitation band cannot be attributed to the conventional allowed transition to the singlet excited state. Emission decay of the order of microseconds suggests that radiative relaxation occurs from the triplet excitation state. The XAFS analysis suggests that these In+ species have shorter In-O distances with lower coordination numbers than in In₂O₃. These results clearly demonstrate that In+ exists in a metastable amorphous network, which is the origin of the observed luminescent properties.

Catalytic Discrimination between Formyl Groups in Regio- and Stereoselective Intramolecular Cross-aldol Reactions

Baba, T.; Yamamoto, J.; Hayashi, K.; Sato, M.; Yamanaka, M.; Kawabata, T.; Furuta, T., *Chem. Sci.*, 7, 3791-3797 (2016).

Catalytic discrimination between inequivalent formyl groups was achieved using an aniline-type acid-base catalyst for the regio-, diastereo-, and enantioselective intramolecular cross-aldol reactions of enolizable dials. Although L-proline gave a mixture of the regio- and

stereoisomeric products in the presence of an *N*-containing 1,6-dial, the aniline-type catalyst afforded *anti*-3,4-disubstituted pyrrolidine in high regio-, and stereoselectivity beyond the background reaction, which led to the regioisomeric 2,3-disubstituted products. The mild reactivity of the aniline-type amine facilitated catalytic discrimination between the inequivalent formyl groups. Kinetic isotope effect studies and reductive amination experiments suggested that the regioselectivity was controlled under the enamine-forming steps.

Rhombic Coulomb Diamonds in a Single-electron Transistor Based on an Au Nanoparticle Chemically Anchored at Both Ends Azuma, Y.; Onuma, Y.; Sakamoto, M.; Teranishi, T.; Majima, Y., *Nanoscale*, **8**, 4720-4726 (2016).

Abstract

Rhombic Coulomb diamonds are clearly observed in a chemically anchored Au nanoparticle single-electron transistor. The stability diagrams show stable Coulomb blockade phenomena and agree with the theoretical curve calculated using the orthodox model. The resistances and capacitances of the double-barrier tunneling junctions between the source electrode and the Au core (R_1 and C_1 , respectively), and those between the Au core and the drain electrode (R_2 and C_2 , respectively), are evaluated as 4.5 M Ω , 1.4 aF, 4.8 M Ω , and 1.3 aF, respectively. This is determined by fitting the theoretical curve against the experimental Coulomb staircases. Two-methylene-group short octanedithiols (C8S2) in a C8S2/hexanethiol (C6S) mixed self-assembled monolayer is concluded to chemically anchor the core of the Au nanoparticle at both ends between the electroless-Au-plated nanogap electrodes even when the Au nanoparticle is protected by decanethiol (C10S). This is because the R_1 value is identical to that of R_2 and corresponds to the tunneling resistances of the octanedithiol chemically bonded with the Au core and the Au electrodes. The dependence of the Coulomb diamond shapes on the tunneling resistance ratio (R_1/R_2) is also discussed, especially in the case of the rhombic Coulomb diamonds. Rhombic Coulomb diamonds result from chemical anchoring of the core of the Au nanoparticle at both ends between the electroless-Au-plated nanogap electrodes.

Soliton-like Magnetic Domain Wall Motion Induced by the Interfacial Dzyaloshinskii-Moriya Interaction

Yoshimura, Y.; Kim, K.-J.; Taniguchi, T.; Tono, T.; Ueda, K.; Hiramatsu, R.; Moriyama, T.; Yamada, K.; Nakatani, Y.; Ono, T., *Nat. Phys.*, **12**, 157-161 (2016).

Abstract

Topological defects such as magnetic solitons, vortices and skyrmions have started to play an important role in modern magnetism because of their extraordinary stability, which can be exploited in the production of memory devices. Recently, a type of antisymmetric exchange interaction, namely the Dzyaloshinskii–Moriya interaction (DMI; refs), has been uncovered and found to influence the formation of topological defects. Exploring how the DMI affects the dynamics of topological defects is therefore an important task. Here we investigate the dynamics of the magnetic domain wall (DW) under a DMI by developing a real time DW detection scheme. For a weak DMI, the DW velocity increases with the external field and reaches a peak velocity at a threshold field, beyond which it abruptly decreases. For a strong DMI, on the other hand, the velocity reduction is completely suppressed and the peak velocity is maintained constant even far above the threshold field. Such a distinct trend of the velocity can

be explained in terms of a magnetic soliton, the topology of which is protected during its motion. Our results therefore shed light on the physics of dynamic topological defects, which paves the way for future work in topology-based memory applications.

VISITING PROFESSORS' ACTIVITIES IN ICR



Vis Prof MATSUO, Yutaka (D Sc)

Laboratory of Polymer Controlled Synthesis

Project Professor, School of Engineering, The University of Tokyo (7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656)

Lecture at *ICR*

Materials Science of Organic Semiconductor Functionalization of Fullerene/Carbon Nanotube and Development of Organic Thin Film Solar Cell



Vis Prof MURAYAMA, Masafumi (D Sc)

Laboratory of Hydrospheric Environment Analytical Chemistry

Professor, Center for Advanced Marine Core Research, Kochi University (Monobeotsu, Nankoku, Kochi 783-8502)

Lecture at ICR

What occurred on the Shallow and Deep Seafloor by the Great East Japan Earthquake



Vis Prof SUGIOKA, Koji (D Eng)

Laboratory of Laser Matter Interaction Science

Unit Leader, RIKEN-SIOM Joint Research Unit, RIKEN (2-1 Hirosawa, Wako, Saitama 351-0198)

Lecture at ICR

High-performance Materials Processing by Using Manipulated Laser Beams



Vis Prof AKIYAMA, Yutaka (D Eng)

Laboratory of Mathematical Bioinformatics

Professor, Department of Computer Science School of Computing, Tokyo Institute of Technology (2-12-1 Ookayama, Meguro-ku, Tokyo 152-8552)

Lecture at ICR

Massively Parallel Bioinformatics Computing on K-computer, TSUBAME, and Azure: Metagenome Analysis and Exhaustive Proteinprotein Interaction Prediction



Vis Prof DE LA HIGUERA, Colin Manuel (Ph D)

Laboratory of Mathematical Bioinformatics

Professor, UFR de Sciences et Techniques, University of Nantes (2 rue de la Houssinière, BP 92208, 44322 Nantes Cedex 03, France)

Lecture at ICR

Machine Learning is about Intelligent Forgetting



Vis Assoc Prof FURUKAWA, Ko (D Sc)

Laboratory of Structural Organic Chemistry

Associate Professor, Centre for Instrumental Analysis, Niigata University (Nishi-ku, Niigata 950-2181)

Lecture at ICR

Spin Dynamics Study for Functional Materials by Advanced ESR Spectroscopy



Vis Assoc Prof KAWANO, Ryuji (D Eng)

Laboratory of Biofunctional Design-Chemistry

Associate Professor, Tokyo University of Agriculture and Technology (2-24-16 Naka-cho, Koganei-shi, Tokyo 184-8588)

Lecture at ICR

Construction and Application of Artificial Cell Membranes Using Micro-Sized Processing Techniques



Vis Assoc Prof TAKAHASHI, Yoshiaki (D Eng)

Laboratory of Molecular Rheology

Associate Professor, Institute for Materials Chemistry and Engineering, Kyushu University (6-1 Kasuga Koen, Kasuga, Fukuoka

816-8581)



Dissolution Condition of Cellulose into Ionic Liquids and Characterization of the Solutions by Rheological Measurement



Vis Assoc Prof OGAWA. Yoshihiro (D Sc)

Laboratory of Nanophotonics

Associate Professor, Graduate School of Education, Joetsu University of Education (1, Yamayashiki-machi, Joetsu-shi, Niigata 943-8512)

Lecture at ICR

Coherent Spectroscopy and its Application to the Coherent Control of Quantum States in Semiconductors



Vis Assoc Prof ZHU, Shanfeng (D Eng)

Laboratory of Bio-knowledge Engineering

Associate Professor, Fudan University (No.825, Zhangheng Road, Shanghai, China)

Lecture at ICR

DeepMeSH and MeSHLabeler: Recent Progress in Large-scale MeSH Indexing DrugE-Rank: Improving Drug-target Interaction Prediction of New Candidate Drugs or Targets by Ensemble Learning to Rank



Prof Em KANEHISA, Minoru (D Sc)

Appointed as Res (pt) at ICR, 1 April 2012-31 March 2017

Personal

Awards



IZUHATA, Yoshiyuki Progress Award in Silicon Chemistry, Japan

"New Progress in the Chemistry of Aromatic Compounds Containing a Group 14 element"

The Society of Silicon Chemistry, Japan

8 October 2016



SUZUKI, Yuko ICR Award for Graduate Students

Institute for Chemical Research, Kyoto

"Isolation and Ambident Reactivity of a Chlorogermylenoid"

University

2 December 2016



SUGAHARA, Tomohiro
Best Presentation Award

The 4th Presentation of the Master's Thesis
"Synthesis and Reactivity of Stable Multiple-bond Compounds between Germanium

Atoms'

Grant-in-Aid for Scientific Research on Innovative Areas, New Polymeric Materials Based on Element-Blocks

4 March 2016



HANG, Rui
ICR Award for Graduate Students

"Synthesis of a Distinct Water Dimer inside Fullerene C_{70} "

Institute for Chemical Research, Kyoto

University

2 December 2016



UEDA, Yoshihiro
The Pharmaceutical Society of Japan, Kinki
Branch

Award for Young Scientists

66th Annual Meeting of the PSJ Kinki-Branch

"Final-Stage Site-Selective Acylation for the Total Synthesis of Natural Glycosides"

The Pharmaceutical Society of Japan, Kinki-Branch

8 January 2016



LU, Wenjie Best Presentation Award

The 136th Annual Meeting of the Pharmaceutical Society of Japan

"Preparation of Functionalized Rh Carboxylate Catalyst and Its Catalytic Activity"

The Pharmaceutical Society of Japan

31 March 2016



KASAMATSU, Koji

Best Presentation Award

The 136th Annual Meeting of the Pharmaceutical Society of Japan

"Long-Lived C-N Axially Chiral Enolate Generated from Amino Acid Derivatives"

The Pharmaceutical Society of Japan

31 March 2016



SUJII, Yoshinobu
The Cellulose Society of Japan Award

"Fabrication of Functional Cellulosic Materials through Interfacial Chemical Approach"

The Cellulose Society of Japan

14 July 2016



S AKAKIBARA, Keita

The Young Researcher Award of The Society of Fiber Science and Technology, Japan

"Development and Application of Polymer Monolith Materials with Well-Defined Morphol-

ogy and High Performance"

The Society of Fiber Science and Technology, Japan

8 June 2016



AKAMURA, Yasuyuki ICR Award for Young Scientists

"Termination Mechanism in the Radical Polymerization of Methyl Methacrylate and Styrene Determined by the Reaction of Structurally

Well-defined Polymer End Radicals"

Institute for Chemical Research, Kyoto University

2 December 2016



ASAI, Hirokazu International Exchange Encouragement Award: The Motoharu Kurata Award The Ceramic Society of Japan

3 June 2016



l ORIMOTO, Aya **Presentation Award for Young Scientists** Workshop on Next-generation Optical Materials and their Applications

Borate Glasses"

20 December 2016



Y OSHIMURA, Yoko President Prize of Kyoto University

"Study of Spin Dynamics in Ferromagnetic Nanowire"

Kyoto University

14 March 2016



URAYAMA, Tomo

The 34th European Peptide Symposium and the 8th International Peptide Symposium

"Membrane-remodeling Amphipathic Peptides Enhance Direct Penetration of Octaarginine"

The European Peptide Society

22 June 2016



HKANDA, Junko The 21st Incentive Award

"Rational Design of Synthetic Molecules that Control Protein-protein Interactions"

The Society of Japanese Women Scientists

6 April 2016



ATSUDA, Yousuke The Best Presentation Award

The 10th Symposium on Biorelevant Chemistry "Exploration of RNA G-quadruplex Structures by an RNA G-quadruplex Ligand"

The Chemical Society of Japan

8 September 2016



The Best Presentation Award for Young **Scientists**

The 22nd Regular Meeting of Japan OLED Forum

"Solution-Processable Small Molecule TADF Emitters for Highly Efficient (Host-Free) Organic Light-Emitting Diodes"

Japan OLED Forum

17 November 2016



OHRIN, Yoshiki

2016 The Oceanographic Society of Japan

The 2016 Spring Meeting of the Oceanographic Society of Japan

"Development of Precise Analytical Methods for Trace Elements and Its Application to Marine Chemistry"

The Oceanographic Society of Japan

1 April 2016



HIMOAKA, Takafumi

2016 Young Scientist Presentation Award of The Spectroscopical Society of Japan

2016 Annual Meeting of the Spectroscopical Society of Japan

"Structural and Quantitative Analysis of Three Water Species Involved in Nafion Membrane by Using Infrared and NMR Spectroscopy"

The Spectroscopical Society of Japan

2 June 2016



HIOYA, Nobutaka **CSJ Student Presentation Award 2016**

The 96th CSJ Annual Meeting

"Experimental Optimization of IR pMAIRS by Using an Optically Isotropic Thin Film Exhibit-

ing Berreman's Effect"

The Chemical Society of Japan

13 April 2016

Young Scientist Presentation Award

The 63rd JSAP Spring Meeting, 2016

"Structure Analysis of an Organic Semiconductor Thin Film Using IR pMAIRS"

The Japan Society of Applied Physics

13 September 2016



Award for the Presentation of an Excellent Paper

The 36th Annual Meeting of the Laser Society of Japan

"Terahertz Wave Generation in the Interaction of Two-pulse Laser with Clusters"

The Laser Society of Japan

31 May 2016

Award for the Presentation of an Excellent Paper

"THz Waves Generation in the Interaction between Two-pulse Lasers and Cluster Plasmas"

The Institute of Electric Engineers of Japan

5 September 2016



ERAMOTO, Kensuke
Award for the Presentation of an Excellent
Paper

"Surface Waves Generated on a Metal Wire by Intense Laser Plasma"

The Institute of Electric Engineers of Japan

5 September 2016



Furukawa, Yuki

Award for the Presentation of an Excellent Paper

The 36th Annual Meeting of the Laser Society of Japan

"Formation of LIPSS on Titanium Irradiated by Femtosecond Double Pulse Laser~Features of Structures Produced at Delay Time from 100fs to Sub Nanosecond~"

The Laser Society of Japan

31 May 2016



HARUTA, Mitsutaka
ICR Award for Young Scientists

"Determination of Elemental Ratio in an Atomic Column by Electron Energy Loss Spectroscopy" Institute for Chemical Research, Kyoto

University

2 December 2016



YOSHIDA, Ryota Student Presentation Award The 61st Lignin Symposium

"Ruthenium Complex-Bound Amino Acid-Catalyzed Oxidative Degradation of Lignin and

Model Compounds"

Research Institute for Sustainable Humanosphere, Kyoto University

28 October 2016



SHIMAKAWA, Yuichi; SAITO, Takashi Daiwa Adrian Prizes 2016



"Materials Discovery using Extreme Conditions" Daiwa ANGLO-JAPANESE FOUNDATON 15 November 2016



HOSAKA, Yoshiteru
The Best Student Presentation Award

2016 E-MRS Fall Meeting and Exhibit
"A New Layered Double Perovskite Ca₂FeMnO₆
Showing Charge Disproportionation of Unusual-

ly High Valence Fe4+"

European Materials Research Society

21 September 2016



TAHARA, Hirokazu

NF Foundation R&D Encouragement Award

"Highly Sensitive Photocurrent Spectroscopy Using Femtosecond Phase-locked Pulses and Photocurrent Generation Dynamics in

Semiconductors" NF Foundation

25 November 2016



YAMADA, Takumi ICR Award for Graduate Students

"Dynamic Optical Properties of CH₃NH₃PbI₃ Single Crystals As Revealed by One- and Two-Photon Excited Photoluminescence

Measurements'

Institute for Chemical Research, Kyoto University

2 December 2016



AKUTSU, Tatsuya

Fellov

2016 General Meeting of Information Processing Society of Japan

"Studies on Mathematical Models and Algorithms for Bioinformatics"

Information Processing Society of Japan

3 June 2016

AKAJIMA, Natsu

IPSJ Yamashita SIG Research Award

"Genetic Network Completion for Static Gene Expression Data"

The Information Processing Society of Japan 10 March 2016

Paper Awards



SUGAHARA, Tomohiro; GUO, Jing-Dong; SASAMORI, Takahiro; TOKITOH, Norihiro *et al*.



"Reaction of a Stable Digermyne with Acetylenes: Synthesis of a 1,2-Digermabenzene and a 1,4-Digermabarrelene"



Sugahara, T.; Guo, J.-D.; Sasamori, T.; Karatsu, Y.; Furukawa, Y.; Espinosa Ferao, A.; Nagase, S.; Tokitoh, N.

The Chemical Society of Japan

BCSJ Award Article

15 November 2016





WAKAMIYA, Atsushi *et al*.

The Innovative PV Award

"Perovskite Solar Cells: Elucidation of Structures and Properties of Intermediates in Solution Process and Development of Fabrication Method for Highly Efficient Cells"

Wakamiya, A.; Shimazaki, A.; Nakaike, Y.; Rafieh, A.; Jung, M.; Maruyama, N.; Aharen, T.; Murata, Y.

JSPS 175 Committee on Future Direction of Photovoltaics

20 May 2016



MORISHITA, Hiroki; MIZUOCHI, Norikazu

38th Paper Award of the Japan Society of Applied Physics



"Perfect Selective Alignment of Nitrogen-vacancy Centers in Diamond"

Fukui, T.; Doi, Y.; Miyazaki, T.; Miyamoto, Y.; Kato, H.; Matsumoto, T.; Makino, T.; Yamasaki, S.; Tokuda, N.; Hatano, M.; Sakagawa, Y.; Morishita, H.; Tashima, T.; Miwa, S.; Suzuki, Y.;

Mizuochi, N.
The Japan Society of Applied Physics

13 September 2016



KAJI, Hironori; WADA, Yoshimasa; SHIZU, Katsuyuki; KUBO, Shosei; SUZUKI, Katsuaki; FUKUSHIMA, Tatsuya; ADACHI, Chihaya¹



The 23rd International Display Workshops in Conjunction with Asia Display 2016

"Design of Emitters and Charge Transporters for Highly Efficient Organic LEDs"

International Display Workshops

Best Paper Award

22 December 2016



¹ Department of Applied Chemistry, Faculty of Engineering, Kyushu University







HASHIDA, Masaki; MIYASAKA, Yasuhiro; SHIMIZU, Masahiro¹; SAKAGAMI, Hitoshi²; **INOUE**, Shunsuke; SAKABE, Shuji LSJ Award for Distinguished Achievements in



"Periodic Surface Structures Self-organized on Solid Materials Ablated by Femtosecond Laser Pulses"

The Laser Society of Japan.

Research

31 May 2016

- ¹ Department of Material Chemistry, Graduate School of Engineering, Kyoto University
- ² National Institute for Fusion Science





AMADA, Makoto The Best Paper Award

The 9th ACM international Conference on Web Search and Data Mining (WSDM)

"Beyond Ranking: Optimizing Whole-Page

Presentation"

ACM (Association for Computing Machinery)

22 February 2016

Poster Awards



FUJIMORI, Shiori

The 27th International Conference on Organometallic Chemistry (ICOMC 2016)

"Reaction of Germabenzenylpotassium with [Cp*RuCl]₄: Unexpected Formation of Ruthenium-substituted Germabenzene"

21 July 2016



Sugahara, Tomohiro
Chemical Communications Poster Prizes

The 20th Symposium of the Society of Silicon Chemistry Japan

"Synthesis and Structure of Cyclic Compounds Containing Germanium Atoms Utilizing a Digermacyclobutene Derivative"

Royal Society of Chemistry/The Society of Silicon Chemistry Japan

8 October 2016



TUTAGOISHI, Tsukasa
27th Symposium on Physical Organic
Chemistry

"Synthesis and Properties of an Open-Cage C₆₀ Derivative Encapsulating Molecular Oxygen"

The Society of Physical Organic Chemistry

3 September 2016



OZAKI, Masashi Best Poster Award, 2nd Prize

AEARU 6th Energy and Environmental Workshop

"Development of $CH_3NH_3PbI_3\cdot DMF$ as a Perovskite Precursor for Highly Efficient Perovskite Solar Cells"

The Association of East Asian Research Universities

27 August 2016

Best Poster Award, First Place

The 1st Frontier Solar Cells Seminar

"Fabrication of Highly Efficient Perovskite Solar Cells Using Highly Purified Precursor Material"

The 1st Frontier Solar Cells Seminar Executive Committee

18 November 2016



SUKAO, Masahiro
Chemistry Letters Young Awards

The 10th Symposium of Organic π -Electron Systems

"Synthesis and Properties of Aluminum Complexes with π -Extended Quinolinol Ligands"

The Japanese Association for Organic π -Electron Systems

17 December 2016



Kasamatsu, Koji

Best Poster Award

42nd Symposium on Progress in Organic Reactions and Synthesis -Application in the Life Science-

"Asymmetric α-Fluorination of Amino Acid Derivatives via Racemization-Resistant C-N Axially Chiral Enolates"

Organizing Committee of 42nd Symposium on Progress in Organic Reactions and Synthesis - Application in the Life Science-

7 November 2016



OTSUKI, Haruka

Best Poster Award

Molecular Chirality Asia 2016

"Mechanistic Investigation of Stereochemical Diversity in Asymmetric Cyclization via

Memory of Chirality "

Molecular Chirality Research Organization (MCRO)

23 April 2016

Best Poster Award

110th Symposium on Organic Synthesis

"Investigation of Stereochemical Diversity in Asymmetric Intramolecular Alkylation via Memory of Chirality"

The Society of Synthetic Organic Chemistry, Japan

11 November 2016



EGUCHI, Daichi

Poster Award

"Investigation of the Photochemical Behavior Between Semiconductor Quantum Dots and Porphyrin Derivatives"

The 14th Annual Meeting of The Society of Nano Science and Technology

The Society of Nano Science and Technology

15 June 2016



Annual Meeting Poster Award 2016

65th SPSJ Annual Meeting

"Dispersibility of Polymer-brush-afforded Particles in Liquid Crystals"

The Society of Polymer Science, Japan

27 May 2016



AWANO, Kenichi

The 14th Chinese International Peptide Symposium & The 5th Asia-pacific International Peptide Symposium

"Facilitated Cell Penetration of Dipicolylamine-modified R8 by Metal Complex Formation"

CPS2016 Committee

7 July 2016



UJIMORI, Tomoya

The Society of Polymer Science, Japan Symposium on Macromolecules Poster Award

65th Symposium on Macromolecules

'Modification of Capillary-inner Surface with Thick and Concentrated Polymer Brushes for Control of Microfluidic Flow"

The Society of Polymer Science, Japan

16 September 2016



KISHIBA, Misao

The 14th Chinese International Peptide Symposium & The 5th Asia-pacific International Peptide Symposium

'Modification of Natural Hemolytic Peptide Achieving Intracellular Delivery of Antibodies"

CPS2016 Committee

7 July 2016



EKI, Toyomitsu The Best Poster Award

The 10th International Symposium on Integrated Synthesis (ISONIS-10)

"Development of an Efficient Method for Tellurium Recovery and Recycling from Polymer-end by Using Fluorous Tellurol and Fluorous Extraction"

Organizing Committee of The 10th International Symposium on Integrated Synthesis (ISONIS-10)

19 November 2016



ASUDA, Toshihiro **Award for Excellence at Poster Presentation**

The 53rd Japanese Peptide Symposium

"Novel Peptide Tool to Induce Cell Movement by Affecting Membrane Tension"

The Japanese Peptide Society

27 October 2016



ORIMOTO, Aya The Best Poster Presentation Award

The Ceramic Society of Japan Fall Meeting 2016 "Emission Properties of Ce-doped Barium Borate Glasses Prepared from Different Starting

Materials"

Random Materials Session, The Ceramic Society of Japan

9 September 2016

The Best Poster Presentation Award

JSAP 77th JSAP Autumn Meeting 2016

"Study of Emission Properties of Cerium-doped Barium Borate Glasses Depending on the Starting Materials"

JSAP Ionization Radiation Division

15 September 2016



L AGUCHI, Hiro-omi

5th JACI/GSC Symposium

"Hydrosilylation of CO2 Catalyzed by Noninnocent PNP-Pincer Type Phosphaalkene Complex

with Iridium(I)"

Japan Association for Chemical Innovation (JACI)

3 June 2016

Poster Award

20th International Symposium on Homogeneous Catalysis (ISHCXX)

"Synthesis of Ir(I) and Rh(I) Complexes Bearing a Noninnocent PNP-Pincer Type Phosphaalkene Ligand (Eind-PPEP) and Their Application for Catalytic Hydrosilylation of CO2"

ISHCXX Organizers

15 July 2016



HIROSHIGE, Nao

Region5 Poster Awards

The Physical Society of Japan 2016 Autumn Meeting

"Cascade-emission Spectroscopy on Single Giant CdSe/CdS Nanocrystals: Enhancement of Biexciton Emis-sion through Coulomb Interaction"

The Physical Society of Japan

15 September 2016

Publications International RESEARCH COLLABORATIONS SELECTED GRANTS

Selected grants
Theses

PUBLICATIONS

DIVISION OF SYNTHETIC CHEMISTRY — Organoelement Chemistry —

- Agou, T.; Ikeda, S.; Sasamori, T.; Tokitoh, N., Synthesis and Structure of Lewis-Base-Free Phosphinoalumane Derivatives, *Eur. J. Inorg. Chem.*, 623-627 (2016).
- Sasamori, T.; Franco, J. M. V.; Guo, J.-D.; Sugamata, K.; Nagase, S.; Streubel, R.; Tokitoh, N., Selenium-Substituted Phosphaalkenes Obtained through 1,2-Elimination of Chlorosilanes from Selenenylchlorophosphines, *Eur. J. Inorg. Chem.*, 678-684 (2016).
- Klein, M.; Schnakenburg, G.; Ferao, A. E.; Tokitoh, N.; Streubel, R., Reactions of Li/Cl Phosphinidenoid Complexes with 1,3,4,5-Tetramethylimidazol-2-ylidene: A New Route to *N*-Heterocyclic Carbene Adducts of Terminal Phosphinidene Complexes and an Unprecedented Transformation of an Oxaphosphirane Complex, *Eur. J. Inorg. Chem.*, 685-690 (2016).
- Guo, J.-D.; Sasamori, T.; Yamamoto, Y.; Matsubara, H.; Nagase, S.; Yorimitsu, H., Computational Picture of Silyl Transfer from Silylsilatranes to Arylpalladium Chloride, *Bull. Chem. Soc. Jpn.*, **89**, 192-194 (2016).
- Tokitoh, N.; Agou, T.; Wasano, T.; Sasamori, T., Synthesis and Properties of Stable Alumoles, *Phosphorus, Sulfur and Silicon and the Related Elements*, **191**, 584-587 (2016).
- Agou, T.; Nagata, K.; Sasamori, T.; Tokitoh, N., Reactivities of a Barrelene-type Dialumane as an Equivalent of an Al=Al Doubly-bonded Species, *Phosphorus, Sulfur and Silicon and the Related Elements*, **191**, 588-590 (2016).
- Suzuki, Y.; Sasamori, T.; Guo, J.-D.; Nagase, S.; Tokitoh, N., Isolation and Ambident Reactivity of a Chlorogermylenoid, *Chem. Eur. J.*, **22**, 13784-13788 (2016).
- Yamaguchi, K.; Murai, T.; Guo, J.-D.; Sasamori, T.; Tokitoh, N., Acid-Responsive Absorption and Emission of 5-N-Arylaminothiazoles: Emission of White Light from a Single Fluorescent Dye and a Lewis Acid, *ChemistryOpen*, **5**, 434-438 (2016).
- Nagata, K.; Murosaki, T.; Agou, T.; Sasamori, T.; Matsuo, T.; Tokitoh, N., Activation of Dihydrogen by Masked Doubly Bonded Aluminum Species, *Angew. Chem. Int. Ed.*, **55**, 12877-12880 (2016).
- Agou, T.; Yanagisawa, T.; Sasamori, T.; Tokitoh, N., Synthesis and Structure of an Iron-Bromoalumanyl Complex with a Tri-Coordinated Aluminum Center, *Bull. Chem. Soc. Jpn.*, **89**, 1184-1186 (2016).
- Mizuhata, Y.; Yanagisawa, T.; Sasamori, T.; Tokitoh, N., [K(18-crown-6)][FeCp*(CO)₂] data, *IUCrData*, **1**, x160881 (2016).
- Sugahara, T.; Guo, J.-D.; Sasamori, T.; Karatsu, Y.; Furukawa, Y.; Ferao, A. E.; Nagase, S.; Tokitoh, N., Reaction of a Stable Digermyne with Acetylenes: Synthesis of a 1,2-Digermabenzene and a 1,4-Digermabarrelene, *Bull. Chem. Soc. Jpn.*, **89**, 1375-1384 (2016).

- Structural Organic Chemistry -

- Makuta, S.; Liu, M. N.; Endo, M.; Nishimura, H.; Wakamiya, A.; Tachibana, Y., Photo-Excitation Intensity Dependent Electron and Hole Injections from Lead Iodide Perovskite to Nanocrystalline TiO₂ and Spiro-OMeTAD, *Chem. Commun.*, **52**, 673-676 (2016).
- Yamada, T.; Yamada, Y.; Nishimura, H.; Nakaike, Y.; Wakamiya, A.; Murata, Y.; Kanemitsu, Y., Fast Free-Carrier Diffusion in CH₃NH₃PbBr₃ Single Crystals Revealed by Time-Resolved One-and Two-Photon Excitation Photoluminescence Spectroscopy, *Adv. Electron. Mater.*, **2**, [1500290-1]-[1500290-5] (2016).
- Lee, J.; Jung, H.; Shin, H.; Kim, J.; Yokoyama, D.; Nishimura, H.; Wakamiya, A.; Park, J., Excimer Emission Based on the Control of Molecular Structure and Intermolecular Interactions, *J. Mater. Chem. C*, **4**, 2784-2792 (2016).
- Kamimoto, N.; Nakamura, N.; Tsutsumi, A.; Mandai, H.; Mitsudo, K.; Wakamiya, A.; Murata, Y.; Hasegawa, J. Y.; Suga, S., Facile Synthesis of 1,4-Bis(diaryl)-1,3-butadiynes Bearing Two Amino Moieties by Electrochemical Reaction-Site Switching, and Their Solvatochromic Fluorescence, *Asian J. Org. Chem.*, **5**, 373-379 (2016).
- Tahara, H.; Endo, M.; Wakamiya, A.; Kanemitsu, Y., Experimental Evidence of Localized Shallow States in Orthorhombic Phase of CH₃NH₃PbI₃ Perovskite Thin Films Revealed by Photocurrent Beat Spectroscopy, *J. Phys. Chem. C*, **120**, 5347-5352 (2016).
- Zhang, R.; Murata, M.; Aharen, T.; Wakamiya, A.; Shimoaka, T.; Hasegawa, T.; Murata, Y., Synthesis of a Distinct Water Dimer inside Fullerene C₇₀, *Nat. Chem.*, **8**, 435-441 (2016).
- Nishimura, H.; Fukushima, T.; Wakamiya, A.; Murata, Y.; Kaji, H., The Influence of Quasiplanar Structures of Partially Oxygen-Bridged Triphenylamine Dimers on the Properties of Their Bulk Films, *Bull. Chem. Soc. Jpn.*, **89**, 726-732 (2016).
- Hashikawa, Y.; Murata, M.; Wakamiya, A.; Murata, Y., Synthesis and Properties of Endohedral Aza[60] fullerenes: $H_2O@C_{59}N$ and $H_2@C_{59}N$ as Their Dimers and Monomers, *J. Am. Chem. Soc.*, **138**, 4096-4104 (2016).
- Satou, M.; Nakamura, T.; Aramaki, Y.; Okazaki, S.; Murata, M.; Wakamiya, A.; Murata, Y., Near-infrared Emissive Donor-Acceptor-type Molecules Containing Thiazole-fused Benzothiadiazole as an Electron-acceptor Moiety, *Chem. Lett.*, **45**, 892-894 (2016).
- Murata, M.; Kaji, S.; Nishimura, H.; Wakamiya, A.; Murata, Y., Efficient Synthesis of One- and Two-Dimensional Multimetallic Gold-Bis(dithiolene) Complexes, *Eur. J. Inorg. Chem.*, **2016**, 3228-3232 (2016).
- Hashikawa, Y.; Murata, M.; Wakamiya, A.; Murata, Y., Water Entrapped inside Fullerene Cages: A Potential Probe for Evaluation of Bond Polarization, *Angew. Chem., Int. Ed.*, **55**, 13109-13113 (2016).

- Wang, F. J.; Endo, M.; Mouri, S.; Miyauchi, Y.; Ohno, Y.; Wakamiya, A.; Murata, Y.; Matsuda, K., Highly Stable Perovskite Solar Cells with an All-Carbon Hole Transport Layer, *Nanoscale*, **8**, 11882-11888 (2016).
- Yamada, Y.; Yamada, T.; Shimazaki, A.; Wakamiya, A.; Kanemitsu, Y., Interfacial Charge-Carrier Trapping in CH₃NH₃PbI₃-Based Heterolayered Structures Revealed by Time-Resolved Photoluminescence Spectroscopy, *J. Phys. Chem. Lett.*, **7**, 1972-1977 (2016).
- Le, P. Q.; Yamada, Y.; Nagai, M.; Maruyama, N.; Wakamiya, A.; Kanemitsu, Y., Free Carriers versus Excitons in CH₃NH₃PbI₃ Perovskite Thin Films at Low Temperatures: Charge Transfer from the Orthorhombic Phase to the Tetragonal Phase, *J. Phys. Chem. Lett.*, **7**, 2316-2321 (2016).
- Yamashita, D.; Handa, T.; Ihara, T.; Tahara, H.; Shimazaki, A.; Wakamiya, A.; Kanemitsu, Y., Charge Injection at the Heterointerface in Perovskite CH₃NH₃PbI₃ Solar Cells Studied by Simultaneous Microscopic Photoluminescence and Photocurrent Imaging Spectroscopy, *J. Phys. Chem. Lett.*, **7**, 3186-3191 (2016).
- Ishida, N.; Wakamiya, A.; Saeki, A., Quantifying Hole Transfer Yield from Perovskite to Polymer Layer: Statistical Correlation of Solar Cell Outputs with Kinetic and Energetic Properties, *ACS Photonics*, **3**, 1678-1688 (2016).
- Wakamiya, A.; Nishimura, H.; Murata, Y., Partially Oxygen-Bridged Triphenylamines with a Quasiplanar Structure as a Key Scaffold for Hole-Transporting Materials, *J. Synth. Org. Chem. Jpn.*, **74**, 1128-1135 (2016).
- Bottari, G.; Trukhina, O.; Kahnt, A.; Frunzi, M.; Murata, Y.; Rodriguez-Fortea, A.; Poblet, J. M.; Guldi, D. M.; Torres, T., Regio-, Stereo-, and Atropselective Synthesis of C₆₀ Fullerene Bisadducts by Supramolecular-Directed Functionalization, *Angew. Chem., Int. Ed.*, **55**, 11020-11025 (2016).

- Synthetic Organic Chemistry -

- Ueda, Y., Final-stage Site-selective Acylation for The Total Synthesis of Natural Glycosides, *YAKUGAKU ZASSHI*, **136**, 1631-1639 (2016) (in Japanese).
- Alezra, V.; Kawabata, T., Recent Progress in Memory of Chirality (MOC): An Advanced Chiral Pool, *Synthesis*, 2997-3016 (2016).
- Tomohara, K.; Kasamatsu, K.; Yoshimura, T.; Furuta, T.; Kawabata, T., Asymmetric Synthesis of Multisubstituted Dihydrobenzofurans by Intramolecular Conjugate Addition of Short-Lived C-O Axially Chiral Enolates, *Chem. Pharm. Bull.*, **64**, 899-906 (2016).
- Yamada, T.; Suzuki, K.; Hirose, T.; Furuta, T.; Ueda, Y.; Kawabata, T.; Omura, S.; Sunazuka, T., Organocatalytic Site-Selective Acylation of Avermectin B2a, a Unique Endectocidal Drug, *Chem. Pharm. Bull.*, **64**, 856-864 (2016).
- Mishiro, K.; Takeuchi, H.; Furuta, T.; Kawabata, T., A Concise Access to *C*₂-Symmetric Chiral 4-Pyrrolidinopyridine Catalysts with Dual Functional Side Chains, *Chem. Pharm. Bull.*, **64**, 1073-1077 (2016).

- Imayoshi, A.; Yamanaka, M.; Sato, M.; Yoshida, K.; Furuta, T.; Ueda, Y.; Kawabata, T., Insights into the Molecular Recognition Process in Organocatalytic Chemoselective Monoacylation of 1,5-Pentanediol, *Adv. Synth. Catal.*, **358**, 1337-1344 (2016).
- Baba, T.; Yamamoto, J.; Hayashi, K.; Sato, M.; Yamanaka, M.; Kawabata, T.; Furuta, T., Catalytic Discrimination Among Formyl Groups in Regio- and Stereoselective Intramolecular Cross-Aldol Reactions, *Chem. Sci.*, **7**, 3791-3797 (2016).
- Hyakutake, R.; Gondo, N.; Ueda, Y.; Yoshimura, T.; Furuta, T.; Kawabata, T., Catalyst-Controlled Regiodivergent Vinylogous Aza-Morita-Baylis-Hillman Reactions, *Tetrahedron Lett.*, **57**, 1321-1324 (2016).
- Yanagi, M.; Ninomiya, R.; Ueda, Y.; Furuta, T.; Yamada, T.; Sunazuka, T.; Kawabata, T., Organocatalytic Site-Selective Acylation of 10-Deacetylbaccatin III, *Chem. Pharm. Bull.*, **64**, 907-912 (2016).
- Furuta, T.; Kawabata, T., Chiral Imidazole and Pyridines as Asymmetric Organocatalysis, *RSC Green Chemistry, Sustainable Catalysis, M. North, Ed., The Royal Society of Chemistry*, 351-380 (2016).
- Ueda, Y.; Kawabata, T., Organocatalytic Site-Selective Acylation of Carbohydrates and Polyol Compounds, *Top. Curr. Chem.*, **372**, 203-231 (2016).
- Yoshida, K.; Suzuki, H.; Inoue, H.; Fujino, Y.; Itatsu, Y.; Takao, K., Organocatalytic Regioselective Chlorosilylation of Oxiran Derivatives: Mild and Effective Insertion of the Bulky Silyl Chloride by Using 4-Methoxy Pyridine *N*-oxide, *Adv. Synth. Catal.*, **358**, 1886-1891 (2016).
- Yoshida, K.; Itatsu, Y.; Fujino, Y.; Inoue, H.; Takao, K., Enantioselective Organocatalytic Construction of Spiroindane Derivatives by Intramolecular Friedel-Crafts-type 1,4-Addition, *Angew. Chem. Int. Ed.*, **55**, 6734-5738 (2016).
- Miyazaki, S.; Sasazawa, Y.; Mogi, T.; Suzuki, T.; Yoshida, K.; Dohmae, N.; Takao, K.; Simizu, S., Identification of Secoclavilactonoe B as a Small-molecule Actin Polymerization Inhibitor, *FEBS Letters*, **590**, 1163-1173 (2016).
- Yoshida, K.; Fujino, Y.; Itatsu, Y.; Inoue, H.; Kanoko, Y.; Takao, K., Amine Free Silylation of Alcohols under 4-Methylpyridine *N*-Oxide-Catalyzed Conditions, *Tetrahedron Lett.*, **57**, 627-631 (2016).
- Kawabata, T., *Site-Selective Catalysis*, Gewerbestrasse 11 CH-6330 Cham (ZG) Switzerland: Springer Interntional Publishing (2016).
- Sikervar, V.; Ueda, Y.; Kawabata, T., L-Trytophan *N,N'*-[[(2*S,SS*)-1-(4-pyridinyl)-2,5-pyrrolidinediyl]dicarbonyl]bis-3,3'-dioctyl Ester, *e-EROS Encyclopedia of Reagents for Organic Synthesis; John Wiley & Sons, Ltd.*, DOI: 10.1002/047084289X. rn01815.pub2 (2016).
- Yoshimura, T.; Kawabata, T., Lithium tert-Butyl (trimethylsilyl) amide, e-EROS Encyclopedia of Reagents for Organic Synthesis; John Wiley & Sons, Ltd., DOI: 10.1002/047084289X. rn01797 (2016).
- Ueda, Y.; Kawabata, T., D-2-Naphthylalanine N, N'-[[(2S,5S)-1-(4-pyridinyl)-2,5-pyrrolidinediyl]dicarbonyl]bis-3,3'-dioctyl Ester, e-EROS Encyclopedia of Reagents for Organic Synthesis; John Wiley & Sons, Ltd., DOI: 10.1002/047084289X. rn01986 (2016).

Koeduka, T.; Kajiyama, M.; Furuta, T.; Suzuki, H.; Tsuge, T.; Matsui, K., Benzenoid Biosynthesis in The Flowers of *Eriobotrya Japonica*: Molecular Cloning and Functional Characterization of *p*-Methoxybenzoic Acid Carboxyl Methyltransferase, *Planta*, **244**, 725-736 (2016).

Koeduka, T.; Kajiyama, M.; Furuta, T.; Suzuki, H.; Tsuge, T.; Matsui, K., Characterization of An O-Methyltransferase Specific to Guaiacoltype Benzenoids from The Flowers of Loquat (*Eriobotrya Japonica*), *J. Biosci. Bioeng.*, **122**, 679-684 (2016).

— Advanced Inorganic Synthesis —

Chen, L.; Nakamoto, M.; Haruta, M.; Nemoto, T.; Sato, R.; Kurata, H.; Teranishi, T., Tin Ion Directed Morphology Evolution of Copper Sulfide Nanoparticles and Tuning of Their Plasmonic Properties via Phase Conversion, *Langmuir*, **32**, 7582-7587 (2016).

Sakamoto, M.; Inoue, K.; Okano, M.; Saruyama, M.; Kim, S.; So, Y.-G.; Kimoto, K.; Kanemitsu, Y.; Teranishi, T., Light-Stimulated Carrier Dynamics of CuInS₂/CdS Heterotetrapod Nanocrystals, *Nanoscale*, **8**, 9517-9520 (2016).

Wu, H.-L.; Sato, R.; Yamaguchi, A.; Kimura, M.; Haruta, M.; Kurata, H.; Teranishi, T., Formation of Pseudomorphic Nanocages from Cu₂O Nanocrystals through Anion Exchange Reactions, *Science*, **351**, 1306-1310 (2016).

Azuma, Y.; Onuma, Y.; Sakamoto, M.; Teranishi, T.; Majima, Y., Rhombic Coulomb Diamonds in Single-Electron Transistor based on Au Nanoparticle Chemically Anchored at Both Ends, *Nanoscale*, **8**, 4720-4726 (2016).

Masuo, S.; Kanetaka, K.; Sato, R.; Teranishi, T., Direct Observation of Multiphoton Emission Enhancement from a Single Quantum Dot Using AFM Manipulation of a Cubic Gold Nanoparticle, *ACS Photonics*, **3**, 109-116 (2016).

DIVISION OF MATERIALS CHEMISTRY — Chemistry of Polymer Materials —

Huang, Y.; Sasano, T.; Tsujii, Y.; Ohno, K., Well-Defined Polymer-Brush-Coated Rod-Shaped Particles: Synthesis and Formation of Liquid Crystals, *Macromolecules*, **49**, 8430-8439 (2016).

Sakakibara, K.; Yano, H.; Tsujii, Y., Surface Engineering of Cellulose Nanofiber by Adsorption of Diblock Copolymer Dispersant for Green Nanocomposite Materials, *ACS APPL. MATER. INT.*, **8,** 24893-24900 (2016).

Goto, A.; Tsujii, Y., Surface-Initiated Living Radical Polymerizations Using Iodine, Organotellurium, and Organic catalysts, *Adv. Polym. Sci.*, **270**, 107-122 (2016).

Ferrier, R. C.; Huang, Y.; Ohno, K.; Composto, R. J., Dispersion of PMMA-grafted, Mesoscopic Iron-oxide Rods in Polymer Films, *Soft Matter*, **12**, 2550-2556 (2016).

Tom, J.; Ohno, K.; Perrier, S., Surface-Initiated SET Living Radical Polymerisation for the Synthesis of Silica-Polymer Core-Shell Nanoparticles, *Polym. Chem.*, **7**, 6075-6083 (2016).

Morinaga, T.; Honma, S.; Ishizuka, T.; Kamijo, T.; Sato, T.; Tsujii, Y., Synthesis of Monodisperse Silica Particles Grafted with Concentrated Ionic Liquid-Type Polymer Brushes by Surface-Initiated Atom Transfer Radical Polymerization for Use as a Solid State Polymer Electrolyte, *Polymers*, **8**, 141-146 (2016).

Horikawa, Y.; Imai, T.; Abe, K.; Sakakibara, K.; Tsujii, Y.; Mihashi, A.; Kobayashi, Y.; Sugiyama, J., Assessment of Endoglucanase Activity by Analyzing the Degree of Cellulose Polymerization and High-throughput Analysis by Near-infrared Spectroscopy, *Cellulose*, **23**, 1565-1572 (2016).

Li, L.; Nakaji-Hirabayashi, T.; Kitano, H.; Ohno, K.; Kishioka, T.; Usui, Y., Gradation of Proteins and Cells Attached to the Surface of Bio-inert Zwitterionic Polymer Brush, *Colloids Surf. B*, **144**, 180-187 (2016).

Kawata, Y.; Yamamoto, T.; Kihara, H.; Yamamura, Y.; Saito, K.; Ohno, K., Three Gel States of Colloidal Composites Consisting of Polymer-Brush-Afforded Silica Particles and a Nematic Liquid Crystal with Distinct Viscoelastic and Optical Properties, *ACS APPL. MATER. INT.*, **8**, 29649-29657 (2016).

Ogawa, Y.; Sakakibara, K.; Nakatsubo, F.; Kamitakahara, H.; Takano, T., Factors Affecting Photocurrent Generation Performances of Langmuir-Blodgett Films of Tetraphenylporphyrin-bound Acyl Celluloses, *Cellulose*, **23**, 3875-3883 (2016).

[Others]

Tsujii, Y.; Nakanishi, Y.; Ishige, R.; Ohno, K.; Morinaga, T.; Sato, T., Development of Novel Nano-systems for Electrochemical Devices by Hierarchizing Concentrated Polymer Brushes, *Intelligent Nanosystems for Energy, Information and Biological Technologies*, 195-215 (2016).

Tsujii, Y.; Sakakibara, K., Tribology Controlled by Polymer Brushes, J. Adhes. Soc. Jpn., **52**, 224-229 (2016) (in Japanese).

Sakakibara, K., Development and Application of Polymer Monolith Materials with Controlled Morphology, *Sen'i Gakkaishi*, **72**, 428-430 (2016) (in Japanese).

— Polymer Controlled Synthesis —

Alvarez, M. P.; Delgado, M. C. R.; Taravillo, M.; Baonza, V. G.; Navarrete, J. T. L.; Evans, P.; Jasti, R.; Yamago, S.; Kertesz, M.; Casado, J., The Raman Fingerprint of Cyclic Conjugation: the Case of the Stabilization of Cations and Dications in Cycloparaphenylenes, *Che. Sci.*, 7, 3494-3499 (2016).

Kayahara, E.; Patel, V. K.; Mercier, A.; Kundig, E. P.; Yamago, S., Regioselective Synthesis and Characterization of Multinuclear Convex-Bound Ruthenium-[n] Cycloparaphenylene (n=5 and 6) Complexes, *Angew. Chem. Int. Ed.*, **55**, 302-306 (2016).

Kayahara, E.; Kouyama, T.; Kato, T.; Yamago, S., Synthesis and Characterization of [n]CPP (n=5, 6, 8, 10, and 12) Radical Cation and Dications: Size-Dependent Absorption, Spin, and Charge Delocalization, *J. Am. Chem. Soc.*, **138**, 338-344 (2016).

Nakamura, Y.; Ogihara, T.; Yamago, S., Mechanism of Cu(I)/Cu(0)-Mediated Reductive Coupling Reactions of Bromine-Terminated Polyacrylates, Polymethacrylates, and Polystyrene, *ACS Macro Lett.*, **5**, 248-252 (2016).

- Nakamura, Y.; Lee, R.; Coote, M. L.; Yamago, S., Termination Mechanism of the Radical Polymerization of Acrylates, Macromol. Rapid Commun., 37, 506-513 (2016).
- Kuroda, Y.; Sakamoto, Y.; Suzuki, T.; Kayahara, E.; Yamago, S., Tetracyclo(2,7-carbazole)s: Diatropicity and Paratropicity of Inner Regions of Nanohoops, J. Org. Chem., 81, 3356-3363 (2016).
- Alvarez, M. P.; Qiu, L. L.; Taravillo, M.; Baonza, V. G.; Delgado, M. C. R.; Yamago, S.; Jasti, R.; Navarrete, J. T. L.; Casado, J.; Kertesz, M., From Linear to Cyclic Oligoparaphenylenes: Electronic and Molecular Changes Traced in the Vibrational Raman Spectra and Reformulation of the Bond Length Alternation Pattern, Phys. Chem. Chem. Phys., 18, 11683-11692 (2016).
- Kayahara, E.; Fukayama, K.; Nishinaga, T.; Yamago, S., Size Dependence of [n]Cycloparaphenylenes (n=5-12) in Electrochemical Oxidation, Chem. Asian. J., 11, 1793-1797 (2016).
- Kayahara, E.; Zhai, X.; Yamago, S., Synthesis and Physical Properties of [4]Cyclo-3,7-Dibenzo[b,d]Thiophene and Its S,S-Dioxide, Can. J. Chem., 10.1139/cjc-2016-0474 (2016).
- Fan, W.; Nakamura, Y.; Yamago, S., Synthesis of Multivalent Organotellurium Chain Transfer Agents by Post-modification and Their Applications in Living Radical Polymerization, Chem. Eur. J., **22,** 17006-17010 (2016).
- Nakamura, Y.; Ogihara, T.; Hatano, S.; Abe, M.; Yamago, S., Control of the Termination Mechanism in Radical Polymerization by Viscosity: Selective Disproportionation in Viscous Media, Chem. Eur. J., 10.1002/chem.201604659 (2016).

[Others]

Yamago, S., Radical Polymerization (3) Reversible Chain Transfer, Recent Controlled Polymerization: Excellent Control and Advanced Practical Use (Current View 20), 75-81 (2016) (in Japanese).

- Inorganic Photonics Materials -

- Matsuzaki, Y.; Shimo-Oka, T.; Tanaka, H.; Tokura, Y.; Semba, K.; Mizuochi, N., Hybrid Quantum Magnetic Field Sensor with an Electron Spin and a Nuclear Spin in Diamond, Phys. Rev. A, 94, [052330-1]-[052330-6] (2016).
- Fujiwara, M.; Yoshida, K.; Noda, T.; Takashima, H.; Schell, A. W.; Mizuochi, N.; Takeuchi, S., Manipulation of Single Nanodiamonds to Ultrathin Fiber-taper Nanofibers and Control of NV-spin States Toward Fiber-integrated λ-systems, Nanotechnology, 27, 455202 (2016).
- Fukui, N.; Morishita, H.; Kobayashi, S.; Miwa, S.; Mizuochi, N.; Suzuki, Y., Ferromagnetic Resonance Induced Electromotive Forces in Ni₈₁Fe₁₉|p-type Diamond, Solid state commun., 243, 44-48 (2016).
- Matsuzaki, Y.; Morishita, H.; Tashima, T.; Kakuyanagi, K.; Semba, K.; Munro, W. J.; Yamaguchi, H.; Mizuochi, N.; Saito, S., Optically Detected Magnetic Resonance of High-density Ensemble of NV Centers in Diamond, J. Phys.: Condens. Matter, 28, 275302 (2016).
- Doi, Y.; Fukui, T.; Kato, H.; Makino, T.; Yamasaki, S.; Tashima, T.; Morishita, H.; Miwa, S.; Jelezko, F.; Suzuki, Y.; Mizuochi, N., Pure Negatively Charged State of NV Center in n-type Diamond, Phys. Rev. B, 93, [081203-1]-[081203-6] (2016).

- Goto, M.; Nawaoka, K.; Miwa, S.; Hatanaka, S.; Mizuochi, N.; Suzuki, Y., Electric Field Modulation of Tunneling Anisotropic Magnetoresistance in Tunnel Junctions with Antiferromagnetic Electrodes, Jpn. J. Appl. Phys., 55, [080304-1]-[080304-4] (2016).
- Kuro, T.; Okada, G.; Kawaguchi, N.; Fujimoto, Y.; Masai, H.; Yanagida, T., Scintillation Properties of Rare-earth Doped NaPO₃-Al(PO₃)₃ Glasses, Opt. Mater., 62, 561-568 (2016).
- Tatsumi, H.; Okada, G.; Yanagida, T.; Masai, H., Radiophotoluminescence and Thermally-stimulated Luminescence of Ag-doped Li₃PO₄-Al(PO₃)₃ Glass, J. Lumin., 179, 545-549 (2016).
- Okada, G.; Masai, H.; Torimoto, A.; Kasap, S.; Yanagida, T., X-ray Induced Effects in Sm₃₊-doped ZnO-P₂O₅ Glass for Radiation Measurements, J. Ceram. Process. Res., 17, 148-151 (2016).
- Masai, H.; Ueda, Y.; Yanagida, T.; Okada, G.; Tokuda, Y., Photo- and Radioluminescence of Sn2+ Centers in Alkaline Earth-Substituted Zinc Phosphate Glass, Sensor. Mater., 28, 871-879 (2016).
- Masai, H.; Matsumoto, S.; Ueda, Y.; Koreeda, A., Correlation between Valence State of Tin and Elastic Modulus of Sn-doped Li₂O-B₂O₃-SiO₂ Glasses, J. Appl. Phys., 119, 185104 (2016).
- Masai, H.; Koreeda, A.; Fujii, Y.; Ohkubo, T.; Kohara, S., Photoluminescence of Sn2+-centre as Probe of Transient State of Supercooled Liquid, Opt. Mater. Express, 6, 1827-1836 (2016).
- Torimoto, A.; Masai, H.; Tokuda, Y.; Yanagida, T.; Okada, G.; Mibu, K., Correlation between the Emission Properties of Sn²⁺ Center and the Chemical Composition of ZnO-P₂O₅ Glasses, J. Ceram. Soc. Jpn., **124,** 554-558 (2016).
- Tatsumi, H.; Okada, G.; Yanagida, T.; Masai, H., Scintillation and Dosimeter Properties of Ce-doped Li₃PO₄-Al(PO₃)₃ Glasses, J. Ceram. Soc. Jpn., 124, 550-553 (2016).
- Yanagida, T.; Fujimoto, Y.; Masai, H., Radiation linduced Luminescence Properties of Pure and Sn-doped 60ZnO-40P₂O₅ Glass, Phys. Chem. Glasses, 57, 158-162 (2016).
- Tatsumi, H.; Okada, G.; Yanagida, T.; Masai, H., Radiophotoluminescence of Ag-doped Li₃PO₄-Al(PO₃)₃ Glass, Chem. Lett., 45, 280-282 (2016).
- Yanagida, T.; Ueda, J.; Masai, H.; Fujimoto, Y.; Tanabe, S., Optical and Scintillation Properties of Ce-doped 34Li₂O-5MgO-10Al₂O₃-51SiO₂ Glass, J. Non-Cryst. Solids, 431, 140-144 (2016).
- Masai, H.; Yanagida, T., Photoluminescence of ns2-type Centercontaining Zinc Borate Glasses, J. Non-Cryst. Solids, 431, 83-87
- Tokuda, Y.; Norikawa, Y.; Masai, H.; Ueda, Y.; Nihei, N.; Fujimura, S.; Ono, Y., Nuclear Magnetic Resonance Study of Cs Adsorption onto Clay Minerals, Radiological Issues for Fukushima's Revitalized Future, 3-11 (2016).

[Others]

Kobayashi, S.; Morishita, H.; Matsuzaki, Y.; Miwa, S.; Suzuki, Y.; Mizuochi, N., Enhancement of the Spin Coherence Time in NV Center Under Electric Field, NEW DIAMOND, 32, 11-15 (2016) (in Japanese).

- Nanospintronics -

Ogi, H.; Masuda, S.; Nagakubo, A.; Nakamura, N.; Hirao, M.; Kondou, K.; Ono, T., Impact of Interface Stiffness in Surface-wave Resonances on Nanostrip-attached Substrates, Phys. Rev. B, 93, [037204-1]-[037204-5] (2016).

Yoshimura, Y.; Kim, K. J.; Taniguchi, T.; Tono, T.; Ueda, K.; Hiramatsu, R.; Moriyama, T.; Yamada, K.; Nakatani, Y.; Ono, T., Soliton-like Magnetic Domain Wall Motion Induced by the Interfacial Dzyaloshinskii-Moriya Interaction, Nat. Phys., 12, 157-161 (2016).

Takeshita, S.; Matsuo, S.; Tanaka, T.; Nakaharai, S.; Tsukagoshi, K.; Moriyama, T.; Ono, T.; Arakawa, T.; Kobayashi, K., Anomalous Behavior of 1/f Noise in Graphene Near the Charge Neutrality Point, Appl. Phys. Lett., 108, [103106-1]-[103106-4] (2016).

Tanabe, K.; Matsumoto, R.; Ohe, J. I.; Murakami, S.; Moriyama, T.; Chiba, D.; Kobayashi, K.; Ono, T., Observation of Magnon Hall-like Effect for Sample-edge Scattering in Unsaturated YIG, Phys. Status Solidi B, 253, 783-787 (2016).

Kakizakai, H.; Ando, F.; Koyama, T.; Yamada, K.; Kawaguchi, M.; Kim, S.; Kim, K. J.; Moriyama, T.; Chiba, D.; Ono, T., Switching Local Magnetization by Electric-field-induced Domain Wall Motion, Appl. Phys. Exp., 9, [063004-1]-[063004-3] (2016).

Okuno, T.; Kim, K. J.; Tono, T.; Kim, S.; Moriyama, T.; Yoshikawa, H.; Tsukamoto, A.; Ono, T., Temperature Dependence of Magnetoresistance in GdFeCo/Pt Heterostructure, Appl. Phys. Exp., 9, [073001-1]-[073001-4] (2016).

Oda, K.; Moriyama, T.; Kawaguchi, M.; Kamiya, M.; Tanaka, K.; Kim, K. J.; Ono, T., Exchange Bias Controlled by Electric Current: Interplay of Joule Heating and the Induced Field, Jpn. J. Appl. Phys., **55**, [070304-1]-[070304-4] (2016).

Okuno, T.; Taniguchi, T.; Kim, S.; Baek, S. H. C.; Park, B. G.; Moriyama, T.; Kim, K. J.; Ono, T., Temperature Dependence of Spin Hall Magnetoresistance in W/CoFeB Bilayer, Jpn. J. Appl. Phys., 55, [080308-1]-[080308-4] (2016).

Stigloher, J.; Decker, M.; Korner, H. S.; Tanabe, K.; Moriyama, T.; Taniguchi, T.; Hata, H.; Madami, M.; Gubbiotti, G.; Kobayashi, K.; Ono, T.; Back, C. H., Snell's Law for Spin Waves, Phys. Rev. Lett., **117**, [037204-1]-[037204-5] (2016).

DIVISION OF BIOCHEMISTRY — Biofunctional Design-Chemistry —

Nakase, I.; Noguchi, K.; Fujii, I.; Futaki, S., Vectorization of Biomacromolecules into Cells Using Extracellular Vesicles with Enhanced Internalization Induced by Macropinocytosis, Sci. Rep., 6, [34937] (2016).

Tsuji, S.; Futaki, S.; Imanishi, M., Sequence-Specific Recognition of Methylated DNA by an Engineered Transcription Activator-Like Effector Protein, Chem. Commun., 52, 14238-14241 (2016).

Backlund, C.; Takeuchi, T.; Futaki, S.; Tew, G., Relating Structure and Internalization for ROMP-Based Protein Mimics, Biochim. Biophys. Acta., 1858, 1443-1450 (2016).

Kawaguchi, Y.; Takeuchi, T.; Kuwata, K.; Chiba, J.; Hatanaka, Y.; Nakase, I.; Futaki, S., Syndecan-4 Is a Receptor for Clathrin-Mediated Endocytosis of Arginine-Rich Cell-Penetrating Peptides, Bioconjug. Chem., 27, 1119-1130 (2016).

Murayama, T.; Pujals, S.; Hirose, H.; Nakase, I.; Futaki, S., Effect of Amino Acid Substitution in the Hydrophobic Face of Amphiphilic Peptides on Membrane Curvature and Perturbation: N-Terminal Helix Derived From Adenovirus Internal Protein VI As a Model, Biopolymers, 106, 430-439 (2016).

Takeuchi, T.; Futaki, S., Current Understanding of Direct Translocation of Arginine-Rich Cell-Penetrating Peptides and Its Internalization Mechanisms, Chem. Pharm. Bull., 64, 1431-1437 (2016).

Miyajima, R.; Tsuda, Y.; Inokuma, T.; Shigenaga, A.; Imanishi, M.; Futaki, S.; Otaka, A., Preparation of Peptide Thioesters from Naturally Occurring Sequences Using Reaction Sequence Consisting of Regioselective S-Cyanylation and Hydrazinolysis, Biopolymers, **106**, 531-546 (2016).

Motoyama, K.; Nishiyama, R.; Maeda, Y.; Higashi, T.; Kawaguchi, Y.; Futaki, S.; Ishitsuka, Y.; Kondo, Y.; Irie, T.; Era, T.; Arima H., Cholesterol-Lowering Effect of Octaarginine-Appended β-Cyclodextrin in Npc1-Trap-CHO Cells, Biol. Pharm. Bull., 39, 1823-1829 (2016).

Chemistry of Molecular Biocatalysts —

Kamiyama, A.; Nakajima, M.; Han, L.; Wada, K.; Mizutani, M.; Tabuchi, Y.; Kojima-Yuasa, A.; Matsui-Yuasa, I.; Suzuki, H.; Fukuyama, K.; Watanabe, B.; Hiratake, J., Phosphonate-Based Irreversible Inhibitors of Human γ-Glutamyl Transpeptidase (GGT). GGsTop is a Non-Toxic and Highly Selective Inhibitor with Critical Electrostatic Interaction with an Active-Site Residue Lys562 for Enhanced Inhibitory Activity, Bioorg. Med. Chem., 24, 5340-5352 (2016).

Tamura, K.; Hayashi, N.; George, J.; Toshikuni, N.; Arisawa, T.; Hiratake, J.; Tsuchishima, M.; Tsutsumi, M., GGsTop, a Novel and Specific γ-Glutamyl Transpeptidase Inhibitor, Protects Hepatic Ischemia-Reperfusion Injury in Rats, Am. J. Physiol. Gastrointest. Liver Physiol., 311, G305-G312 (2016).

— Molecular Biology —

Wu, Z.; Zhu, D.; Lin, X.; Miao, J.; Gu, L.; Deng, X.; Yang, Q.; Zhu, D.; Cao, X.; Tsuge, T.; Dean, C.; Aoyama, T.; Gu, H.; Qu, L.-J., RNA Binding Proteins RZ-1B and RZ-1C Play Critical Roles in Regulating Pre-mRNA Splicing and Gene Expression during Development in Arabidopsis, Plant Cell, 28, 55-73 (2016).

Tatsumi, K.; Yano, M.; Kaminade, K.; Sugiyama, A.; Sato, M.; Toyooka, K.; Aoyama, T.; Sato, F.; Yazaki, K., Characterization of Shikonin Derivative Secretion in Lithospermum erythrorhizon Hairy Roots as a Model of Lipid-Soluble Metabolite Secretion from Plants, Front. Plant Sci., 7, [01066-1]-[01066-11] (2016).

Nagata, C.; Miwa, C.; Tanaka, N.; Kato, M.; Suito, M.; Tsuchihira, A.; Sato, Y.; Segami, S.; Maeshima, M., A Novel-Type Phosphatidylinositol Phosphate-Interactive, Ca-Binding Protein PCaP1 in Arabidopsis thaliana: Stable Association with Plasma Membrane and Partial Involvement in Stomata Closure, J. Plant Res., 129, 539-550 (2016).

Koeduka, T.; Kajiyama, M.; Suzuki, H.; Furuta, T.; Tsuge, T.; Matsui, K., Benzenoid Biosynthesis in the Flowers of *Eriobotrya japonica*: Molecular Cloning and Functional Characterization of p-Methoxybenzoic Acid Carboxyl Methyltransferase, *Planta*, **244**, 725-736 (2016).

- Chemical Biology -

Usuda, K.; Biswas, T.; Yamaguchi, T.; Akagi, Y.; Yasui, K.; Uesugi, M.; Shimizu, I.; Hosokawa, S.; Nagasawa, K., Synthesis of Diastereomers of 1,3-cis-25-Dihydroxy-19-norvitamin D-3, *Chem. Pharm. Bull.*, **64**, 1990-1995 (2016).

Matsumoto, Y.; Ito, A.; Uesugi, M.; Kittaka, A., Efficient N-Acyldopamine Synthesis, *Chem. Pharm. Bull.*, **64**, 935-940 (2016).

Katsuda, Y.; Sato, S.; Asano, L.; Morimura, Y.; Furuta, T.; Sugiyama, H.; Hagihara, M.; Uesugi, M., A small Molecule that Represses Translation of G-Quadruplex-Containing mRNA, *J. Am. Chem. Soc.*, **138**, 9037-9040 (2016).

Biswas, T.; Akagi, Y.; Usuda, K.; Yasui, K.; Shimizu, I.; Okamoto, M.; Uesugi, M.; Hosokawa, S.; Nagasawa, K., Synthesis of 24,24-Difluoro-1,3-Cis-25-Dihydroxy-19-Norvitamin D3 Derivatives and Evaluation of their Vitamin D Receptor-Binding Affinity, *Biol. Pharm. Bull.*, **39**, 1387-1391 (2016).

Sakano, D.; Choi, S.; Kataoka, M.; Shiraki, N.; Uesugi, M.; Kume, K.; Kume, S., Dopamine D2 Receptor-Mediated Regulation Pancreatic Beta Cell Mass, *Stem Cell Rep.*, 7, 95-109 (2016).

Takemoto, Y.; Ito, A.; Niwa, H.; Okamura, M.; Fujiwara, T.; Hirano, T.; Handa, N.; Umehara, T.; Sonoda, T.; Ogawa, K.; Tariq, M.; Nishino, N.; Dan, S.; Kagechika, H.; Yamori, T.; Yokoyama, S.; Yoshida, M., Identification of Cyproheptadine as an Inhibitor of SET Domain Containing Lysine Methyltransferase 7/9 (Set7/9) That Regulates Estrogen-Dependent Transcription, *J. Med. Chem.*, **59**, 3650-60 (2016).

Ursu, A.; Illich, D. J.; Takemoto, Y.; Porfetye, A. T.; Zhang, M.; Brockmeyer, A.; Janning, P.; Watanabe, N.; Osada, H.; Vetter, I. R.; Ziegler, S.; Scholer, H. R.; Waldmann, H., Epiblastin A Induces Reprogramming of Epiblast Stem Cells Into Embryonic Stem Cells by Inhibition of Casein Kinase 1, *Cell Chem. Biol.*, **23**, 494-507 (2016).

[Others]

Takaya, J.; Uesugi, M., Chemical Biological Analysis of TRPA1 Activation Mechanism, *J. SYN. ORG. CHEM. JPN.*, **74**, 505-511 (2016) (in Japanese).

DIVISION OF ENVIRONMENTAL CHEMISTRY — Molecular Materials Chemistry —

Uratani, H.; Kubo, S.; Shizu, K.; Suzuki, F.; Fukushima, T.; Kaji, H., Detailed Analysis of Charge Transport in Amorphous Organic Thin Layer by Multiscale Simulation without Any Adjustable Parameters, *Sci. Rep.*, **6**, 39128 (2016).

Shimizu, T.; Kanamori, K.; Maeno, A.; Kaji, H.; Doherty, C. M.; Falcaro, P.; Nakanishi, K., Transparent, Highly Insulating Polyethyland Polyvinylsilsesquioxane Aerogels: Mechanical Improvements by Vulcanization for Ambient Pressure Drying, *Chem. Mater.*, **28**, 6860-6868 (2016).

Shizu, K.; Kaji, H., Organic Electroluminescent Materials Realizing Efficient Conversion from Electricity to Light, *J. Photopolym. Sci. Technol.*, **29**, 305-310 (2016).

Nishimura, H.; Fukushima, T.; Wakamiya, A.; Murata, Y.; Kaji, H., The Influence of Quasiplanar Structures of Partially Oxygen-Bridged Triphenylamine Dimers on the Properties of Their Bulk Films, *Bull. Chem. Soc. Jpn.*, **89**, 726-732 (2016).

Hayase, G.; Nonomura, K.; Kanamori, K.; Maeno, A.; Kaji, H.; Nakanishi, K., Boehmite Nanofiber-Polymethylsilsesquioxane Core-Shell Porous Monoliths for a Thermal Insulator under Low Vacuum Conditions, *Chem. Mater.*, **28**, 3237-3240 (2016).

Wada, Y.; Shizu, K.; Kubo, S.; Fukushima, T.; Miwa, T.; Tanaka, H.; Adachi, C.; Kaji, H., Highly Efficient Solution-Processed Host-Free Organic Light-Emitting Diodes Showing an External Quantum Efficiency of Nearly 18% with a Thermally Activated Delayed Fluorescence Emitter, *Appl. Phys. Express*, **9**, [032102-1]-[032102-3] (2016).

Hayase, G.; Kanamori, K.; Maeno, A.; Kaji, H.; Nakanishi, K., Dynamic Spring-Back Behavior in Evaporative Drying of Polymethylsilsesquioxane Monolithic Gels for Low-Density Transparent Thermal Superinsulators, *J. Non-Cryst. Solids*, **434**, 115-119 (2016).

Shizu, K.; Noda, H.; Tanaka, H.; Taneda, M.; Uejima, M.; Sato, T.; Tanaka, K.; Kaji, H.; Adachi, C., Highly Efficient Blue Electroluminescence Using Delayed-Fluorescence Emitters with Large Overlap Density between Luminescent and Ground States, *J. Phys. Chem. C*, **119**, 26283-26289 (2015).

Shimizu, T.; Kanamori, K.; Maeno, A.; Kaji, H.; Nakanishi, K., Transparent Ethylene-Bridged Polymethylsiloxane Aerogels and Xerogels with Improved Bending Flexibility, *LANGMUIR*, **32**, 13427-13434 (2016).

- Hydrospheric Environment Analytical Chemistry -

Nakashima, Y.; Shimizu, A.; Maruo, M.; Sohrin, Y., Trace Elements Influenced by Environmental Changes in Lake Biwa: (I) Seasonal Variations under Suboxic Hypolimnion Conditions during 2007 and 2009, *Limnol.*, **17**, 151-162 (2016).

Sohrin, Y.; Nakashima, Y.; Maruo, M., Trace Elements Influenced by Environmental Changes in Lake Biwa: (II) Chemical Variations in the Hypolimnion over the Last Half-Century, *Limnol.*, **17**, 163-173 (2016).

— Solution and Interface Chemistry —

Shimoaka, T.; Hasegawa, T., Molecular Structural Analysis of Hydrated Ethylene Glycol Accounting for the Antifreeze Effect by Using Infrared Attenuated Total Reflection Spectroscopy, *J. Mol. Liquids*, **223**, 621-627 (2016).

Shimoaka, T.; Tanaka, Y.; Shioya, N.; Morita, K.; Sonoyama, M.; Amii, H.; Takagi, T.; Kanamori, T.; Hasegawa, T., Surface Properties of a Single Perfluoroalkyl Group on Water Surfaces Studied by Surface Potential Measurements, *J. Colloid Interf. Sci.*, **483**, 353-359 (2016).

- Hada, M.; Shioya, N.; Shimoaka, T.; Eda, K.; Hada, M.; Hasegawa, T., Comprehensive Understanding of Structure-Controlling Factors of a Zinc Tetraphenylporphyrin Thin Film Using pMAIRS and GIXD Techniques, *Chem. Eur. J.*, **22**, 16539-16546 (2016).
- Zhang, R.; Murata, M.; Aharen, T.; Wakamiya, A.; Shimoaka, T.; Hasegawa, T.; Murata, Y., Synthesis of a Distinct Water Dimer Inside Fullerene C₇₀, *Nature Chem.*, **8**, 435-441 (2016).
- Wakai, C.; Shimoaka, T.; Hasegawa, T., Characterization of Adsorbed Molecular Water on the Surface of a Stretched Polytetrafluoroethylene Tape Analyzed by ¹H NMR, *J. Phys. Chem. B*, **120**, 2538-2543 (2016).
- Iwasa, J.; Kumazawa, K.; Aoyama, K.; Suzuki, H.; Norimoto, S.; Shimoaka, T.; Hasegawa, T., In Situ Observation of a Self-Assembled Monolayer Formation of Octadecyltrimethoxysilane on a Silicon Oxide Surface Using a High-Speed Atomic Force Microscope, *J. Phys. Chem. C*, **120**, 2807-2813 (2016).
- Noda, T.; Kondo, S.; Kado, S.; Hasegawa, T.; Yamada, N., Formation of Polyglycine II Structure from Fatty Acid Derivatives Containing Mono-, Di- and Tri-Glycinate, *KOBUNSHI RONBUNSHU*, **73**, 69-75 (2016) (in Japanese).

[Others]

- Shimoaka, T.; Hasegawa, T., Chemical Structural Analysis of Antifreeze Solution by Using Infrared Spectroscopy with an Aid of Chemometrics, *Transactions of The Research Institute of Oceanochemistry*, **29(2)**, 79-84 (2016) (in Japanese).
- Matsuno, T.; Tabuchi, Y.; Hosomi, R.; Hasegawa, T.; Kometani, N.; Tsuji, K., Sample Preparation for TXRF Analysis of Metal Particles in Used Machine Oils and Preliminary Research for Application of Principal Component Analysis, *Adv. X-ray Anal.*, **59**, 112-119 (2016).

- Molecular Microbial Science -

- Ito, T.; Gong, C.; Kawamoto, J.; Kurihara, T., Development of a Versatile Method for Targeted Gene Deletion and Insertion by Using the *pyrF* Gene in the Psychrotrophic Bacterium, *Shewanella living-stonensis* Ac10, *J. Biosci. Bioeng.*, **122**, 645-651 (2016).
- Ohke, Y.; Maruyama, S.; Tarui, J.; Wang, Y.; Kawamoto, J.; Kurihara, T., A Phosphoporin Homolog Plays a Critical Role in the Dissimilatory Iron-Respiration Linked to Iron (III) Reduction by a Cold-adapted Bacterium, *Shewanella livingstonensis* Ac10, *Trace Nutr. Res.*, 33, 35-42 (2016).
- Sugiura, M.; Park, J.; Kawamoto, J.; Esaki, N.; Kurihara, T., Regulatory Mechanism of Membrane Protein Production in an EPA-Producing Bacterium, *Shewanella livingstonensis* Ac10, *Trace Nutr. Res.*, **33**, 63-72 (2016).

[Others]

Kurihara, T., Pentose Phosphate Pathway; Heterolactic Fermentation Via Phosphoketolase Pathway; Stickland Reaction, *Applied Microbiology, 3rd Ed.*, 94-97 (2016) (in Japanese).

DIVISION OF MULTIDISCIPLINARY CHEMISTRY — Polymer Materials Science —

- Nishida, K.; Saiga, M.; Fujishima, Y.; Ogawa, H.; Kanaya, T., Quasi-Equilibrium Gelation Temperature of Aqueous Methylcellulose, *J. Fiber Sci. Technol.*, **72**, 251-257 (2016).
- Henmi, K.; Sato, H.; Matsuba, G.; Tsuji, H.; Nishida, K.; Kanaya, T.; Toyohara, K.; Oda, A.; Endou, K., Isothermal Crystallization Process of Poly(L lactic acid)/Poly(D lactic acid) Blends after Rapid Cooling from the Melt, *ACS Omega*, **1**, 476-482 (2016).
- Kanaya, T.; Ogino, Y.; Mizuike, A.; Miyazaki, T.; Inoue, R.; Nishida, K., A study on Polymerization-Induced Phase Separation (PIPS) of Epoxy Solution in Oligoethylene Glycol Aiming at Controlling the Characteristic Length, *J. Fiber Sci. Technol.*, **72**, 126-131 (2016).
- Hirai, Y.; Terashima, T.; Takenaka, M.; Sawamoto, M., Precision Self-Assembly of Amphiphilic Random Copolymers into Uniform and Self-Sorting Nanocompartments in Water, *Macromolecules*, **49**, 5084-5091 (2016).
- Wang, Y.-C.; Kim, M.-I.; Akasaka, S.; Saijo, K.; Hasegawa, H.; Hikima, T.; Takenaka. M., Fddd Structure in Polystyrene-block-polyisoprene Diblock Copolymer/Polystyrene Homopolymer Blends, *Macromolecules*, **49**, 2257-2261 (2016).
- Matsumoto, K.; Terashima, T.; Sugita, T.; Takenaka, M.; Sawamoto, M., Amphiphilic Random Copolymers with Hydrophobic/Hydrogen-Bonding Urea Pendants: Self-Folding Polymers in Aqueous and Organic Media, *Macromolecules*, **49**, 7917-7927 (2016).
- Ishige, R.; Ohta, N.; Ogawa, H.; Tokita, M.; Takahara, A., Fully Liquid-Crystalline ABA Triblock Copolymer of Fluorinated Side-Chain Liquid-Crystalline A Block and Main-Chain Liquid-Crystalline, B Block: Higher Order Structure in Bulk and Thin Film States, *Macromolecules*, **49**, 6061-6074 (2016).
- Ogawa, H.; Takenaka, M.; Miyazaki, T.; Fujiwara, A.; Lee, B.; Shimokita, K.; Nishibori, E.; Takata, M., Direct Observation on Spin-Coating Process of PS-*b*-P2VP Thin Films, *Macromolecules*, **49**, 3471-3477 (2016).

[Others]

Nishida, K., Observation of Structures in µm-scale by Small-angle Light Scattering, in Ishii, T. Ed., *Information for Consultants in Colloid and Surface Science*, 493-498 (2016) (in Japanese).

- Molecular Rheology -

- Ankiewicz, S.; Orbey, H.; Watanabe, H.; Lentzakis, H.; Dealy, J., On the Use of Continuous Relaxation Spectra to Characterize Model Polymers, *J. Rheol.*, **60**, 1115-1120 (2016).
- Doi, Y.; Takano, A.; Matsushita, Y., Synthesis and Characterization of Dumbbell-Shaped Polystyrene, *Polymer*, **106**, 8-13 (2016).
- Matsumiya, Y.; Watanabe, H.; Urakawa, O.; Inoue, T., Experimental Test for Viscoelastic Relaxation of Polyisoprene Undergoing Monofunctional Head-to-Head Association and Dissociation, *Macromolecules*, **49(18)**, 7088-7095 (2016).

Matsumiya, Y.; Watanabe, H., Nonlinear Stress Relaxation of Miscible Polyisoprene/Poly(p-tert-butylstyrene) Blends in Pseudomonodisperse State, *Macromolecules*, **49(12)**, 4544-4556 (2016).

Kwon, Y.; Matsumiya, Y.; Watanabe, H., Viscoelastic and Orientational Relaxation of Linear and Ring Rouse Chains Undergoing Reversible End-Association and Dissociation, *Macromolecules*, **49(9)**, 3593-3607 (2016).

Doi, Y.; Iwasa, Y.; Watanabe, K.; Nakamura, M.; Takano, A.; Takahashi, Y.; Matsushita, Y., Synthesis and Characterization of Comb-Shaped Ring Polystyrenes, *Macromolecules*, **49(8)**, 3109-3115 (2016).

Rakkapao, N.; Watanabe, H.; Matsumiya, Y.; Masubuchi, Y., Dielectric Relaxation and Ionic Conductivity of a Chitosan/Poly(ethylene oxide) Blend Doped with Potassium and Calcium Cations, *J. Soc. Rheol. Japan*, **44(2)**, 89-97 (2016).

Masubuchi, Y.; Watanabe, H., Stress-Optical Relationship in Bead-Spring Simulations for Entangled Polymers under Start-up Shear Flows, *J. Soc. Rheol. Japan*, **44(1)**, 65-68 (2016).

ADVANCED RESEARCH CENTER FOR BEAM SCIENCE — Particle Beam Science —

Fuwa, Y.; Iwashita, Y.; Tongu, H.; Inoue, S.; Hashida, M.; Sakabe, S.; Okamura, M.; Yamazaki, A., RF Synchronized Short Pulse Laser Ion Source, *Rev. Sci. Instrum.*, **87**, [02A911-1]-[02A911-4] (2016).

Iwashita, Y.; Tongu, H.; Fuwa, Y.; Ichikawa, M., Compact Permanent Magnet H⁺ ECR Ion Source with Pulse Gas Valve, *Rev. Sci. Instrum.*, **87(2)**, [02A718-1]-[02A718-3] (2016).

Imajo, S.; Mishima, K.; Kitaguchi, M.; Iwashita, Y.; Yamada, N. L.; Hino, M.; Oda, T.; Ino, T.; Shimizu, H. M.; Yamashita, S.; Katayama, R., Pulsed Ultra-cold Neutron Production Using a Doppler Shifter at J-PARC, *Prog. Theor. Exp. Phys.*, **2016(1)**, 013C02-1-22 (2016).

Otani, M.; Mibe, T.; Yoshida, M.; Hasegawa, K.; Kondo, Y.; Hayashizaki, N.; Iwashita, Y.; Iwata, Y.; Kitamura, R.; Saito, N., Interdigital H-mode Drift-tube Linac Design with Alternative Phase Focusing for Muon Linac, *Phys. Rev. Accel. Beams*, **19**, [040101-1]-[040101-8] (2016).

Adeva, B.; Afanasyev, L.; Allkofer, Y.; Amsler, C.; Anania, A.; Aogaki, S.; Benelli, A.; Brekhovskikh, V.; Caragheorgheopol, Gh.; Cechak, T.; Chiba, M.; Chliapnikov, P.; Ciocarlan, C.; Constantinescu, S.; Detraz, C.; Doskarova, P.; Drijard, D.; Dudarev, A.; Duma, M.; Dumitriu, D.; Fluerasu, D.; Gorin, A.; Gorchakov, O.; Gritsay, K.; Guaraldo, C.; Gugiu, M.; Hansroul, M.; Hons, Z.; Horikawa, S.; Iwashita, Y.; Karpukhin, V.; Kluson, J.; Kobayashi, M.; Kruglov, V.; Kruglova, L.; Kulikov, A.; Kulish, E.; Kuptsov, A.; Lamberto, A.; Lanaro, A.; Lednicky, R.; Mariñas, C.; Martincik, J.; Nemenov, L.; Nikitin, M.; Okada, K.; Olchevskii, V.; Pentia, M.; Penzo, A.; Plo, M.; Prusa, P.; Rappazzo, G. F.; Romero Vidal, A.; Ryazantsev, A.; Rykalin, V.; Saborido, J.; Schacher, J.; Sidorov, A.; Smolik, J.; Takeutchi, F.; Tauscher, L.; Trojek, T.; Trusov, S.; Ullaland, O.; Urban, T.; Vrba, T.; Yazkov, V.; Yoshimura, Y.; Zhabitsky, M.; Zrelov, P., Upgraded DIRAC Spectrometer at CERN PS for the Investigation of $\pi\pi$ and πK Atoms, Nucl. Instrum. Meth., A839, 52-85 (2016).

[Others]

Otani, M.; Mibe, T.; Yoshida, M.; Hasegawa, K.; Kondo, Y.; Hayashizaki, N.; Iwashita, Y.; Iwata, Y.; Kitamura, R.;Saito, N., APF IH-DTL Design for the Muon LINAC in the J-PARC Muon g-2/EDM Experiment, *Proceedings of the 7th International Particle Accelerator Conference, IPAC'16*, 1539-1542 (2016).

Otani, M.; Mibe, T.; Naito, F.; Saito, N.; Yoshida, M.; Hasegawa, K.; Ito, T.; Kondo, Y.; Hayashizaki, N.; Iwashita, Y.; Iwata, Y.; Kitamura, R., Development of Muon LINAC for the Muon g-2/EDM Experiment at J-PARC, *Proceedings of the 7th International Particle Accelerator Conference, IPAC'16*, 1543-1546 (2016).

Kubo, T.; Saeki, T.; Cenni, E.; Hajima, R.; Sawamura, M.; Hokonohara, H.; Iwashita, Y.; Tongu, H., Multipactor Simulations in 650 MHz Superconducting Spoke Cavity for an Electron Accelerator, *Proceedings of the 7th International Particle Accelerator Conference, IPAC'16*, 2161-2163 (2016).

Saeki, T.; Hayano, H.; Kubo, T.; Iwashita, Y.; Oikawa, H., Construction of Measurement System for Superconducting Characteristics on Thin-film Samples at KEK, *Proceedings of the 7th International Particle Accelerator Conference, IPAC'16*, 2167-2169 (2016).

Tongu, H.; Hokonohara, H.; Iwashita, Y.; Hajima, R.; Sawamura, M.; Hayano, H.; Kubo, T.; Saeki, T.; Yamamoto, Y., Study on Nondestructive Inspections for Super-conducting Cavity, *Proceedings of the 7th International Particle Accelerator Conference, IPAC'16*, 2174-2176 (2016).

Sawamura, M.; Hajima, R.; Hokonohara, H.; Iwashita, Y.; Tongu, H.; Kubo, T.; Saeki, T., Fabrication of Superconducting Spoke Cavity for Laser Compton Scattered Photon Sources, *Proceedings of the 7th International Particle Accelerator Conference, IPAC'16*, 2177-2179 (2016).

Kondo, Y.; Hasegawa, K.; Ito, T.; Artikova, S.; Otani, M.; Mibe, T.; Naito, F.; Yoshida, M.; Kitamura, R.; Iwashita, Y.; Iwata, Y.; Hayashizaki, N.; Saito, N., The Muon Linac for the Precise Measurement of Muon g-2/EDM at J-PARC, *Proceedings of the 13th Annual Meeting of Particle Accelerator Society of Japan*, MOOM04 (2016) (in Japanese).

Fuwa, Y.; Iwashita, Y.; Inoue, S.; Hashida, M.; Sakabe, S.; Tongu, H., Development of RF Synchronized Short Pulse Laser Ion Source, *Proceedings of the 13th Annual Meeting of Particle Accelerator Society of Japan*, WEOM04 (2016) (in Japanese).

Imajo, S.; Iwashita, Y.; Mishima, K.; Kitaguchi, M.; Shimizu, H. M.; Ino, T.; Yamashita, S., Present Status of Ultracold Neutron Rebuncher, *Proceedings of the 13th Annual Meeting of Particle Accelerator Society of Japan*, MOP009 (2016) (in Japanese).

Oikawa, H.; Higashiguchi, T.; Iwashita, Y.; Hino, M.; Kubo, T.; Kato, S.; Saeki, T.; Hayano, H., Design Study of Niobium Mushroom-shaped Cavity for Evaluating RF Critical Magnetic Field of Thin-Film Superconductor, *Proceedings of the 13th Annual Meeting of Particle Accelerator Society of Japan*, MOP021 (2016) (in Japanese).

Iwashita, Y.; Tongu, H.; Fuwa, Y.; Saeki, T.; Kubo, T., Characteristics Measurements on Sc Thin Foil, *Proceedings of the 13th Annual Meeting of Particle Accelerator Society of Japan*, TUP014 (2016) (in Japanese).

- Otani, M.; Hasegawa, K.; Hayashizaki, N.; Iwashita, Y.; Iwata, Y.; Kitamura, R.; Kondo, Y.; Mibe, T.; Saito, N.; Yoshida, M., Interdigital H-mode Drift-tube Linac Design with Alternative Phase Focusing for Muon Linac, *Proceedings of the 13th Annual Meeting of Particle Accelerator Society of Japan*, TUP017 (2016) (in Japanese).
- Sawamura, M.; Hajima, R.; Saeki, T.; Kubo, T.; Iwashita, Y.; Tongu, H.; Hokonohara, H., Press Forming Test of Superconducting Spoke Cavity, *Proceedings of the 13th Annual Meeting of Particle Accelerator Society of Japan*, TUP024 (2016) (in Japanese).
- Fuwa, Y.; Iwashita, Y.; Tongu, H., Simulation of Klystron Beam Focusing by Permanent Magnet, *Proceedings of the 13th Annual Meeting of Particle Accelerator Society of Japan*, TUP030 (2016) (in Japanese).
- Imajo, S.; Iwashita, Y.; Mishima, K.; Kitaguchi, M.; Shimizu, H. M.; Ino, T.; Yamashita, S., Development of Ultracold Neutron Accelerator for Time Focusing of Pulsed Neutrons, *Proceedings of 28th Linear Accelerator Conference*, MOPRC001 (2016).

- Laser Matter Interaction Science -

- Fuwa, Y.; Iwashita, Y.; Tongu, H.; Inoue, S.; Hashida, M.; Sakabe, S.; Okamura, M.; Yamazaki, A., RF Synchronized Short Pulse Laser Ion Source, *Rev. Sci. Instrum.*, **87**, [02A911-1]-[02A911-4] (2016).
- Hashida, M.; Nishii, T.; Miyasaka, Y.; Sakagami, H.; Shimizu, M.; Inoue, S.; Sakabe, S., Orientation of Periodic Grating Structures Controlled by Double-pulse Irradiation, *Appl. Phys. A*, **122**, [1]-[5] (2016).
- Furukawa, Y.; Sakata, R.; Konishi, K.; Ono, K.; Matsuoka, S.; Watanabe, K.; Inoue, S.; Hashida, M.; Sakabe, S., Demonstration of Periodic Nanostructure Formation with Less Ablation by Double-pulse Laser Irradiation on Titanium, *Appl. Phys. Lett.*, **108**, [264101-1]-[264101-4] (2016).
- Inoue, S.; Maeda, K.; Tokita, S.; Mori, K.; Teramoto, K.; Hashida, M.; Sakabe, S., Single Plasma Mirror Providing 10(4) Contrast Enhancement and 70% Reflectivity for Intense Femtosecond Lasers, *Appl. Opt.*, **55**, [5647]-[5651] (2016).
- Inoue, S.; Maeda, K.; Tokita, S.; Mori, K.; Teramoto, K.; Hashida, M.; Sakabe, S., Single Plasma Mirror Providing 10(4) Contrast Enhancement and 70% Reflectivity for Intense Femtosecond Lasers: Publisher's Note (vol 55, pg 5647, 2016), *Appl. Opt.*, **55**, [6435]-[6435] (2016).
- Arikawa, Y.; Kojima, S.; Morace, A.; Sakata, S.; Gawa, T.; Taguchi, Y.; Abe, Y.; Zhang, Z.; Vaisseau, X.; Lee, S. H.; Matsuo, K.; Tosaki, S.; Hata, M.; Kawabata, K.; Kawakami, Y.; Ishida, M.; Tsuji, K.; Matsuo, S.; Morio, N.; Kawasaki, T.; Tokita, S.; Nakata, Y.; Jitsuno, T.; Miyanaga, N.; Kawanaka, J.; Nagatomo, H.; Yogo, A.; Nakai, M.; Nishimura, H.; Shiraga, H.; Fujioka, S.; Azechi, H.; Sunahara, A.; Johzaki, T.; Ozaki, T.; Sakagami, H.; Sagisaka, A.; Ogura, K.; Pirozhkov, A. S.; Nishikino, M.; Kondo, K.; Inoue, S.; Teramoto, K.; Hashida, M.; Sakabe, S., Ultrahigh-contrast Kilojoule-class Petawatt LFEX Laser Using a Plasma Mirror, *Appl. Opt.*, **55**, [6850]-[6857] (2016).
- Hashida, M.; Miyasaka, Y.; Nishii, T.; Shimizu, M.; Inoue, S.; Sakabe, S., Threshold Fluence for Femtosecond Laser Nanoablation for Metals, *Electron. Comm. Jpn.*, **99**, [88]-[95] (2016).

- Electron Microscopy and Crystal Chemistry -

- Tsai, C. H.; Chen, S. Y.; Song, J. M.; Haruta, M.; Kurata, H., Effect of Ag Templates on the Formation of Au-Ag Hollow/Core-Shell Nanostructures, *Nanoscale Res. Lett.*, **10**, [438-1]-[438-9] (2015).
- Zhu, Y.; Shimizu, T.; Kitajima, T.; Morisato, K.; Moitra, N.; Brun, N.; Kiyomura, T.; Kanamori, K.; Takeda, K.; Kurata, H.; Tafu, M.; Nakanishi, K., Synthesis of Robust Hierarchically Porous Zirconium Phosphate Monolith for Efficient Ion Adsorption, *New J. Chem.*, **39**, 2444-2450 (2015).
- Zhu, Y.; Morisato, K.; Hasegawa, G.; Moitra, N.; Kiyomura, T.; Kurata, H.; Kanamori, K.; Nakanishi, K., High-performance Liquid Chromatography Separation of Unsaturated Organic Compounds by a Monolithic Silica Column Embedded with Silver Nanoparticles, *J. Sep. Sci.*, **38**, 2841-2847 (2015).
- Zhu, Y.; Yoneda, K.; Kanamori, K.; Takeda, K.; Kiyomura, T.; Kurata, H.; Nakanishi, K., Hierarchically Porous Titanium Phosphate Monoliths and Their Crystallization Behavior in Ethylene Glycol, *New J. Chem.*, **40**, 4153-4159 (2016).
- Shirai, K.; Sugimoto, T.; Watanabe, K.; Haruta, M.; Kurata, H.; Matsumoto, Y., Effect of Water Adsorption on Carrier Trapping Dynamics at the Surface of Anatase TiO₂ Nanoparticles, *Nano Lett.*, **16**, 1323-1327 (2016).
- Wu, H. L.; Sato, R.; Yamaguchi, A.; Kimura, M.; Haruta, M.; Kurata, H.; Teranishi, T., Formation of Pseudomorphic Nanocages from Cu₂O Nanocrystals through Anion Exchange Reactions, *Science*, **351**, 1306-1310 (2016).
- Kan, D.; Aso, R.; Sato, R.; Haruta, M.; Kurata, H.; Shimakawa, Y., Tuning Magnetic Anisotropy by Interfacially Engineering the Oxygen Coordination Environment in a Transition Metal Oxide, *Nature Mater.*, **15**, 432-437 (2016).
- Hasegawa, G.; Kanamori, K.; Kiyomura, T.; Kurata, H.; Abe, T.; Nakanishi, K., Hierarchically Porous Carbon Monoliths Comprising Ordered Mesoporous Nanorod Assemblies for High-Voltage Aqueous Supercapacitors, *Chem. Mater.*, **28**, 3944-3950 (2016).
- Haruta, M.; Hosaka, Y.; Ichikawa, N.; Saito, T.; Shimakawa, Y.; Kurata, H., Determination of Elemental Ratio in an Atomic Column by Electron Energy Loss Spectroscopy, *ACS Nano*, **10**, 6680-6684 (2016).
- Shoji, S.; Ogawa, T.; Hashishin, T.; Ogasawara, S.; Watanabe, H.; Usami, H.; Tamiaki, H., Nanotubes of Biomimetic Supramolecules Constructed by Synthetic Metal Chlorophyll Derivatives, *Nano Lett.*, **16**, 3650-3654 (2016).
- Chen, L. H.; Sakamoto, M.; Haruta, M.; Nemoto, T.; Sato, R.; Kurata, H.; Teranishi, T., Tin Ion Directed Morphology Evolution of Copper Sulfide Nanoparticles and Tuning of Their Plasmonic Properties via Phase Conversion, *Langmuir*, **32**, 7582-7587 (2016).

— Structural Molecular Biology —

Fujii, T.; Sato, A.; Okamoto, Y.; Yamauchi, T.; Kato, S.; Yoshida, M.; Oikawa, T.; Hata, Y., The Crystal Structure of Maleylacetate Reductase from *Rhizobium* sp. strain MTP-10005 Provides Insights into the Reaction Mechanism of Enzymes in its Original Family, *Proteins*, **84**, 1029-1042 (2016).

INTERNATIONAL RESEARCH CENTER FOR ELEMENTS SCIENCE

— Synthetic Organotransformation —

Yoshida, R.; Isozaki, K.; Yokoi, T.; Yasuda, N.; Sadakane, K.; Iwamoto, T.; Takaya, H.; Nakamura, M., ONO-Pincer Ruthenium Complex-Bound Norvaline for Efficient Catalytic Oxidation of Methoxybenzenes with Hydrogen Peroxide, *Org. Biomol. Chem.*, **14**, 7468-7479 (2016).

Naito, W.; Yasuda, N.; Morimoto, T.; Shigeta, Y.; Takaya, H.; Hisaki, I.; Maeda, H., Doubly *N*-Methylated Porphyrinoids, *Org. Lett.*, **18**, 3006-3009 (2016).

Bando, Y.; Haketa, Y.; Sakurai, T.; Matsuda, W.; Seki, S.; Takaya, H.; Maeda, H., Ion-Pairing Assemblies Based on Pentacyano-Substituted Cyclopentadienide as a π -Electronic Anion, *Chem. Eur. J.*, **22**, 7843-7850 (2016).

Yamaguchi, A.; Isozaki, K.; Nakamura, M.; Takaya, H.; Watanabe, T., Discovery of 12-Mer Peptides That Bind to Wood Lignin, *Sci. Rep.*, **6**, [21833-1]-[21833-11] (2016).

Isozaki, K.; Yokoi, T.; Yoshida, R.; Ogata, K.; Hashizume, D.; Yasuda, N.; Sadakane, K.; Takaya, H.; Nakamura, M., Synthesis and Applications of ONO-Pincer Ruthenium-Complex-Bound Norvalines, *Chem. Asian. J.*, **11**, 1076-1091 (2016).

[Others]

Takaya, H.; Nakajima, S.; Nakamura, M., Synchrotron X-ray Absorption Spectroscopy and Solution-Phase Structure of Organoiron Catalysts, *Chemistry & Chemical Industry*, **69**, 391-393 (2016) (in Japanease).

- Advanced Solid State Chemistry -

Seki, H.; Hosaka, Y.; Saito, T.; Mizumaki, M.; Shimakawa, Y., Ferromagnetism Induced by Substitution of the Iron(IV) Ion by an Unusual High-valence Nickel(IV) Ion in Antiferromagnetic SrFeO₃, *Angew. Chem. Int. Ed.*, **55**, 1360-1363 (2016).

Kawakami, T.; Sekiya, Y.; Miura, A.; Kobayashi, K.; Tokumichi, K.; Yamada, I.; Mizumaki, M.; Kawamura, N.; Shimakawa, Y.; Ohishi, Y.; Hirao, N.; Ishimatsu, N.; Hayashi, N.; Takano, M., Two-step Suppression of Charge Disproportionation in CaCu₃Fe₄O₁₂ under High Pressure, *J. Phys. Soc. Jpn.*, **85**, [034716-1]-[034716-6] (2016).

Kan, D.; Orikasa, Y.; Nitta, K.; Tanida, H.; Kurosaki, R.; Nishimura, T.; Sasaki, T.; Guo, H.; Ozaki, Y.; Uchimoto, Y.; Shimakawa, Y., Overpotential-induced Introduction of Oxygen Vacancy in La_{0.67}Sr_{0.33}MnO₃ Surface and Its Impact on Oxygen Reduction Reaction Catalytic Activity in Alkaline Solution, *J. Phys. Chem. C*, **120**, 6006-6010 (2016).

Heo, Y.; Kan, D.; Shimakawa, Y.; Seidel, J., Resistive Switching Properties of Epitaxial BaTiO_{3-δ}Thin Films Tuned by After-growth Oxygen Cooling Pressure, *Phys. Chem. Chem. Phys.*, **18**, 197-204 (2016).

Kan, D.; Aso, R.; Sato, R.; Haruta, M.; Kurata, H.; Shimakawa, Y., Tuning Magnetic Anisotropy by Interfacially Engineering the Oxygen Coordination Environment in a Transition Metal Oxide, *Nature Materials*, **15**, 432-437 (2016).

Haruta, M.; Hosaka, Y.; Ichikawa, N.; Saito, T.; Shimakawa, Y.; Kurata, H., Determination of Elemental Ratio in an Atomic Column by Electron Energy Loss Spectroscopy, *ACS Nano*, **10**, 6680-6684 (2016).

Xiong, P.; Seki, H.; Guo, H.; Hosaka, Y.; Saito, T.; Mizumaki, M.; Shimakawa, Y., Geometrical Spin Frustration of Unusually High Valence Fe⁵⁺ in the Double Perovskite La₂LiFeO₆, *Inorg. Chem.*, **55**, 6218-6222 (2016).

Hosaka, Y.; Ichikawa, N.; Saito, T.; Mizumaki, M.; Shimakawa, Y., A Layered Double Perovskite Ca₂FeMnO₆ with Unusually High Valence Fe⁴⁺ Obtained by Low-Temperature Topotactic Oxidation, *J. Jpn. Soc. Powder Powder Metallurgy*, **63**, 605-608 (2016).

Seki, H.; Saito, T.; Shimakawa, Y., High Pressure Synthesis of SrFe_{1-x}Ni_xO₃, *J. Jpn. Soc. Powder Powder Metallurgy*, **63**, 609-612 (2016)

Kan, D.; Wakabayashi, Y.; Tajiri, H.; Shimakawa, Y., Interfacially Engineered Oxygen Octahedral Rotations and Their Impact on Strain Relief in Coherently Grown SrRuO₃ Films, *Phys. Rev. B*, **94**, [024112-1]-[024112-6] (2016).

Hosaka, Y.; Ichikawa, N.; Saito, T.; Attfield, J. P.; Shimakawa, Y., Charge and Spin Order in the Perovskite CaFe_{0.5}Mn_{0.5}O₃: Charge Disproportionation Behavior of Randomly Arranged Fe⁴⁺, *Phys. Rev. B*, **94**, [104429-1]-[104429-5] (2016).

Gustin, L.; Hosaka, Y.; Tassel, C.; Aharen, T.; Shimakawa, Y.; Kageyama, H.; Wiley, J. B., From Tetrahedral to Octahedral Iron Coordination: Layer Compression in Topochemically Prepared FeLa₂Ti₃O₁₀, *Inorg. Chem.*, **55**, 11529-11537 (2016).

Manabe, K.; Ichikawa, N.; Saito, T.; Kan, D.; Shimakawa, Y., Crystal Structures and B-site Cation Valences of A-site Layered Perovskites, GdBa(Fe_{0.5}Mn_{0.5})₂O_x (x=5,6), *J. Jpn. Soc. Powder Powder Metallurgy*, **63**, 33-37 (2016) (in Japanese).

Kan, D.; Shimakawa, Y., Direct Observation and Engineering of Oxygen Coordination Environments in Oxide Heterostructures, *J. Jpn. Soc. Powder Powder Metallurgy*, **63**, 829-834 (2016) (in Japanese).

[Others]

Shimakawa, Y., A Novel Half-metallic Transition-metal Oxide with a High Transition Temperature: Control of the Spin Structure by Cation Ordering in Oxides, *Oyo Buturi*, **85**, 212-217 (2016) (in Japanese).

— Organometallic Chemistry —

Nakajima, Y.; Tsuchimoto, T.; Chang, Y.-H.; Takeuchi, K.; Ozawa, F., Reactions of [Cu(X)(BPEP-Ph)] (X = PF₆, SbF₆) with Silyl Compounds. Cooperative Bond Activation Involving Noncoordinating Anions, *Dalton Trans.*, **45**, 2079-2084 (2016).

Chang, Y.-H.; Tanigawa, I.; Taguchi, H.; Takeuchi, K.; Ozawa, F., Iridium(I) Complexes Bearing a Noninnocent PNP-Pincer-Type Phosphaalkene Ligand: Catalytic Application in the Base-Free *N*-Alkyl ation of Amines with Alcohols, *Eur. J. Inorg. Chem.*, **2016**, 754-760 (2016).

- Iizuka, E.; Wakioka, M.; Ozawa, F., Mixed-Ligand Approach to Palladium-Catalyzed Direct Arylation Polymerization: Effective Prevention of Structural Defects Using Diamines, *Macromolecules*, **49**, 3310-3317 (2016).
- Taguchi, H.; Sasaki, D.; Takeuchi, K.; Tsujimoto, S.; Matsuo, T.; Tanaka, H.; Yoshizawa, K.; Ozawa, F., Unsymmetrical PNP-Pincer Type Phosphaalkene Ligands Protected by a Fused-Ring Bulky Eind Group: Synthesis and Applications to Rh(I) and Ir(I) Complexes, *Organometallics*, **35**, 1526-1533 (2016).
- Pouliot, J.-R.; Wakioka, M.; Ozawa, F.; Li, Y.-N.; Leclerc, M., Structural Analysis of Poly(3-hexylthiophene) Prepared via Direct Heteroarylation Polymerization, *Macromol. Chem. Phys.*, **217**, 1493-1500 (2016).
- Ozawa, F.; Nakajima, Y., PNP-Pincer-Type Phosphaalkene Complexes of Late Transition Metals, *Chem. Rec.*, **16**, 2314-2323 (2016).
- Takeuchi, K.; Taguchi, H.; Tanigawa, I.; Tsujimoto, S.; Matsuo, T.; Tanaka, H.; Yoshizawa, K.; Ozawa, F., A Square-Planar Complex of Platinum(0), *Angew. Chem. Int. Ed.*, **55**, 15347-15350 (2016).

— Nanophotonics —

- Le, P. Q.; Nakaike, Y.; Wakamiya, A.; Kanemitsu, Y., Free Excitons and Exciton-Phonon Coupling in CH₃NH₃PbI₃ Single Crystals Revealed by Photocurrent and Photoluminescence Measurements at Low Temperatures, *J. Phys. Chem. Lett.*, **7**, 4905-4910 (2016).
- Tex, D. M.; Akahane, K.; Kanemitsu, Y., Intrinsic Trade-off between Up-Conversion and Trapping Rates in InAs Quantum Dots for Intermediate-Band Solar Cells, *Phys. Rev. Applied*, **6**, [044003-1]-[044003-10] (2016).
- Ibuki, H.; Ihara, T.; Kanemitsu, Y., Spectral Diffusion of Emissions of Excitons and Trions in Single CdSe/ZnS Nanocrystals: Charge Fluctuations in and around Nanocrystals, *J. Phys. Chem. C*, **120**, 23772-23779 (2016).
- Yamashita, D.; Handa, T.; Ihara, T.; Tahara, H.; Shimazaki, A.; Wakamiya, A.; Kanemitsu, Y., Charge Injection at the Heterointerface in Perovskite CH₃NH₃PbI₃ Solar Cells Studied by Simultaneous Microscopic Photoluminescence and Photocurrent Imaging Spectroscopy, *J. Phys. Chem. Lett.*, **7**, 3186-3191 (2016).
- Ihara, T., Biexciton Cascade Emission Reveals Absolute Absorption Cross Section of Single Semiconductor Nanocrystals, *Phys. Rev. B*, **93**, [235442-1]-[235442-7] (2016).
- Le, P. Q.; Yamada, Y.; Nagai, M.; Maruyama, N.; Wakamiya, A.; Kanemitsu, Y., Free Carriers versus Excitons in CH₃NH₃PbI₃ Perovskite Thin Films at Low Temperatures: Charge Transfer from the Orthorhombic Phase to the Tetragonal Phase, *J. Phys. Chem. Lett.*, **7**, 2316-2321 (2016).
- Yamada, Y.; Yamada, T.; Shimazaki, A.; Wakamiya, A.; Kanemitsu, Y., Interfacial Charge-Carrier Trapping in CH₃NH₃PbI₃-Based Heterolayered Structures Revealed by Time-Resolved Photoluminescence Spectroscopy, *J. Phys. Chem. Lett.*, **7**, 1972-1977 (2016).

- Sakamoto, M.; Inoue, K.; Okano, M.; Saruyama, M.; Kim, S.; So, Y.-G.; Kimoto, K.; Kanemitsu, Y.; Teranishi, T., Light-Stimulated Carrier Dynamics of CuInS₂/CdS Heterotetrapod Nanocrystals, *Nanoscale*, **8**, 9517-9520 (2016).
- Handa, T.; Tex, D. M.; Shimazaki, A.; Aharen, T.; Wakamiya, A.; Kanemitsu, Y., Optical Characterization of Voltage-Accelerated Degradation in CH₃NH₃PbI₃ Perovskite Solar Cells, *Opt. Express*, **24**, A917-A924 (2016).
- Zhu, L.; Mochizuki, T.; Yoshita, M.; Chen, S.; Kim, C.; Akiyama, H.; Kanemitsu, Y., Conversion Efficiency Limits and Bandgap Designs for Multi-Junction Solar Cells with Internal Radiative Efficiencies below Unity, *Opt. Express*, **24**, A740-A751 (2016).
- Zhu, L.; Yoshita, M.; Nakamura, T.; Imaizumi, M.; Kim, C.; Mochizuki, T.; Chen, S.; Kanemitsu, Y.; Akiyama, H., Characterization and Modeling of Radiation Damages via Internal Radiative Efficiency in Multi-Junction Solar Cells, *Proc. SPIE*, **9743**, [97430U-1]-[97430U-7] (2016).
- Yoshita, M.; Zhu, L.; Kim, C.; Mochizuki, T.; Nakamura, T.; Imaizumi, M.; Chen, S.; Kubota, H.; Kanemitsu, Y.; Akiyama, H., Calibration Standards and Measurement Accuracy of Absolute Electroluminescence and Internal Properties in Multi-Junction and Arrayed Solar Cells, *Proc. SPIE*, **9743**, [97430D-1]-[97430D-6] (2016).
- Ibuki, H.; Ihara, T.; Kanemitsu, Y., Photoluminescence Blinking and Spectral Diffusion of Single CdSe/ZnS Nanocrystals: Charge Fluctuation Effects, *Proc. SPIE*, **9758**, [97580R-1]-[97580R-8] (2016).
- Handa, T.; Okano, M.; Tex, D. M.; Shimazaki, A.; Aharen, T.; Wakamiya, A.; Kanemitsu, Y., Carrier Injection and Recombination Processes in Perovskite CH₃NH₃PbI₃ Solar Cells Studied by Electroluminescence Spectroscopy, *Proc. SPIE*, **9745**, [97451I-1]-[97451I-7] (2016).
- Tahara, H.; Endo, M.; Wakamiya, A.; Kanemitsu, Y., Experimental Evidence of Localized Shallow States in Orthorhombic Phase of CH₃NH₃PbI₃ Perovskite Thin Films Revealed by Photocurrent Beat Spectroscopy, *J. Phys. Chem. C*, **120**, 5347-5352 (2016).
- Tahara, H.; Kanemitsu, Y., Observation of Localized States in Bulk GaAs Using Ultrafast Photocurrent Beat Spectroscopy, *Appl. Phys. Express*, **9**, [032403-1]-[032403-4] (2016).
- Mochizuki, T.; Kim, C.; Yoshita, M.; Mitchell, J.; Lin, Z.; Chen, S.; Takato, H.; Kanemitsu, Y.; Akiyama, H., Solar-Cell Radiance Standard for Absolute Electroluminescence Measurements and Open-Circuit Voltage Mapping of Silicon Solar Modules, *J. Appl. Phys.*, **119**, [034501-1]-[034501-8] (2016).
- Yamada, T.; Yamada, Y.; Nishimura, H.; Nakaike, Y.; Wakamiya, A.; Murata, Y.; Kanemitsu, Y., Fast Free-Carrier Diffusion in CH₃NH₃PbBr₃ Single Crystals Revealed by Time-Resolved One-and Two-Photon Excitation Photoluminescence Spectroscopy, *Adv. Electron. Mater.*, **2**, [1500290-1]-[1500290-5] (2016).
- Zhu, L.; Yoshita, M.; Chen, S.; Nakamura, T.; Mochizuki, T.; Kim, C.; Imaizumi, M.; Kanemitsu, Y.; Akiyama, H., Characterizations of Radiation Damages in Multi-Junction Solar Cells Focused on Subcell Internal Luminescence Quantum Yields via Absolute Electroluminescence Measurements, *IEEE J. Photovolt.*, **6**, 777-782 (2015).

Nishihara, T.; Yamada, Y.; Okano, M.; Kanemitsu, Y., Dynamics of the Lowest-Energy Excitons in Single-Walled Carbon Nanotubes under Resonant and Nonresonant Optical Excitation, *J. Phys. Chem. C*, **119**, 28654-28659 (2015).

BIOINFORMATICS CENTER

— Chemical Life Science —

Yoshizawa, A. C., Which Database to Use? —Confusions and Puzzles in Database Search and Sequence Analysis—, *Proteome Lett.*, **1**, 63-80 (2016) (in Japanese).

Roux, S.; Brum, J. R.; Dutilh, B. E.; Sunagawa, S.; Duhaime, M. B.; Loy, A.; Poulos, B. T.; Solonenko, N.; Lara, E.; Poulain, J.; Pesant, S.; Kandels-Lewis, S.; Dimier, C.; Picheral, M.; Searson, S.; Cruaud, C.; Alberti, A.; Duarte, C. M.; Gasol, J. M.; Vaque, D.; Tara Oceans Coordinators; Bork, P.; Acinas, S. G.; Wincker, P.; Sullivan M. B., Ecogenomics and Potential Biogeochemical Impacts of Globally Abundant Ocean Viruses, *Nature*, **537**, 689-693 (2016).

Ben Halima, N.; Khemakhem, B.; Fendri, I.; Ogata, H.; Baril, P.; Pichon, C.; Abdelkafi, S., Identification of a New Oat Beta-Amylase by Functional Proteomics, *Biochim. Biophys. Acta*, **1864**, 52-61 (2016).

Moriya, Y.; Yamada, T.; Okuda, S.; Nakagawa, Z.; Kotera, M.; Tokimatsu, T.; Kanehisa, M.; Goto, S., Identification of Enzyme Genes Using Chemical Structure Alignments of Substrate-Product Pairs, *J. Chem. Inf. Model*, **56**, 510-516 (2016).

Mihara, T.; Nishimura, Y.; Shimizu, Y.; Nishiyama, H.; Yoshikawa, G.; Uehara, H.; Hingamp, P.; Goto, S.; Ogata, H., Linking Virus Genomes with Host Taxonomy, *Viruses*, **8**, 66 (2016).

Mihara, T.; Nasr-Eldin, M. A.; Chatchawankanphanich, O.; Bhunchoth, A.; Phironrit, N.; Kawasaki, T.; Nakano, M.; Fujie, M.; Ogata, H.; Yamada, T., A Ralstonia Solanacearum Phage \$\phi\$RP15 is Closely Related to Viunalikeviruses and Encodes 19 TRNA-Related Sequences, *Virology Reports*, 6, 61-73 (2016).

Guidi, L.; Chaffron, S.; Bittner, L.; Eveillard, D.; Larhlimi, A.; Roux, S.; Darzi, Y.; Audic, S.; Berline, L.; Brum, J. R.; Coelho, L. P.; Espinoza, J. C. I.; Malviya, S.; Sunagawa, S.; Dimier, C.; Kandels-Lewis, S.; Picheral, M.; Poulain, J.; Searson, S.; Stemmann, L.; Not, F.; Hingamp, P.; Speich, S.; Follows, M.; Karp-Boss, L.; Boss, E.; Ogata, H.; Pesant, S.; Weissenbach, J.; Wincker, P.; Acinas, S. G.; Bork, P.; de Vargas, C.; Iudicone, D.; Sullivan, M. B.; Raes, J.; Karsenti, E.; Bowler, C.; Gorsky, G., Plankton Networks Driving Carbon Export in the Oligotrophic Ocean, *Nature*, **532**, 465-470 (2016)

Lescot, M.; Hingamp, P.; Kojima, K. K.; Villar, E.; Romac, S.; Veluchamy, A.; Boccara, M.; Jaillon, O.; Iudicone, D.; Bowler, C.; Wincker, P.; Claverie, J. M.; Ogata, H., Reverse Transcriptase Genes are Highly Abundant and Transcriptionally Active in Marine Plankton Assemblages, *ISME J.*, **10**, 1134-1146 (2016).

Blanc-Mathieu, R.; Ogata, H., DNA Repair Genes in the Megavirales Pangenome, *Curr. Opin. Microbiol.*, **31**, 94-100 (2016).

Bhunchoth, A.; Blanc-Mathieu, R.; Mihara, T.; Nishimura, Y.; Askora, A.; Phironrit, N.; Leksomboon, C.; Chatchawankanphanich, O.; Kawasaki, T.; Nakano, M.; Fujie, M.; Ogata, H.; Yamada, T., Two Asian Jumbo Phages, φRSL2 and φRSF1, Infect Ralstonia Solanacearum and Show Common Features of φKZ-Related Phages, *Virology*, **494**, 56-66 (2016).

Forster, D.; Dunthorn, M.; Mahe, F.; Dolan, J. R.; Audic, S.; Bass, D.; Bittner, L.; Boutte, C.; Christen, R.; Claverie, J. M.; Decelle, J.; Edvardsen, B.; Egge, E.; Eikrem, W.; Gobet, A.; Kooistra, W. H. C. F.; Logares, R.; Massana, R.; Montresor, M.; Not, F.; Ogata, H.; Pawlowski, J.; Pernice, M. C.; Romac, S.; Shalchian-Tabrizi, K.; Simon, N.; Richards, T. A.; Santini, S.; Sarno, D.; Siano, R.; Vaulot, D.; Wincker, P.; Zingone, A.; de Vargas, C.; Stoeck, T., Benthic Protists: The Under-Charted Majority, *FEMS Microbiol. Ecol.*, 92, (2016).

Takami, H.; Taniguchi, T.; Arai, W.; Takemoto, K.; Moriya, Y.; Goto, S., An Automated System for Evaluation of the Potential Functionome: MAPLE version 2.1.0, *DNA Res.*, **23**, 467-475 (2016).

Kotera, M.; Goto, S., Metabolic Pathway Reconstruction Strategies for Central Metabolism and Natural Product Biosynthesis, *Biophysics and Physicobiology*, **13**, 195-205 (2016).

— Mathematical Bioinformatics —

Hasegawa, T.; Niida, A.; Mori, T.; Shimamura, T.; Yamaguchi, R.; Miyano, S.; Akutsu, T.; Imoto, S., A Likelihood-free Filtering Method via Approximate Bayesian Computation in Evaluating Biological Simulation Models, *Comput. Stat. Data An.*, **94**, 63-74 (2016).

Jindalertudomdee, J.; Hayashida, M.; Zhao, Y.; Akutsu, T., Enumeration Method for Tree-like Chemical Compounds with Benzene Rings and Naphthalene Rings by Breadth-first Search Order, *BMC Bioinformatics*, **17**, [113-1]-[113-16] (2016).

Nacher, J. C.; Akutsu, T., Minimum Dominating Set-based Methods for Analyzing Biological Networks, *Methods*, **102**, 57-63 (2016).

Hayashida, M.; Akutsu, T., Complex Network-based Approaches to Biomarker Discovery, *Biomarkers in Medicine*, **10**, 621-632 (2016).

Li, C.; Chang, C. H.; Porebski, B. T.; Hayashida, M.; Akutsu, T.; Song, J.; Buckle, A. M., Critical Evaluation of in silico Methods for Prediction of Coiled-coil Domains in Proteins, *Brief. Bioinform.*, **17**, 270-282 (2016).

Ishitsuka, M.; Akutsu, T.; Nacher, J. C., Critical Controllability in Proteome-wide Protein Interaction Network Integrating Transcriptome, *Sci. Rep.*, **6**, [23541-1]-[23541-13] (2016).

Koyano, H.; Hayashida, M.; Akutsu, T., Maximum Margin Classifier Working in a Set of Strings, *P. R. Soc. A*, **472**, [20150551-1]-[20150551-17] (2016).

Takemoto, K.; Akutsu, T., Analysis of the Effect of Degree Correlation on the Size of Minimum Dominating Sets in Complex Networks, *PLoS ONE*, **11**, [e0157868-1]-[e0157868-11] (2016).

Jindalertudomdee, J.; Hayashida, M.; Akutsu, T., Enumeration Method for Structural Isomers Containing User-defined Structures Based on Breadth-first Search Approach, *J. Comput. Biol.*, **23**, 625-640 (2016).

Hou, W.; Tamura, T.; Ching, W. K.; Akutsu, T., Finding and Analyzing the Minimum Set of Driver Nodes in Control of Boolean Networks, *Adv. Complex Syst.*, **19**, [1650006-1]-[1650006-32] (2016).

- He, F.; Hanai, A.; Nagamochi, H.; Akutsu, T., Enumerating Naphthalene Isomers of Tree-like Chemical Graphs, *Proc. 9th International Joint Conference on Biomedical Engineering Systems and Technologies*, 258-265 (2016).
- Tamura, T.; Lin, C. Y.; Yang, J. M.; Akutsu, T., Finding Influential Genes Using Gene Expression Data and Boolean Models of Metabolic Networks, *Proc. IEEE 16th International Conference on Bioinformatics and Bioengineering*, 57-63 (2016).
- Jindalertudomdee, J.; Hayashida, M.; Song, J.; Akutsu, T., Host-pathogen Protein Interaction Prediction Based on Local Topology Structures of a Protein Interaction Network, *Proc. IEEE 16th International Conference on Bioinformatics and Bioengineering*, 7-12 (2016).
- Cheng, X.; Mori, T.; Qiu, Y.; Ching, W. K.; Akutsu, T., Exact Identification of the Structure of a Probabilistic Boolean Network from Samples, *IEEE/ACM Transactions on Computational Biology and Bioinformatics*, **13**, 1545-5963 (2016).
- Bao, Y.; Hayashida, M.; Akutsu, T., LBSizeCleav: Improved Support Vector Machine (SVM)-based Prediction of Dicer Cleavage Sites Using Loop/Bulge Length, *BMC Bioinformatics*, **17**, [487-1]-[487-11] (2016).

[Others]

Akutsu, T., Analyses of Genomic and Chemical Information Using Mathematical Methods and Artificial Intelligence, *Transactions of the Research Institute of Oceanochemistry*, **29**, 85-91 (2016) (in Japanese).

- Bio-knowledge Engineering -

- Yotsukura, S.; Karasuyama, M.; Takigawa, I.; Mamitsuka, H., A Bioinformatics Approach for Understanding Genotype-phenotype Correlation in Breast Cancer, *Big Data Analytics in Genomics*, 397-428 (2016).
- Gao, J.; Yamada, M.; Kaski, S.; Mamitsuka, H.; Zhu, S., A Robust Convex Formulations for Ensemble Clustering, *Proceedings of the 25th International Joint Conference on Artificial Intelligence (IJCAI 2016)*, 1476-1482 (2016).
- Wicker, N.; Nguyen, C. H.; Mamitsuka, H., Some Properties of a Dissimilarity Measure for Labeled Graphs, *Publications Mathématiques de Besançon: Algèbre et Théorie des Nombres*, 81-90 (2016).
- Peng, S.; You, R.; Wang, H.; Zhai, C.; Mamitsuka, H.; Zhu, S., DeepMeSH: Deep Semantic Representation for Improving Large-scale MeSH Indexing, Bioinformatics (Proceedings of the 24th International Conference on Intelligent Systems for Molecular Biology (ISMB 2016)), 32(12), i70-i79 (2016).
- Yuan, Q.-J.; Gao, J.; Wu, D.; Zhang, S.; Mamitsuka, H.; Zhu, S., DrugE-Rank: Improving Drug-Target Interaction Prediction of New Candidate Drugs or Targets by Ensemble Learning to Rank, *Bioinformatics (Proceedings of the 24th International Conference on Intelligent Systems for Molecular Biology (ISMB 2016))*, **32(12)**, i18-i27 (2016).

- Natsume-Kitatani, Y.; Mamitsuka, H., Classification of Promoters based on the Combination of Core Promoter Elements Exhibits Different Histone Modification Patterns, *PLoS One*, **11(3)**, e0151917 (2016).
- Mohamed, A.; Nguyen, C. H.; Mamitsuka, H., NMRPro: An Integrated Web Component for Interactive Processing and Visualization of NMR Spectra, *Bioinformatics*, **32(13)**, 2067-2068 (2016).
- Shinkai-Ouchi, F.; Koyama, S.; Ono, Y.; Hata, S.; Ojima, K.; Shindo, M.; duVerle, D.; Kitamura, F.; Doi, N.; Takigawa, I.; Mamitsuka, H.; Sorimachi, H., Predictions of Cleavability of Calpain Proteolysis by Quantitative Structure-Activity Relationship Analysis Using Newly Determined Cleavage Sites and Catalytic Efficiencies of an Oligopeptide Array, *Molecular and Cellular Proteomics*, **15**, 1262-1280 (2016).
- Johnston, I.; Hancock, T.; Mamitsuka, H.; Carvalho, L., Gene-proximity Models for Genome-Wide Association Studies, *Annals of Applied Statistics*, **10(3)**, 1217-1244 (2016).
- Nguyen, C. H.; Mamitsuka, H., New Resistance Distances with Global Information on Large Graphs, *JMLR Workshop and Conference Proceedings (Proceedings of the Nineteenth International Conference on Artificial Intelligence and Statistics (AISTATS 2016)*), **51**, 639-647 (2016).
- Nakamura, A.; Takigawa, I.; Tosaka, H.; Kudo, M.; Mamitsuka, H., Mining Approximate Patterns with Frequent Locally Optimal Occurrences, *Discrete Applied Mathematics*, **200**, 123-152 (2016).
- Mohamed, A.; Nguyen, C. H.; Mamitsuka, H., Current Status and Prospects of Computational Resources for Natural Product Dereplication: A Review, *Briefings in Bioinformatics*, **17(2)**, 309-321 (2016).
- Iwata, T.; Yamada, M., Multi-view Anomaly Detection via Robust Probabilistic Latent Variable Models, *Advances in Neural Information Processing Systems 30 (NIPS2016)* (2016).
- Kozareva, Z.; Yamada, M., Which Tumblr Post Should I Read Next?, *Proceedings of the Annual Meeting of the Association for Computational Linguistics (ACL 2016)*, 332-336 (2016).
- Chang, Y.; Tang, J.; Yin, D.; Yamada, M.; Liu, Y., Timeline Summarization with Publications Life Cycle Models, *Proceedings of International Joint Conference on Artificial Intelligence (IJCAI 2016)*, 3698-3704 (2016).
- Gao, J.; Yamada, M.; Kaski, S.; Mamitsuka, H.; Zhu, S., A Robust Convex Formulation for Ensemble Clustering, *Proceedings of International Joint Conference on Artificial Intelligence (IJCAI 2016)*, 1476-1482 (2016).
- Wang, Y.; Yin, D.; Luo, J.; Wang, P.; Yamada, M.; Chang, Y.; Mei, Q., Beyond Ranking: Optimizing Whole-Page Presentation, *Proceedings of the 9th ACM International Conference on Web Search and Data Mining (WSDM 2016)*, 103-112 (2016).

[Others]

Xu, Y.; Luo, C.; Mamitsuka, H.; Zhu, S., MetaMHCpan, A Meta Apporach for Pan-specific MHC Peptide Binding Prediction, *Methods in Molecular Biology*, **1404**, 753-760 (2016).

HAKUBI PROJECT

- Algorithmic Graph Theory with Applications to Bioinformatics
- Jansson, J.; Shen, C.; Sung, W.-K., Improved Algorithms for Constructing Consensus Trees, Journal of the ACM, 63, Article 28
- Jansson, J.; Sung, W.-K.; Vu, H.; Yiu, S.-M., Faster Algorithms for Computing the R* Consensus Tree, Algorithmica, 76, 1224-1244
- Asahiro, Y.; Jansson, J.; Miyano, E.; Ono, H., Degree-Constrained Graph Orientation: Maximum Satisfaction and Minimum Violation, Theory of Computing Systems, 58, 60-93 (2016).
- Mori, T.; Takasu, A.; Jansson, J.; Hwang, J.; Tamura, T.; Akutsu, T., Similar Subtree Search Using Extended Tree Inclusion (extended abstract), Proc. of the 2016 IEEE 32nd International Conference on Data Engineering (ICDE 2016), 1558-1559 (2016).
- Jansson, J.; Sung, W.-K., Minimal Phylogenetic Supertrees and Local Consensus Trees, Proc. of the 41st International Symposium on Mathematical Foundations of Computer Science (MFCS 2016), LIPIcs, Article 53, 53:1-53:14 (2016).
- Jansson, J.; Sung, W.-K., Algorithms for Combining Rooted Triplets into a Galled Phylogenetic Network, in Kao, Ming-Yang (ed.), Encyclopedia of Algorithms (Second Edition), U.S.A., Springer -Verlag New York, 48-52 (2016).
- Jansson, J., Directed Perfect Phylogeny (Binary Characters), Ibid., 553-556 (2016).
- Jansson, J.; Sung, W.-K., Maximum Agreement Supertree, Ibid., 1224-1227 (2016).
- Jansson, J., Perfect Phylogeny (Bounded Number of States), Ibid., 1550-1553 (2016).
- Jansson, J., Phylogenetic Tree Construction from a Distance Matrix, Ibid., 1564-1567 (2016).

INTERNATIONAL RESEARCH COLLABORATIONS

[Australia]

RMIT University, School of Aerospace, Mechanical and Manufacturing Engineering

The University of Sydney, Key Centre for Polymers & Colloids

Monash University, Faculty of Pharmacy and Pharmaceutical Sciences

[Canada]

Université Laval, Département de Chimie

University of Waterloo, Department of Chemical Engineering

[China, P.R.]

Peking University, College of Life Science

Fujian Agriculture and Forestry University, Haixia Institute of Science and Technology

Chinese Academy of Science, Institute of Genetics and Developmental Biology

National Plant Gene Research Center

Fudan University, School of Computer Science

Fudan University, Shanghai Key Lab of Intelligent Information Processing

[Finland]

Aalto University, Department of Computer Science, School of

[France]

Université de Renne, Sciences Chimiques de Rennes

Aix Marseille University, Mediterranean Institute of Oceanography

CNRS

Université de Lille 1, Laboratoire Painlevé

[Germany]

Rheinische Friedrich-Wilhelms-Universität Bonn, Institut für Anorganische Chemie

Friedrich-Alexander-Universität Erlangen-Nürnberg, Department of Chemistry and Pharmacy, Interdisciplinary Center for Molecular Materials (ICMM)

The European Molecular Biology Laboratory

[Korea, R.]

The Catholic University of Korea, Department of Chemistry/Display Research Center

[Norway]

University of Bergen, Department of Biology

[Spain]

Universidad Autónoma de Madrid, Departamento de Química Orgánica

Universitat Rovira i Virgili, Departament de Química Física i Inorgànica

[Taiwan]

National Taiwan University of Science and Technology, Graduate Institute of Applied Science and Technology

National Taiwan University of Science and Technology, Department of Materials Science and Engineering

National Chung Hsing University, Department of Materials Science and Engineering

U.K.

The University of Warwick, Department of Chemistry

John Innes Centre for Plant Science Research, Department of Cell and Developmental Biology

The University of Edinburgh, Centre for Science at Extreme Conditions (CSEC) and School of Chemistry

[U.S.A.]

Columbia University, Department of Chemistry

University of Pennsylvania, Department of Materials Science and Engineering

University of Massachusetts, Department of Polymer Science and Engineering

The Ohio State University, Department of Microbiology

Boston University, Department of Mathematics and Statistics

^{*}The list shows the institutions with which papers are co-authored.

SELECTED GRANTS

DIVISION OF SYNTHETIC CHEMISTRY

— Organoelement Chemistry —

Tokitoh, N.

Creation of Novel Catalysts Centered on the Coordination Diversity of Heavy Typical Elements

Grant-in-Aid for Scientific Research on Innovative Area "Stimuliresponsive Chemical Species for the Creation of Functional Molecules"

28 June 2012-31 March 2017

Tokitoh, N

Electron-state Control of Aromatic Compounds Containing Heavier Group 14 Elements by Substituent introduction and Element Substitution

Grant-in-Aid for Scientific Research (B)

1 April 2013-31 March 2016

Tokitoh, N.

Synthesis and Properties of Alumoles Having an Aluminum–Halogen Bond

Grant-in-Aid for Challenging Exploratory Research

1 April 2014-31 March 2016

Tokitoh, N.

Synthesis and Properties of Phenyl Anion Analogs Containing Heavier Group 14 Elements

Grant-in-Aid for Scientific Research (B)

1 April 2016-31 March 2019

Sasamori, T.

Construction of [2]Ferrocenophanes Linked by π -Bond between Heavier Group 14 Elements and Control of Their Ring-opening Polymerization

Grant-in-Aid for Scientific Research on Innovative Area "Emergent Chemistry of Nano-scale Molecular Systems" and "New Polymeric Materials Based on Element-Blocks"

1 April 2013-31 March 2017

Sasamori, T.

Development of Transformations of Small Molecules and Multicomponent Couplings Utilizing Low-valent Compounds of Heavier Group 14 Elements

Grant-in-Aid for Scientific Research (B)

1 April 2015-31 March 2018

Sasamori, T.

Construction of d- π Electron Systems Containing Heavier Group 14 Elements and Their Functionalization

Grant-in-Aid for Challenging Exploratory Research

1 April 2015-31 March 2017

Mizuhata, Y.

Construction of Silicon-containing Dehydroannulenes and Their Aromaticity and Antiaromaticity

Grant-in-Aid for Scientific Research (C)

1 April 2014-31 March 2017

- Structural Organic Chemistry -

Murata. Y.

Synthesis of Tailor-made Nanocarbons and Their Application to Electronic Devices

Grant-in-Aid for Scientific Research (A)

1 April 2011-31 March 2016

Murata, Y.

Molecular Interface Science of π -Conjugated Carbon Complexes on Non-Equilibrated States

PRESTO (Precursory Research for Embryonic Science and Technology), JST

1 October 2012-31 March 2016

Murata, Y.

Spherical π -Figuration Based on Functionalization of Sub-Nano Space

Grant-in-Aid for Scientific Research on Innovative Areas " π -System Figuration"

1 April 2015-31 March 2017

Murata, Y.

Functional Molecular Systems Based on Dynamic Behavior of Active Species

Grant-in-Aid for Scientific Research on Innovative Areas "Stimuliresponsive Chemical Species for the Creation of Functional Molecules"

1 April 2015-31 March 2017

Murata, Y.

Creation of New Reactive Chemical Species by the Ultimate Steric Protection

Grant-in-Aid for Challenging Exploratory Research

1 April 2015-31 March 2017

Wakamiya, A.

Development of Organic Dyes Based on Fine Tuning of π -Orbitals using DFT Calculations

PRESTO (Precursory Research for Embryonic Science and Technology), JST

1 October 2010-31 March 2016

Abbreviations and acronyms

JST: Japan Science and Technology Agency

MEXT: Ministry of Education, Culture, Sports, Science and Technology

METI: Ministry of Economy, Trade and Industry

NEDO: New Energy and Industrial Technology Development Organization

Wakamiya, A.

Creation of Wireless Electric Power Supply Center of Innovation Program (COI) 1 October 2013-31 March 2022

Wakamiya, A.

High Dimensional Structural Control of π -Conjugated Systems and Their Functionalization

Grant-in-Aid for Scientific Research (B)

1 April 2014-31 March 2017

Wakamiya, A.

Development of High Performance and Environmentally Friendly Perovskite Type Solar Cells

Advanced Low Carbon Technology Research and Development Program (ALCA)

16 November 2016-31 March 2021

Murata, M.

Organization of Nanocarbon Molecules Based on Metal Coordination Grant-in-Aid for Challenging Exploratory Research

1 April 2014-31 March 2016

Murata, M.

Development of Neutral Thermoelectric Materials Based on π -Extended Metal-Bis(dithiolene) Complexes

PRESTO (Precursory Research for Embryonic Science and Technology), JST

1 October 2016-31 March 2020

Exploratory Studies on Materials for Energy Conversion Based on Efficient Synthesis of π -Conjugated Multimetallic Complexes Grant-in-Aid for Challenging Exploratory Research 1 April 2016-31 March 2018

Murata, M.

Development of Functional Materials Based on Efficient Synthesis of PAHs Containing Pentagon Rings Grant-in-Aid for Scientific Research (B)

1 April 2016-31 March 2019

Murata, M.

Exploration of Functions of Cyclopenta-Fused Polycyclic Aromatic Hydrocarbons CP-PAHs

Research Encouragement Grants, The Asahi Glass Foundation 1 April 2016-31 March 2018

- Synthetic Organic Chemistry -

Kawabata, T.

Regioselective Molecular Transformation Based on Organocatalytic Molecular Recognition

Grant-in-Aid for Scientific Research on Innovative Area

1 October 2011-31 March 2015

Kawabata, T.

Regioselective Molecular Transformation of Multifunctionalized Molecules

Grant-in-Aid for Scientific Research (S)

1 April 2014-31 March 2018

Furuta, T.

Direct Intra and Intermolecular Aldol Reaction by Catalytic Discrimination of Aldehydes

Grant-in-Aid for Scientific Research (C)

1 April 2014-31 March 2017

Ueda, Y.

Site-Selective Molecular Transformation Promoted by Anion-Exchange of Cationic Intermediates in Nucleophilic Catalysis Grant-in-Aid for Young Scientists (B) 1 April 2015-31 March 2017

Synthesis of Carbohydrate-Related Middle Molecules Based on Sequential Site-Selective Functionalization Grant-in-Aid for Scientific Research on Innovative Areas 1 April 2016-31 March 2018

Yoshida, K.

Total Synthesis of Natural Products by Using Organocatalytic Asymmetric Construction of All-carbon Quaternary Stereocenter Grant-in-Aid for Young Scientists (B) 1 April 2016-31 March 2019

— Advanced Inorganic Synthesis —

Teranishi, T.

Establishment of Deeply Penetrating Photoacoustic Imaging Technology Based on Fucntional Probes: Design and Synthesis of Activatable Probes and Development of in vivo Imaging Technology Industry-Academia Collaborative R&D Programs, Japan Agency for Medical Research and Development 1 December 2011-31 March 2017

Teranishi, T.

Synthesis of Magnetic Nanoparticles for Creating Novel Nanocomposite Magnetic Materials Elements Strategy Initiative, MEXT 1 July 2012-31 March 2022

Teranishi, T.

Research on Nanoscale Phase-Controlled Nanocomposite Magnets Mirai Kaitaku Research Project, NEDO 1 October 2012-31 March 2022

Teranishi, T.

Development of Green Sustainable Chemical Process Mirai Kaitaku Research Project, NEDO 1 November 2012-31 March 2022

Teranishi, T.

Novel Development of Asymmetry Chemistry in Inorganic Nanocrystals Grant-in-Aid for Scientific Research on Innovative Areas

30 June 2016-31 March 2021

Teranishi, T.

Formation of Novel Metallic Phase Nanoparticles and Development of Their Catalytic Properties Grant-in-Aid for Scientific Research (B)

1 April 2016-31 March 2019

Sakamoto, M.

Investigation on the Gain-of-function Process by the Formation of Nanoparticle Assemblage Using DNA Origami

SPIRITS (Supporting Program for Interaction-based Initiative Team Studies)

1 April 2015-31March 2017

Sato, R.

Development of The Novel and Versatile Alloying Process via Nanosized Phosphorus Compounds

Grant-in-Aid for Challenging Exploratory Research

1 April 2015-31March 2017

Saruyama, M.

Orientational Control and Structure-specific Properties of Heterostructural Nanoparticles

Grant-in-Aid for Research Activity Start-up

1 October 2015-31March 2017

DIVISION OF MATERIALS CHEMISTRY — Chemistry of Polymer Materials —

Tsujii, Y.

Reinforcement of Resiliency of Concentrated Polymer Brushes and Its Tribological Applications-Development of Novel "Soft and Resilient Tribology (SRT)" System

ACCEL Program, JST

1 September 2015-31 March 2019

Tsujii, Y.

Super Lubrication of Novel Nano-Brushes

Advanced Environmental Materials of Green Network of Excellence (GRENE) program, MEXT

6 December 2011-31 March 2016

Tsujii, Y.

Development of High-Performance Li-ion Batteries using High-capacity, Low-cost Oxide Electrodes

NEDO Project for Development of Novel Technology in Li-ion Batteries

1 October 2012-27 February 2017

Ohno, K.

Development of Ionic Liquid-Containing Blend Films

PRESTO (Precursory Research for Embryonic Science and Technology), JST

1 October 2013-31 March 2017

Sakakibara, K.

Construction of Cellulosic Bottlebrushes with Regioselectively Substituted Side Chains

Grant-in-Aid for Young Scientist (B)

1 April 2016-31 March 2018

— Polymer Controlled Synthesis —

Yamago, S

New Organic Chemistry and Material Science of Curved π -Conjugated Molecules

Grant-in-Aid for Scientific Research (S)

1 April 2016-31 March 2021

Yamago, S.

Highly Value-added Polymer Material Created by New Living Radical Polymerization Agent

Next Generation Technology Transfer Program (NexTEP), JST 1 April 2014–31 March 2020

- Inorganic Photonics Materials -

Mizuochi, N.

High Sensitive and High Resolution Quantum Nano-sensor by Diamond

Grant-in-Aid for Scientific Research (A)

1 April 2016-31 March 2021

Mizuochi, N.

Innovative Magnetic Sensor Based on Nano-electronics of Carbon Materials

Core Research for Evolutional Science and Technology (CREST), JST

1 April 2014-31 March 2019

Mizuochi, N.

Research on Diamond Quantum Information Device TORAY Science and Technology Research Grant 1 January 2016–31 March 2018

Morishita, H.

Electrical Coherent Detection of Electron Spin of NV Centers in Diamond

Grant-in-Aid for Encouragement of Young Scientists (B)

1 April 2016-31 March 2018

- Nanospintronics -

Ono, T.

Spin-orbitronics and Device Application

New Research Projects under Specially Promoted Research

1 April 2015-31 March 2020

DIVISION OF BIOCHEMISTRY — Biofunctional Design-Chemistry —

Futaki, S.

New Strategies for Intracellular Delivery of Biopharmaceuticals Grant-in-Aid for Science Research (A)

1 April 2015-31 March 2018

Imanishi, M.

Construction of Strategies for Sequence Specific Epigenomic Manipulation

Grant-in-Aid for Science Research (B)

1 April 2016-31 March 2019

— Chemistry of Molecular Biocatalysts —

Watanabe, B.

Synthesis of γ -Glutamyl Transpeptidase-Specific Chemical Probes and Their Application to Cancer Immunotherapy

Grant-in-Aid for Young Scientists (B)

1 April 2015-31 March 2017

- Molecular Biology -

Aoyama, T.

Plant Epidermal Cell Differentiation Regulated by the Transcription

Grant-in-Aid for Scientific Research (B)

1 April 2016-31 March 2020

Aoyama, T.

Mechanism for Establishment of Planar Polarity in Plant Cell Morphogenesis

Grant-in-Aid for Scientific Research (C), Special Field

1 April 2016–31 March 2019

- Chemical Biology -

Uesugi, M.

Control and Analysis of Cells by Synthetic Small Molecules Grant-in-Aid for Scientific Research (S) 30 May 2014-31 March 2019

Chemical Biological Exploration of New Functions of Endogenous Lipid-related Molecules

AMED-CREST

1 October 2014-31 March 2020

DIVISION OF ENVIRONMENTAL CHEMISTRY Molecular Materials Chemistry —

Structure and Function of Organic Thin-Film Solar Cells: Specially-Shaped Polymers and Hierarchical Structure Analysis Grant-in-Aid for Scientific Research (A)

1 April 2013-31 March 2016

Kaii. H.

Adachi Molecular Exciton Engineering Project ERATO (Exploratory Research for Advanced Technology), JST 1 April 2014-31 March 2018

Fukushima, T.

Solid-State NMR Analysis of Bulk Heterostructures toward Highefficiency Organic Solar Cells

Grant-in-Aid for Young Scientists (B) 1 April 2014-31 March 2016

Fukushima, T.

Study on Weather Resistance of Solution-Processable Organic Solar

Suga Weathering Technology Foundation

1 April 2015-31 March 2016

Shizu, K.

A Method of Visualizing Radiative and Non-Radiative Decays and Its Applications to Design for Deep-Blue Organic Emitters Grant-in-Aid for Young Scientists (B)

1 April 2015-31 March 2017

Analysis of Organic Photovoltaics Materials by Solid-State Dynamic Nuclear Polarization

SPIRITS (Supporting Program for Interaction-based Initiative Team Studies)

1 April 2015-31 March 2017

Fukushima, T.

A Fundamental Study on Weather Resistance of Solution-Processed Organic Solar Cells by Solid-State NMR Grant-in-Aid for Scientific Research (C)

1 April 2016-31 March 2018

— Hydrospheric Environment Analytical Chemistry —

Sohrin, Y.

Development of Novel Proxies for Paleoceanography on the Precise Analysis of Stable Isotope Ratios of Heavy Metals Grant-in-Aid for Challenging Exploratory Research

1 April 2014-31 March 2017

Sohrin, Y.

Development of Heavy Metal Stable Isotope Marine Chemistry to Understand Marine Environment and Ecosystems Gtant-in-Aid for Scientific Research (A)

1 April 2015-31 March 2018

— Solution and Interface Chemistry —

Hasegawa, T.

Development of ROA Imaging and its Application to Visualization of Atropisomers for a Study of Fluoroorganic Chemistry Grant-in-Aid for Scientific Research (A)

1 April 2015-31 March 2020

Hasegawa, T.

Development of Novel Analytical Techniques for Revealing Molecular Orientation of Adsorbed Molecules on a Rough Surface or on Nano Particles

Grant-in-Aid for challenging Exploratory Research 1 April 2014-31 March 2016

Hasegawa, T.

Development of a Novel Analytical Spectroscopy for Strategic Molecular Design of a Fluorine-containing Acryl Polymer Enabling Us to Overcome Environmental Regulations

Matching Planner Program: Cooperation Research with a Company 1 June 2016-31 March 2017

Shimoaka, T.

Development of Analytical Techniques for Studying the Structure and Property of a Polymer Influenced by Minute Water Molecules Involved in a Polymer Thin Film

Grant-in-Aid for Young Scientists (B)

1 April 2014-31 March 2017

- Molecular Microbial Science -

Kurihara, T.

Mechanism of Biogenesis of Membrane Microdomain Containing Polyunsaturated Fatty Acids in Bacteria and Its Physiological

Grant-in-Aid for Scientific Research (B)

1 April 2015-31 March 2018

Kurihara, T.

Analysis of Cold-Adaptation Mechanism of Food Spoilage Bacteria and Its Application to Food Industry

Grant-in-Aid for Challenging Exploratory Research

1 April 2015-31 March 2017

Kawamoto, J.

Development of a Membrane Protein Production System by Using Tailor-made Membrane Vesicles Synthesized by Extremophiles Grant-in-Aid for Challenging Exploratory Research 1 April 2016–31 March 2018

Ogawa, T.

Research on Acyltransferases that Create Heterogeneity of Bacterial Biomembranes

Grant-in-Aid for Research Activity Start-up

23 August 2015-31 March 2017

DIVISION OF MULTIDISCIPLINARY CHEMISTRY — Polymer Materials Science —

Takenaka, M.

Photon and Quantum Basic Research Coordinated Development Program, JST

1 September 2013-31 March 2018

Takenaka, M.

Nano-Control Technologies for DSA Nano-Patterning Nano Defect Management Project 1 July 2016–31 March 2018

Ogawa, H.

Development of Visualizing Method Through Cooperative Small Angle X-ray Scattering Coupled with Computed Tomography (SAXS-CT) and Information Science

Strategic Basic Research Programs, PRESTO (Precursory Research for Embryonic Science and Technology), JST

1 September 2016-31 March 2020

- Molecular Rheology -

Watanabe, H.

Relationship Between Chemical Structure and Extensional Behavior of Entangled Polymer Chain

Grant-in-Aid for Scientific Research (B)

1 April 2015-31 March 2018

Matsumiya, Y.

Experimental Test on the Dynamics of Telechelic Polymers Grant-in-Aid for Scientific Research (C)

1 April 2015-31 March 2018

ADVANCED RESEARCH CENTER FOR BEAM SCIENCE — Particle Beam Science —

Iwashita, Y.

Quantum Improvement of the Superconducting Acceleration Cavity Performance by the Laminated Film Structure

Grant-in-Aid for Challenging Exploratory Research

1 April 2014-31 March 2016

Iwashita, Y.

Fundamental Technology Development for High Brightness X-ray Source and the Imaging by Compact Accelerator

Photon and Quantum Basic Research Coordinated Development Program

1 April 2013-31 March 2017

- Laser Matter Interaction Science -

Sakabe, S

Proof of Concept for Electron Optics Using Intense Laser-driven Surface Wave

Grant-in-Aid for Scientific Research (A)

1 April 2016-31 March 2019

Hashida, M.

Stable Formation of Advanced Functionality on Metal Surface Produced by High Electric Field of Laser Pulse Grant-in-Aid for Scientific Research (C)

1 April 2016-31 March 2019

Hashida, M.

Advanced Research Program for Energy and Environmental Technologies/Manufacturing Technologies Development of High Quality Laser Material Processing for Inducing New Functionalities New Energy and Industrial Technology Development Organization 4 January 2016–4 January 2017

Inoue S

Demonstration of Laser-driven Ultrafast and Intense Electron Source with Solid-plasma Hybrid Cathode

Grant-in-Aid for Encouragement of Young Scientists (B)

1 April 2016-31 March 2018

- Electron Microscopy and Crystal Chemistry -

Kurata, H.

Advanced Characterization Nanotechnology Platform at Kyoto University

Nanotechnology Platform Project, MEXT

2 July 2012-31 March 2022

Kurata, H.

State Analysis of Organic Nanomaterials by High-Resolution EELS Grant-in-Aid for Challenging Exploratory Research

1 April 2016-31 March 2018

Haruta, M.

Electronic State Mapping Using Oxygen Grant-in-Aid for Young Scientist (A) 1 April 2014–31 March 2018

Haruta, M.

Basic Research of Atomic Resolution Organic Crystal Image Using STEM

Grant-in-Aid for Challenging Exploratory Research

1 April 2014–31 March 2017

INTERNATIONAL RESEARCH CENTER FOR ELEMENTS SCIENCE

-Synthetic Organotransformation-

Nakamura, M.

Synthesis of Nitrogen-Containing Polycyclic Aromatic Compounds via Iron-catalyzed C-H Amination

Grant-in-Aid for Challenging Exploratory Research

1 April 2015-31 March 2016

Takaya, H.

Solution-Phase Characterization of Paramagnetic Metal Complex by X-ray Absorption Spectroscopy

Grant-in-Aid for Challenging Exploratory Research

1 April 2015-31 March 2017

Isozaki, K.

Development of Catalytic Multi-photon-excited Photoreactions in the Reaction Field Localizing Substrates and Excitation Sources Grant-in-Aid for Scientific Research on Innovative Areas "Application of Cooperative Excitation into Innovative Molecular Systems with High-Order Photofunctions"

1 April 2015-31 March 2017

Iwamoto, T.

Development of Aromatic C-H Functionalization Base on Cation- π Interaction

Grant-in-Aid for Young Scientists (B)

1 April 2015-31 March 2017

- Advanced Solid State Chemistry -

Shimakawa, Y.

Solid-state Chemistry for Transition-metal Oxides: Exploring for New Materials with Novel Functionalities

JSPS Core-to-Core Program

1 April 2016-31 March 2020

— Organometallic Chemistry —

Ozawa, F.

Synthesis and Catalytic Properties of Stimulus-responsible Transition Metal Complexes Bearing Low-coordinate Phosphorus Ligands Grant-in-Aid for Scientific Research on Innovative Areas "Stimuliresponsive Chemical Species for the Creation of Functional Molecules'

1 April 2012-31 March 2017

Synthesis and Catalytic Applications of Non-innocent Phosphaalkene

Grant-in-Aid for Scientific Research (B)

1 April 2014-31 March 2017

Wakioka, M.

Development of Highly Efficient Catalytic Systems for Direct Arylation Polymerization based on Equilibrium between Active and **Dormant Species**

Grant-in-Aid for Young Scientists (B)

1 April 2015-31 March 2017

Ozawa, F.

Development of Highly Efficient Catalysts for Synthesizing of π -Conjugated Polymers via Direct Arylation

ACT-C, JST

1 October 2012-31 March 2018

- Nanophotonics -

Kanemitsu, Y.

Evaluation of Nonradiative Carrier Recombination Loss in Concentrator Heterostructure Solar Cells

CREST(Core Research for Evolutional Science and Technology),

1 October 2011-31 March 2017

BIOINFORMATICS CENTER

— Chemical Life Science —

Ogata, H.

Neo-virology, the Raison D'être of Viruses - Deciphering the Mechanisms of Virus-host Co-existence in Aquatic Environments Grant-in-Aid for Scientific Research on Innovative Areas 30 June 2016-31 March 2021

Ogata, H.

Probabilistic and Statistical Theory on Non-Abelian Topological Semigroup A* and Its Application to Environmental Microbiology and Bioengineering

Grant-in-Aid for Scientific Research (B)

19 July 2016-31 March 2019

Ogata, H.

A Holistic Ecosystemic Investigation on Marine Giruses, Virophages and Their Eukaryotic Hosts

Grant-in-Aid for Scientific Research (C)

1 April 2014-31 March 2017

Ogata, H.

Are Viruses Elementary Particles that Generate and Maintain the Diversity of Marine Organisms?

Pursuit of Ideal, CANON Foundation

1 April 2014-31 March 2017

Development of Integrated Proteome Database jPOST Database Integration Coordination Program, JST 1 April 2015-31 March 2018

Goto, S.

Bioinformatics for Marine Microbial Genomes and Environmental Data

CREST (Core Research for Evolutional Science and Technology),

1 October 2012-31 March 2017

Goto, S.

Elucidation on Evolutionary Mechanisms of Antigenic Variation Gene Families

Grant-in-Aid for Scientific Research (B)

1 April 2014-31 March 2018

— Bio-knowledge Engineering —

Mamitsuka, H.

Reinforcement of Resiliency of Concentrated Polymer Brushes and Its Tribological Applications

Strategic Basic Research Program, ACCEL, JST

1 September 2015-31 March 2020

Mamitsuka, H.

Efficiently Inferring Factors Embedded in Multiple Data Matrices Grant-in-Aid for Scientific Research (B) 1 April 2016-31 March 2019

Yotsukura, S.

Feasibility Study of Allergen Genes Identification and Prediction of the Gene Expression Level Associated with Cultivation Conditions Specific Project Investigation, JST 1 October 2016-31 March 2017

Yamada, M. Nonlinear Feature Selection for Ultra-High Dimensional Data Grant-in-Aid for Young Scientists (B) 1 April 2016–31 March 2018

Yamada, M. Nonlinear Feature Selection for Science Discovery
PRESTO (Precursory Research for Embryonic Science and Technology), JST 1 December 2016–31 March 2020

THESES

BABA, Tomonori

D Pharm Sc, Kyoto University

"Development of Aniline-type Acid Base Catalysts for Asymmetric

Direct Aldol Reactions"

Supervisor: Prof KAWABATA, Takeo

23 March 2016

CHEN, Lihui

D Sc, Kyoto University

"Synthesis and Plasmonic Properties of Copper-based Nanocrystals"

Supervisor: Prof TERANISHI, Toshiharu

23 March 2016

HSU, Shu-Yao

D Eng, Kyoto University

"Fabrication of Well-Defined Architectures of Ultrahigh-Molecular-

Weight Polymers by Living Radical Polymerization"

Supervisor: Prof TSUJII, Yoshinobu

23 March 2016

HYAKUTAKE, Ryuichi

D Pharm Sc, Kyoto University

"Selective Reactions in Conjugate Addition to Unsaturated Carbonyl

Compounds"

Supervisor: Prof KAWABATA, Takeo

23 March 2016

IMAYOSHI, Ayumi

D Pharm Sc, Kyoto University

"Discrimination of Mobile Supramolecular Chirality: Acylative

Molecular Transformation by Organocatalysis'

Supervisor: Prof KAWABATA, Takeo

23 March 2016

JIRA, Jindalertudomdee

D Inf, Kyoto University

"Chemical Compound Enumeration and Host-Pathogen Protein

Interaction Prediction By Graph-Based Approaches"

Supervisor: Prof AKUTSU, Tatsuya

23 September 2016

KAWAGUCHI, Masashi

D Sc, Kyoto University

"Current Induced Effective Magnetic Field in Ferromagnetic

Multilayers"

Supervisor: Prof ONO, Teruo

12 January 2016

KAWAGUCHI, Yoshimasa

D Pharm Sc, Kyoto University

"Identification of an Octaarginine Receptor for Its Intracellular

Uptake Using Photocrosslinking and Elucidation of the Intracellular

Uptake Pathway"

Supervisor: Prof FUTAKI, Shiroh

23 March 2016

MOHAMED, Ahmed Mohamed

D Pharm Sc, Kyoto University

"Development of Computational Analysis Tools for Natural Products

Research and Metabolomics"

Supervisor: Prof MAMITSUKA, Hiroshi

23 March 2016

NAGATA, Koichi

D Sc, Kyoto University

"Synthetic Studies on the Al=Al Doubly Bonded Species and

Elucidation of Their Reactivity"

Supervisor: Prof TOKITOH, Norihiro

23 March 2016

NISHIMURA, Hidetaka

D Eng, Kyoto University

"Development of Functional Organic Materials Using Oxygen-

Bridged Triarylamine Skeleton as Quasiplanar Structure"

Supervisor: Prof MURATA, Yasujiro

23 March 2016

OKU, Akihiko

D Pharm Sc, Kyoto University

"Peptide-Scientific Approaches for Intracellular Delivery of

Biological Macromolecules"

Supervisor: Prof FUTAKI, Shiroh

23 March 2016

SEKI, Hayato

D Sc, Kyoto Uiversity

"High Pressure Synthesis of Heavily Substituted Perovskite Oxides

and Explorations of their Physical Properties"

Supervisor: Prof SHIMAKAWA, Yuichi

23 March 2016

TAKAYA, Junichiro

D Med Sc, Kyoto University

"A Potent and Site-selective Agonist of TRPA1"

Supervisor: Prof UESUGI, Motonari

23 March 2016

TAKEUCHI, Hironori

D Pharm Sc, Kyoto University

"Novel Synthetic Strategy for Natural Glycosides based on Catalytic

Sequential Functionalization of Unprotected Glucose"

Supervisor: Prof KAWABATA, Takeo

23 March 2016

YANAGI, Masanori

D Pharm Sc, Kyoto University

"Novel Strategy for Acceleration of Nucleophilic Catalysis via Counter-anion Control and Catalytic Site-selective Functionalization

of Polyols'

Supervisor: Prof KAWABATA, Takeo

23 March 2016

110 THESES

YOSHIMURA, Yoko D Sc, Kyoto University "Spin Dynamics in Ferromagnetic Nanowire" Supervisor: Prof ONO, Teruo 12 January 2016

YOTSUKURA, Sohiya D Pharm Sc, Kyoto University "Computational Framework for the Dissection of Cancer Genomic Architecture and Its Association in Different Biomarkers" Supervisor: Prof MAMITSUKA, Hiroshi 23 September 2016

THE 116TH ICR ANNUAL SYMPOSIUM

Seminars Meetings and Symposia

THE 116TH ICR ANNUAL SYMPOSIUM

(2 December 2016)

ORAL PRESENTATIONS

TAKAYA, Hikaru (Synthetic Organotransformation)

"Solution-Phase Structure Analysis of X-ray Absorption Spectroscopy"

MIZUHATA, Yoshiyuki (Organoelement Chemistry)

"The Chemistry of 'Heavy Phenyl Anions'

WAKIOKA, Masayuki (Organometallic Chemistry)

"Direct Arylation Polymerization: A New Synthetic Method of π-Conjugated Polymers"

SAITO, Takashi (Advanced Solid State Chemistry)

"High-pressure Syntheses of Transition Metal Oxides Containing Unusually High Valence Cations"

FURUTA, Takumi (Synthetic Organic Chemistry)

"Direct Asymmetric Cross-aldol Reaction of Aliphatic Carbonyl Compounds via Catalytic Discrimination of Carbonyl Groups"

ICR Award for Young Scientists —

NAKAMURA, Yasuyuki (Polymer Controlled Synthesis)

"Termination Mechanism in the Radical Polymerization of Methyl Methacrylate and Styrene Determined by the Reaction of Structurally Well-Defined Polymer End Radicals"

— ICR Award for Young Scientists —

HARUTA, Mitsutaka (Electron Microscopy and Crystal Chemistry) "Determination of Elemental Ratio in an Atomic Column by Electron Energy Loss Spectroscopy"

— ICR Award for Graduate Students —

SUZUKI, Yuko (Organoelement Chemistry)

"Isolation and Ambident Reactivity of a Chlorogermylenoid"

— ICR Award for Graduate Students —

ZHANG, Rui (Structural Organic Chemistry)

"Synthesis of a Distinct Water Dimer inside Fullerene C₇₀"

— ICR Award for Graduate Students —

YAMADA, Takumi (Nanophotonics)

"Dynamic Optical Properties of CH₃NH₃PbI₃ Single Crystals As Reveald by One- and Two-Photon Excited Photoluminescence Measurements"

ICR Grants for Promoting Integrated Research —

IWAMOTO, Takahiro (Synthetic Organotransformation)

"Mechanistic Study of Iron Fluoride-Catalyzed Cross-Coupling Reaction"

KATO, Mariko (Molecular Biology)

"Development of Cell-permeable Probes for Phosphatidylinositol 4,5-bisphosphate in Arabidopsis"

SAKAMOTO, Masanori (Advanced Inorganic Synthesis)

"Investigation on Photoinduced Carrier Dynamics at Nano-hetero Interfaces Using Multiple Laser Pulses"

POSTER PRESENTATIONS

LW: Laboratory Whole Presentation

LT : Laboratory Topic

GE: General Presentation

— Organoelement Chemistry —

W "Studies on the Synthesis and Properties of Novel Organic Compounds Containing Heavier Elements"

— Structural Organic Chemistry —

W "Research Activities in Structural Organic Chemistry Laboratory"

GE OZAKI, Masashi; SHIMAZAKI, Ai; JUNG, Mina; NAKAIKE, Yumi; MARUYAMA, Naoki; AHAREN, Tomoko; RAFIEH, Alwani; SASAMORI, Takahiro; TOKITOH, Norihiro; WAKAMIYA, Atsushi; MURATA, Yasujiro

"Fabrication Protocol of Highly Efficient Perovskite Solar Cells Using Purified Precursor Materials"

GE OKAZAKI, Shuhei; MURATA, Michihisa; WAKAMIYA, Atsushi; MURATA, Yasujiro

"Synthesis of Polymers Containing Imidazole-Fused Benzothiadiazole

— Synthetic Organic Chemistry —

"Current Topics of Fine Organic Synthesis"

GE ARAI, Kenta; UEDA, Yoshihiro; FURUTA, Takumi; KAWABATA,

"Dirhodium-Catalyzed Chemo- and Regioselective C(sp2)-H Amination"

— Advanced Inorganic Synthesis —

"Current Research Activities in Advanced Inorganic Synthesis"

GE CHEN, Lihui; SAKAMOTO, Masanori; TERANISHI, Toshiharu "Tin Ion Directed Morphology Evolution of Copper Sulfide Nanoparticles"

GE KIMURA, Masato; SAKAMOTO, Masanori; FURUBE, Akihiro; ADACHI, Hiroyuki; SUGIMOTO, Toshiki; WATANABE, Kazuya; MATSUMOTO, Yoshiyasu; TERANISHI, Toshiharu

"Visible Light-induced Charge Separation in Heterostructured Au_xS/ ZnS Nanoparticles"

©E MATSUMOTO, Kenshi; SATO, Ryota; TRINH, T. Thuy; SAKUMA, Noritsugu; TERANISHI, Toshiharu

"High-Performance L1₀-FePd/α-Fe Nanocomposite Magnets with Well-Controlled Nanostructures"

— Chemistry of Polymer Materials —

- W "Research Activities in Chemistry of Polymer Materials"
- © KINOSE, Yuji; SAKAKIBARA, Keita; TSUJII, Yoshinobu "Conformation of Well-defined Janus-type Bottle Brush with Cellulosic Main Chain in Solution"
- GE SHIMIZU, Yoshihiko; SAKAKIBARA, Keita; TSUJII, Yoshinobu "Preparation of Bacterial Cellulose-reinforced Elastomers Using Percolation Structure"

— Polymer Controlled Synthesis —

- W "Research Activities in Polymer Controlled Synthesis"
- © SEKI, Toyomitsu; YU, Mengmeng; NAKAMURA, Yasuyuki; YAMAGO, Shigeru
- "New Fluorous Organotellurium Compound for Living Radical Polymerization and Efficient Recovery and Recycle of the Organotellurium Group"
- E HUASHEN, Zhiren; NAKAMURA, Yasuyuki; YAMAGO,
- "Synthesis of Organoheteroatom Chain Transfer Agent via Radical Trapping Reaction and Their Uses for Living Radical Polymerization"

— Inorganic Photonics Materials —

W "Introduction of Research Topic of Inorganic Photonics Materials Group"

— Nanospintronics —

- IT TANAKA, Kensho; MORIYAMA, Takahiro; KAMIYA, Michinari; MIZUNO, Hayato; ONO, Teruo
- "Spin Hall Effect in Ferromagnets Measured by Spin-torque FMR"
- GE TANIGUCHI, Takuya; KIM, Kab-Jin; KOYAMA, Tomohiro; CHIBA, Daichi; ONO, Teruo
- "Observation of Discrete Jump of Domain Wall Segment among Quenched Disorders"

— Biofunctional Design-Chemistry —

- W "Research Activities in Biofunctional Design-Chemistry"
- GE SAKAMOTO, Kentarou; AKISHIBA, Misao; FUTAKI, Shiroh "A Statistical Research on Structure-activity Relationships between Antimicrobial Peptides Derived from a Spider Toxin"
- GE SAKAGAMI, Koki; FUTAKI, Shiroh
- "Mechanism of Cellular Uptake of Cell-penetrating Peptides with Stapled α-Helix"

— Chemistry of Molecular Biocatalysts —

- W "Research Activities in Laboratory of Chemistry of Molecular Biocatalysts"
- Molecular Biology —
- "Research Activity in Laboratory of Molecular Biology"

- Chemical Biology -

- "Create New World of Bioactive Synthetic Small Molecules"
- Molecular Materials Chemistry -
- "Molecular Materials Chemistry"
- © MIURA, Tsutomu; FUKUSHIMA, Tatsuya; KAJI, Hironori "High Performance Bulk Heterojunction Organic Thin Film Solar Cells with Halogen Free Solutions"
- GE OGATA, Isamu; KUBO, Shosei; SUZUKI, Katsuaki; KAJI, Hironori
- "Synthesis and Photophysical Properties of a Deuterated Thermally Activated Delayed Fluorescence Material"

- Hydrospheric Environment Analytical Chemistry -

- "Studies in Laboratory of Hydrospheric Environment Analytical Chemistry"
- © OKITA, Shutaro; UMETANI, Shigeo; SOHRIN, Yoshiki "Solid Phase Extraction and Separation of 13 Group Metal Ions with Acylpyrazolones Loaded XAD7 Resin"

- Solution and Interface Chemistry -

- © IZUMI, Naoki; SHIMOAKA, Takafumi; HASEGAWA, Takeshi "Structural Analysis of Polymer Thin Films of Fluoroacrylate Having an Rf Side Chain with Various Lengths Using IR Spectroscopy"
- © UKAI, Hironori; SHIMOAKA, Takafumi; HASEGAWA, Takeshi "Study on an Aggregation Factor of Amphiphiles Involving a Perfluoroalkyl Group Using IR Spectroscopy"

- Molecular Microbial Science -

- W "Research of Molecular Microbial Science Laboratory"
- © TOYOTAKE, Yosuke; CHO, Hyun-Nam; OGAWA, Takuya; KAWAMOTO, Jun; KURIHARA, Tatsuo
- "Characterization of a Novel Enzyme for Membrane Phospholipid Biosynthesis from a Cold-adapted Bacterium"

— Polymer Materials Science —

- W "Introduction of Polymer Materials Science Laboratory"
- GE NAGANO, Yoshiki; TAKENAKA, Mikihito; JANG, Junhyeok "The Study of Heterogeneous Structure of Acrylonitrile-butadiene Random Copolymer (NBR)"
- (IE) HIGASHINO, Naruhisa; TAKENAKA, Mikihito; YAMAGUCHI, Kaoru; Aisyah
- "Alignment of Ferritin Protein by Using Nano Pattern of Block Copolymers"
- © MIYATAKE, Yuki; OGAWA, Hiroki; NISHIDA, Koji; YAMADA, Norifumi; KANAYA, Toshiji
- "Phase Separation and Dewetting of Deuterated Polystyrene/ poly-2-chlorostyrene Blend Thin Films"

- Molecular Rheology -

"Research Activities in Laboratory of Molecular Rheology"

— Molecular Aggregation Analysis —

LT MURDEY, Richard

"Introduction to Nonlinear Photocurrent Spectroscopy"

— Particle Beam Science —

W "Status of Accelerator Laboratory, Advanced Research Center for Beam Science"

© MIYAWAKI, Eisuke; FUWA, Yasuhiro; TONGU, Hiromu; IWASHITA, Yoshihisa

"Study of Ion Sources for Future Accelerators"

GE FUWA, Yasuhiro; IWASHITA, Yoshihisa

"Beam Focusing System for Klystrons with Permanent Mangets"

© KITAHARA, Ryunosuke; IWASHITA, Yoshihisa; NOP Group "Measurement of 14N(n, p)14C Cross Section Using Cold Neutron Beam at J-PARC'

— Laser Matter Interaction Science —

W "Research Activities in Laboratory of Laser Matter Interaction Science"

© FURUKAWA, Yuki; TERAMOTO, Kensuke; MORI, Kazuaki; NAKAMIYA, Yoshihide; INOUE, Shunsuke; HASHIDA, Masaki; SAKABE, Shuji

"Creation of Nanostructures on Metal Surface by Double-pulse Femtosecond Laser Irradiation"

(III) MORI, Kazuaki; HASHIDA, Masaki; NAGASHIMA, Takeshi; LI, Dazhi; TERAMOTO, Kensuke; INOUE, Shunsuke; SAKABE, Shuii

"THz Waves Generation from Interaction between a Target of Argon Clusters and Intense Double-pulse Laser"

© TERAMOTO, Kensuke; INOUE, Shunsuke; TOKITA, Shigeki; YASUHARA, Ryo; TERAO, Tokinori; KAWANAKA, Junji; NAGASHIMA, Takeshi; HASHIDA, Masaki; SAKABE, Shuji "Generation of Intense THz Surface Waves on a Metal Wire by Intense Laser"

- Electron Microscopy and Crystal Chemistry -

W "Research Activities in Division of Electron Microscopy and Crystal Chemistry"

GE YAMAGUCHI, Atsushi; HARUTA, Mitsutaka; NEMOTO, Takashi; KURATA, Hiroki

"Detecting the Directionality of Local Electronic States in SrTiO3 by Momentum-selected STEM-EELS"

- Structural Molecular Biology -

"Introduction of Structural Molecular Biology Laboratory"

— Synthetic Organotransformation —

"Introduction of Synthetic Organotransformation Laboratory"

GE OKUZONO, Chiemi; IWAMOTO, Takahiro; ADAK, Laksmikanta; JIN, Masayoshi; TAKAYA, Hikaru; NAKAMURA, Masaharu "Iron-Catalyzed Enantioselective Cross-Coupling of Haloester with Aryl Boron Reagent"

GE UENO, Ryo; ISHIBASHI, Kosuke; ISOZAKI, Katsuhiro; TAKAYA, Hikaru; NAKAMURA, Masaharu

"Synthesis and Photocatalytic Property of Peptide Dendron Thiolate-Functionalized Gold Clusters"

- Advanced Solid State Chemistry -

W "Introduction of Advanced Solid State Chemistry Laboratory"

© YUU, Hou; SEKI, Hayato; KAKU, Kaisen; HOSAKA, Yoshiteru; SAITO, Takashi; SHIMAKAWA, Yuichi

"Geometrical Spin Frustration of Unusually High Valence Fe⁵⁺ in the Double Perovskite La₂LiFeO₆"

- Organometallic Chemistry -

"Activity Report: Organometallic Chemistry Laboratory"

- Nanophotonics -

W "Recent Research Topics of Nanophotonics Group"

- Chemical Life Science -

"Research Activities in Laboratory of Chemical Life Science"

GE LI, Yanze; WATAI, Hiroyasu; HINGAMP, Pascal; YOSHIDA, Takashi; OGATA, Hiroyuki

"Amplicon Analysis of Megaviridae DNA Polymerase Gene"

- Mathematical Bioinformatics -

LT JIRA, Jindalertudomdee

"Enumeration Method for Structural Isomers Containing User-Defined Structures Based on Breadth-First Search Approach'

"LBSizeCleav: Improved Support Vector Machine (SVM)-based Prediction of Dicer Cleavage Sites Using Loop/Bulge Length"

- Bio-knowledge Engineering -

GE YAMADA, Makoto; WIMALAWARNE, Kishan "Nonlinear Feature Selection for Biological Data"

- ICR Grants for Promoting Integrated Research -

MURATA, Michihisa; TSUKAO, Masahiro; WAKAMIYA, Atsushi; MURATA, Yasujiro; SUZUKI, Katsuaki; KAJI, Hironori "Elucidation of Solid-state Structures and Functions of Metal Complexes with 3D π -Systems"

KAYAHARA, Eiichi; FUKUSHIMA, Tatsuya

"Application of Cyclic π -Conjugated Moleculars to Oraganic Electronics"

CHO, Hyun-Nam; KAWAMOTO, Jun

"Analysis of the Fatty Acid-Recognition Mechanism of an Anti-Fatty Acid Antibodies and Their Application"

NGUYEN, Hao Canh "Advanced Computational Tools for Knowledge Discovery in Ecology and Biology"

SEMINARS

Prof AJAYAGHOSH, Ayyappanpillai

CSIR-NIST, India

"Self-Sorted Donor-Acceptor Assemblie"

19 October 2016

Prof AKIYAMA, Yutaka

Tokyo Institute of Technology, Tokyo, Japan

"Massively Parallel Bioinformatics Computing on K-computer, TSUBAME, and Azure: Metagenome Analysis and Exhaustive

Protein-Protein Interaction Prediction"

1 July 2016

Assoc Prof AU, Kin Fai

University of Iowa, U.S.A.

"Transcriptome Analysis at the Gene Isoform Level Using Hybrid

Sequencing'

11 October 2016

Prof BERKESSEL, Albrecht

Department of Chemistry, University of Cologne, Cologne, Germany

"Carbene Catalysis and the Breslow Intermediate"

1 March 2016

Dr BURLINA, Fabienne

Laboratoire des Biomolécules, UMR 7203, Université P. et M. Curie

- CNRS - ENS, Paris, France

"Design and Study of Cyclic Cell-Penetrating Peptides"

25 October 2016

Prof CHEN, Hongyu

Nanyang Technological University (Nanjin Tech University),

Singapore (China, P.R.)

"Exploiting Polymer in Nanosynthesis"

19 December 2016

Dr CHEN, Sophia

Laboratoire D'utilisation des Lasers Intenses, Ecole Polytrchnique,

"What Would You Do with 1022 W/cm2 Laser Intensities?"

1 July 2016

Mr. CHENG, Didier

Laboratoire des Biomolécules, UMR 7203, Université P. et M. Curie

- CNRS - ENS, Paris, France

"Comparative Studies between Antimicrobial Peptides and Cell-

Penetrating Peptides"

25 October 2016

Prof CLAYDEN, Jonathan

School of Chemistry, University of Bristol, Bristol, U.K.

"Unconventional Reactivity from Organolithium Compounds"

25 April 2016

Dr CONWAY, Timothy

Institute of Geochemistry and Peterology, Department of Earth

Sciences, ETH Zürich, Switzerland

"Investigating the Role of Dust in the Marine Iron Cycle Using Iron

Isotopes'

8 July 2016

Prof DE LA HIGUERA, Colin Manuel

University of Nantes, Nantes, France

"Machine Learning is about Intelligent Forgetting"

14 April 2016

Senior Researcher ENAMI, Shinichi

National Institute for Environmental Studies, Ibaraki, Japan

"Physical Chemistry on an Interfacial Phenomena: (1) Mechanism of

a Chemical Reaction at an Air/Liquid Interface"

19 December 2016

Dr ENDO, Hisashi

Hokkaido University, Hokkaido, Japan

"Contrasting Biogeography and Diversity Patterns between Diatoms

and Haptophytes in the Central Pacific Ocean"

22 December 2016

Dr FUCHS, Julien

LULI, Ecole Polytechnique, CNRS, CEA, UPMC, France

"Intense Laser Pulses and Strong Magnetization as Tools for

Investigating Astrophysical Phenomena"

1 July 2016

Assoc Prof FURUMI, Seiichi

Faculty of Science Division I, Department of Applied Chemistry,

Tokyo University of Science, Tokyo, Japan

"Soft Photonic Devices Using Polymeric Self-Organization"

27 October 2016

Dr GOLOVKO, Vladimir

Department of Chemistry, Canterbury University, New Zealand

"Uncovering Structure-property Relationship: From Studies of Pure and Supported Metal Clusters Colloids to Their Use in Catalysis and Sensing"

6 July 2016

Assoc Prof GROMIHA, M. Michael

Indian Institute of Technology Madras, India

"Protein Interactions: Integrating Computational Methods and

Experimental Data for Understanding the Binding Specificity'

24 November 2016

Assist Prof HAMA, Tetsuya

Institute of Low Temperature Science, Hokkaido University,

Hokkaido, Japan

"Physical Chemistry on an Interfacial Phenomena: (2) Hydrogenbonding Dynamics of H₂O Molecules at a Cryogenic Temperature"

19 December 2016

Prof HAMADA, Hiroki

Department of Life Science, Okayama University of Science, Okayama, Japan

"Organic Synthesis by Plant Cell Culture"

16 April 2016

Prof HASSAGER, Ole

Technical University of Denmark, Denmark

"Applications of Controlled Filament Stretching in Extensional Rheology"

5 February 2016

Researcher HENDEL, Gregor

Zuse Institute Berlin, Berlin, Germany

"How to Solve Integer Optimization Problems with SCIP" 21 June 2016

Dr HERVE, Marie

Karlsruhe Institute of Technology, Germany

"Spin Polarized Scanning Tunneling Microscopy: From Probing Non Collinear Magnetic Configuration to the Investigation of Their High Frequency Dynamic Properties"

18 August 2016

Prof HOVEYDA, Amir H.

Department of Chemistry, Boston College, U.S.A.

"New Concepts, Catalysts and Methods for High-Impact Chemical Synthesis'

4 November 2016

Dr ILLIEN, Françoise

Laboratoire des Biomolécules, UMR 7203, Université P. et M. Curie - CNRS - ENS, Paris, France

"A New Method to Quantify Cell-Penetrating Peptides Translocation in Cells Based on Fluorescence: Application to Penetratin, R9 and Tat"

25 October 2016

Dr INGLESON, Michael

The University of Manchester, U.K.

"Direct Electrophilic Borylation for the Synthesis of Emissive Boron Materials and C(sp2)-Boronate Esters"

11 March 2016

Mr. ISHIDA, Toshinori

Division of Quantum Science and Engineering Faculty of Engineering, Hokkaido University, Hokkaido, Japan

"Development and Measurement of The Small-Angle Scattering Device in The Neutron Source with The Compact Accelerator" 20 September 2016

Prof ITO, Kimihito

Hokkaido University, Hokkaido, Japan

"Combining Population Genetics of Pathogens and Epidemiology of Infectious Diseases"

4 October 2016

Prof JANEŽIČ, Dušanka

University of Primorska, Slovenia

"Protein Binding Sites Dynamics in Drug Discovery" 16 November 2016

Dr JOHNSON, Brett C.

School of Physics, The University of Melbourne, Australia

"Single Photon Sources in Silicon Carbide"

23 September 2016

Dr JUNGFLEISCH, Benjamin

Argonne National Laboratory, U.S.A.

"Spin-torque Driven Magnetization Dynamics"

30 November 2016

Assist Prof KAGEYAMA, Yoshiyuki

Department of Chemistry, Faculty of Science, Hokkaido University, Hokkaido, Japan

"Mimic of Nature-Interdisciplinary Research between Mechanical Property and Dissipative Ordered Structure"

5 August 2016

Senior Researcher KINO, Koichi

Research Institute for Measurement and Analytical Instrumentation, National Institute of Advanced Industrial Science and Technology, Ibaraki, Japan

"Pulse Neutron Beam Imaging of The Lithium Ion Rechargeable Battery"

20 September 2016

Prof KLAUI, Mathias

University of Mainz, Germany

"Topology in Magnetic Materials"

25 November 2016

Prof KNAPP, Ernst-Walter

Free University of Berlin, Germany

"ProPairs: A Data Set for Protein-Protein Docking"

4 August 2016

Prof LEMIÈGRE, Loïc

Ecole Nationale Supérieure de Chimie de Rennes, France "Chemical and Chemoenzymatic Synthesis of Glycolipids" 7 November 2016

Prof MURAYAMA, Masafumi

Center for Advanced Marine Core Research, Kochi University, Nankoku, Japan

"What Occurred on the Shallow and Deep Seafloor by the Great East Japan Earthquake 2011?"

2 December 2016

Dr MUROOKA, Yoshie

Laboratory for Ultrafast Microscopy and Electron Scattering, École Polytechnique Fédérale de Lausanne (EPFL), Switzerland

"Filming Plasmonic Field and Magnetic Dynamics in Real Space and Time using ultrafast TEM"

12 July 2016

Dr NAKANISHI, Waka

MANA, National Institute for Materials Science, Ibaraki, Japan "Control of Molecular Structure at Interfaces" 28 June 2016

Assist Prof OTANI, Misato

Graduate School of Biological Sciences, Nara Institute of Science and Technology, Nara, Japan

"Regulation of RNA Metabolism Related with Efficient Plant Regeneration"

22 January 2016

Prof OZAWA, Takeaki

School of Science, The University of Tokyo, Tokyo, Japan

"Luminescent Sensors and Optical Switches for Single Cell Analysis" 28 January 2016

Dr RUBIO, Vicente

Centro Nacional de Biotecnología-CSIC, Madrid, Spain

"ABA Strikes Back: Inhibition of Plant CRL4 Ubiquitin Ligases Mediated by ABA"

28 November 2016

Assoc Prof SAEKI, Takayuki

High Energy Accelerator Research Organization, Ibaragi, Japan

"Production of the Superconducting Cavity"

10 October 2015

Prof SANDAA, Ruth-Anne

Bergen University, Norway, Norway

"Marine Viruses: Small but Important"

26 September 2016

Dr SCHAPER, Andreas Karl

Philipps University of Marburg, Germany

"Microscopy of Distortion in Crystals and of Hierarchical Ordering

in Macromolecular Systems"

14 October 2016

Prof SCHWECHHEIMER, Claus

Plant Systems Biology, Technische Universität München, Freising,

Germany

"PIN-ning Down the Function of D6PK Protein Kinases in Auxin

Transport'

28 November 2016

Prof SHIMIZU, Hirohiko M.

Laboratory for Particle Properties, Department of Physics, Nagoya

University, Aichi, Japan

"The Transmission Imaging by Stable Neutron Source"

15 January 2016

"Construction of the Core Technology of a Compact Neutron Source

for the Industrial Application"

11 March 2016

Dr SHINANO, Yuji

Zuse Institute Berlin, Berlin, Germany

"Towards Using over a Million CPU Cores to Solve Previously

Unsolved Mixed Integer Programming Problem Instances"

21 June 2016

Dr SONG, Jiangning

Monash University, Clayton, Australia

"Systematic Identification of Secreted Effector Proteins of Bacterial

Secretion System IV: A Bioinformatics Study'

30 November 2016

Prof STANG, Peter J.

The Universit of Utah, U.S.A.

"Abiological Self-Assembly: Predesigned Metallacycles and

Metallacages via Coordination"

22 April 2016

Dr TESILEANU, Ovidiu

Extreme Light Infrastructure - Nuclear Physics, Czech R.

"Project of Extreme Light Infrastructure for Nuclear Physics in

Czech Republic"

25 May 2016

Prof WANG, Yuan

College of Chemistry and Molecular Engineering, Peking University,

China, P.R.

"Molecular Engineering of Functional Systems based on

Nanoparticles"

16 September 2016

Dr YAMAMOTO, Kei

University of Mainz, Germany

"Spin Torque and Fluctuation-dissipation at FM-AFM Tunneling

Junctions'

18 November 2016

Dr YAMAMOTO, Noboru

Eisai Co., Ltd., Ibaraki, Japan

"Drug Discovery Created by Chemists-From Drug Design to

Biomarker"

20 May 2016

Prof YAMANAKA, Masahiro

Department of Chemistry, Rikkyo University, Tokyo, Japan

"Investigation of Stereoselectivity in Organocatalysis: Toward

Rational Design Based on Transition-State Model"

8 December 2016

Prof YANG, Jinn-Moon

National Chiao Tung University, Hsinchu, Taiwan

"NetPharmalogs for Drug Discovery and Applications"

20 April 2016

Dr YASUDA, Nobuyoshi

Merck Research Institute, MERCK&CO., INC., U.S.A.

"The Art of Process Chemistry"

8 November 2016

Assoc Prof ZHU, Shanfeng

Fudan University, China, P.R.

"DeepMeSH and MeSHLabeler: Recent Progress in Large-scale

MeSH Indexing"

29 August 2016

"DrugE-Rank: Improving Drug-target Interaction Prediction of New

Candidate Drugs or Targets by Ensemble Learning to Rank"

2 September 2016

Dr ZINOVYEV, Andrei

Computational Systems Biology of Cancer group, Institut Curie,

France

"The Shortest Path is not the One You Know: Application of

Biological Networks in Precision Oncology Research"

13 December 2016

MEETINGS AND SYMPOSIA

Asian Chemical Biology Initiative 2016 Jakarta Meeting

Organized by UESUGI, Motonari 29 January-1 February 2016 (Jakarta, Indonesia)

The 11th International Workshop for East Asian Young Rheologists (IWEAYR-11)

Co-organized by WATANABE, Hiroshi 4-7 February 2016 (Shenzhen, China, P.R.)

7th Thin-film Solar Cells Seminar

Organized by YOSHIDA, Tsukasa; WAKAMIYA, Atsushi 13-15 May 2016 (Yonezawa, Japan)

The 11th Organoelement Chemistry Seminar

Organized by TOKITOH, Norihiro; SASAMORI, Takahiro; MIZUHATA, Yoshiyuki 3-4 June 2016 (Uji, Japan)

The 5th Main Group Element Chemistry Seminar

Organized by WAKAMIYA, Atsushi 3 August 2016 (Mie, Japan)

17th International Congress on Rheology

Chaired by WATANABE, Hiroshi 8-13 August 2016 (Kyoto, Japan)

Catalyst Society of Japan

"The 44th Organometallic Seminar"

Organized by OZAWA, Fumiyuki; NAKAMURA, Masaharu 24 October 2016 (Uji, Japan)

2016 JSME Annual Meeting

"Symposium S4: Would You Like to Discuss the Raison D'être for Viruses?"

Organized by OGATA, Hiroyuki; NAGASAKI, Keizo 24 October 2016 (Yokosuka, Japan)

The 1st Frontier Solar Cells Seminar

Organized by WAKAMIYA, Atsushi 17-18 November 2016 (Uji, Japan)

Workshop on pMAIRS: Innovative Analytical Technique for an Amorphous Thin Film Device with a Surface Roughness

Organized by HASEGAWA, Takeshi 2 December 2016 (Tokyo, Japan)

The 39th Annual Meeting of the Molecular Biology Society of

"Symposium 3AS11: Where Did They Come From? Where Do They Habit? and What Do They Do in Our World?" Organized by OGATA, Hiroyuki; TAKEMURA, Masaharu 2 December 2016 (Yokohama, Japan)

The 8th Asian Conference on Organic Electronics (A-COE 2016)

Organized by KAJI, Hironori 5–7 December 2016 (Uji, Japan)

INDEX

NAME INDEX

[A]		ELIDLIKAWA Ko	76	IHADA Tochiyadi	56
[A] ADAK, Laksmikanta	50	FURUKAWA, Ko FURUKAWA, Takahiro	34	IHARA, Toshiyuki IINO, Masatomo	56 20
AGATA, Ryosuke	50	FURUKAWA, Yuki	44, 80	IKEDA, Hideki	42
AHAREN, Tomoko	6	FURUTA, Takumi	8	IMAJO, Sohei	42
AKISHIBA, Misao	20, 85	FURUTA, Tomoyuki	26	IMANISHI, Miki	20
AKIYAMA, Yutaka	76	FUTAGOISHI, Tsukasa	6, 84	INOUE, Shota	46
AKIZUKI, Maika	26	FUTAKI, Shiroh	20	INOUE, Shunsuke	44, 83
AKUTSU, Tatsuya	60, 80	FUWA, Takuto	36	ISE, Shoko	20
ANDO, Fuyuki	18	FUWA, Yasuhiro	42	ISHIBASHI, Mio	18
AOKI, Yuma	50			ISHIBASHI, Takuma	38
AOYAMA, Takashi	24	[G]		ISHIKURA, Yasuhisa	6
ARAFILES, Jan Vincent	20	GOETZE, Sebastian Andreas	26	ISHIMOTO, Ryushin	38
ARAI, Kenta	8	GONDOH, Naruhiro	8	ISHIMOTO, Shintaro	16
ARAKAWA, Noriko	6	GOTO, Kakeru	24	ISOZAKI, Katsuhiro	50
ARAMAKI, Takuya	58	GOTO, Susumu	58	ITO, Shota	22
ASANO, Risa	26	GUO, Haichuan	52	ITO, Yoshiaki	48
ASHIDA, Takaki	16	GUO, Jing-Dong	4, 82	IWAMOTO, Takahiro	50
[12]		IIII		IWASHITA, Yoshihisa	42 32
[B] BAO, Yu	60	[H] HABUKA, Yume	28	IZUMI, Naoki	32
BLANC-MATHIEU, Romain	58	HADA, Miyako	32	[J]	
BLANC-MATTILO, Romani	30	HAKARIYA, Hayase	26	JAMESON, Robert A.	42
[C]		HANDA, Taketo	56	JANSSON, Jesper	66
CAO, Yu	60	HARADA, Kazuki	54	JEONG, Jaeyoung	14
CHAOLUMEN	6	HARAYAMA, Seiya	12	JIRA, Jindalertudomdee	60
CHEN, Chen	34	HARISADHAN, Ghosh	8	JUNG, Mina	6
CHEN, Gong	8	HARUTA, Mitsutaka	46, 80	JUNG, Yejin	26
CHEN, Lihui	10	HASEGAWA, Takeshi	32		
CHEN, Tsai-Min	34	HASHIDA, Masaki	44, 83	[K]	
CHENG, Xiaoqing	60	HASHIKAWA, Yoshifumi	6	KAJI, Hironori	28, 82
CHUNG, Watson Xie Khim	26	HASHIMOTO, Hisashi	8	KAJI, Shoji	6
		HASHIMOTO, Sigma	14	KAKIZAKAI, Haruka	18
[D]		HATSUHARA, Hikari	34	KAMIYA, Michinari	18
DANJYO, Takuya	16	HAYASAKA, Manami	8	KAN, Daisuke	52
DE LA HIGUERA, Colin Manue		HAYASHI, Kan	16	KANEHISA, Minoru	76
DOI, Yuya	38	HAYASHI, Kazuhiro	8	KANEKO, Syunichi	16
rm.		HAYASHI, Shuhei	20	KANEMITSU, Yoshihiko	56
[E]	10	HAYASHI, Toshiki	14	KASAMATSU, Koji	8, 78, 84
EGAWA, Shigehisa	10	HAYASHIDA, Morihiro	60	KATSUDA, Yousuke	26, 79
EGI, Kohki	12	HIGAKI, Sotaro	28	KAWABATA, Takeo	8
EGUCHI, Daichi	10, 84	HIGASHINO, Naruhisa	36 46	KAWAI, Soichiro	34
[F]		HIGUCHI, Hiroki HIRANO, Koki	40	KAWAKAMI, Yuko KAWAMOTO, Jun	60 34
FABIO, Denis Romero	52	HIRANO, Toshiko	4	KAWANO, Kenichi	20, 85
FAN, Lulu	50	HIRATA, Atsushi	8	KAWANO, Ryuji	76
FAN, Weijia	14	HIRATA, Yushou	18	KAWAWAKI, Tokuhisa	10
FUJIHASHI, Akiko	8	HIRATAKE, Jun	22	KAYAHARA, Eiichi	14
FUJII, Tomomi	48	HIROSHIGE, Nao	56, 86	KIDO, Tsuyoshi	28
FUJIMORI, Shiori					
FILIMODI T	4, 84	HISATOMI, Tatsuya	12	KIM, Duck-Ho	18
FUJIMOKI, Iomoya	4, 84 12, 85	HISATOMI, Tatsuya HOKONOHARA, Hisao		KIM, Duck-Ho KIM, Sanghoon	18 18
FUJIMORI, Tomoya FUJIMOTO, Junji			12	KIM, Duck-Ho KIM, Sanghoon KIM, Sungwon	
	12, 85	HOKONOHARA, Hisao	12 42	KIM, Sanghoon	18
FUJIMOTO, Junji FUJIMOTO, Syou FUJIMOTO, Takumi	12, 85 18 16 8	HOKONOHARA, Hisao HONJO, Kazuma	12 42 4	KIM, Sanghoon KIM, Sungwon	18 10
FUJIMOTO, Junji FUJIMOTO, Syou FUJIMOTO, Takumi FUJITA, Shuuhei	12, 85 18 16 8 36	HOKONOHARA, Hisao HONJO, Kazuma HONORE, Jean-Claude HOSAKA, Yoshiteru HOSOKAWA, Atsushi	12 42 4 14 52, 80 50	KIM, Sanghoon KIM, Sungwon KIMURA, Masato KIMURA, Syunsaku KINOSE, Yuji	18 10 10 34 12
FUJIMOTO, Junji FUJIMOTO, Syou FUJIMOTO, Takumi FUJITA, Shuuhei FUJITA, Takehiro	12, 85 18 16 8 36 14	HOKONOHARA, Hisao HONJO, Kazuma HONORE, Jean-Claude HOSAKA, Yoshiteru HOSOKAWA, Atsushi HOU, Wenpin	12 42 4 14 52, 80 50 60	KIM, Sanghoon KIM, Sungwon KIMURA, Masato KIMURA, Syunsaku KINOSE, Yuji KIRIHATA, Tatsuro	18 10 10 34 12 62
FUJIMOTO, Junji FUJIMOTO, Syou FUJIMOTO, Takumi FUJITA, Shuuhei FUJITA, Takehiro FUJIWARA, Masanori	12, 85 18 16 8 36 14 16	HOKONOHARA, Hisao HONJO, Kazuma HONORE, Jean-Claude HOSAKA, Yoshiteru HOSOKAWA, Atsushi HOU, Wenpin HSU, Shih-Chen	12 42 4 14 52, 80 50 60 10	KIM, Sanghoon KIM, Sungwon KIMURA, Masato KIMURA, Syunsaku KINOSE, Yuji KIRIHATA, Tatsuro KISE, Ryuma	18 10 10 34 12 62 32
FUJIMOTO, Junji FUJIMOTO, Syou FUJIMOTO, Takumi FUJITA, Shuuhei FUJITA, Takehiro FUJIWARA, Masanori FUJIWARA, Takashi	12, 85 18 16 8 36 14 16 24	HOKONOHARA, Hisao HONJO, Kazuma HONORE, Jean-Claude HOSAKA, Yoshiteru HOSOKAWA, Atsushi HOU, Wenpin	12 42 4 14 52, 80 50 60	KIM, Sanghoon KIM, Sungwon KIMURA, Masato KIMURA, Syunsaku KINOSE, Yuji KIRIHATA, Tatsuro KISE, Ryuma KITAHARA, Ryunosuke	18 10 10 34 12 62 32 42
FUJIMOTO, Junji FUJIMOTO, Syou FUJIMOTO, Takumi FUJITA, Shuuhei FUJITA, Takehiro FUJIWARA, Masanori FUJIWARA, Takashi FUJIWARA-KATO, Mariko	12, 85 18 16 8 36 14 16 24 24	HOKONOHARA, Hisao HONJO, Kazuma HONORE, Jean-Claude HOSAKA, Yoshiteru HOSOKAWA, Atsushi HOU, Wenpin HSU, Shih-Chen HUA SHEN, Zhiren	12 42 4 14 52, 80 50 60 10	KIM, Sanghoon KIM, Sungwon KIMURA, Masato KIMURA, Syunsaku KINOSE, Yuji KIRIHATA, Tatsuro KISE, Ryuma KITAHARA, Ryunosuke KITAYAMA, Kaori	18 10 10 34 12 62 32 42 34
FUJIMOTO, Junji FUJIMOTO, Syou FUJIMOTO, Takumi FUJITA, Shuuhei FUJITA, Takehiro FUJIWARA, Masanori FUJIWARA, Takashi FUJIWARA-KATO, Mariko FUJIYOSHI, Yoshifumi	12, 85 18 16 8 36 14 16 24 24 46	HOKONOHARA, Hisao HONJO, Kazuma HONORE, Jean-Claude HOSAKA, Yoshiteru HOSOKAWA, Atsushi HOU, Wenpin HSU, Shih-Chen HUA SHEN, Zhiren	12 42 4 14 52, 80 50 60 10 14	KIM, Sanghoon KIM, Sungwon KIMURA, Masato KIMURA, Syunsaku KINOSE, Yuji KIRIHATA, Tatsuro KISE, Ryuma KITAHARA, Ryunosuke KITAYAMA, Kaori KIYOMURA, Tsutomu	18 10 10 34 12 62 32 42 34 46
FUJIMOTO, Junji FUJIMOTO, Syou FUJIMOTO, Takumi FUJITA, Shuuhei FUJITA, Takehiro FUJIWARA, Masanori FUJIWARA, Takashi FUJIWARA-KATO, Mariko	12, 85 18 16 8 36 14 16 24 24	HOKONOHARA, Hisao HONJO, Kazuma HONORE, Jean-Claude HOSAKA, Yoshiteru HOSOKAWA, Atsushi HOU, Wenpin HSU, Shih-Chen HUA SHEN, Zhiren	12 42 4 14 52, 80 50 60 10	KIM, Sanghoon KIM, Sungwon KIMURA, Masato KIMURA, Syunsaku KINOSE, Yuji KIRIHATA, Tatsuro KISE, Ryuma KITAHARA, Ryunosuke KITAYAMA, Kaori	18 10 10 34 12 62 32 42 34

KONO, Taro	14	MORI, Teppei	16	OKAMOTO, Masanori	36
KOTAKE, Kenjiro	26	MORIKI, Yoshito	12	OKAMOTO, Shu	6
KUBO, Kosei	38	MORISHITA, Hiroki	16, 82	OKAMOTO, Yasuo	10
KUBO, Shosei	28, 82	MORITA, Hazuki	54	OKAZAKI, Shuhei	6
KUMAGAI, Fumihito	34	MORIYAMA, Takahiro	18	OKITA, Shutaro	30
KUMEGAWA, Ryo	8	MOTOMURA, Yoshihiro	38	OKUDA, Satoshi	26
, ,	46				18
KURATA, Hiroki		MURAI, Takuya	8	OKUNO, Takaya	
KURIBAYASHI, Toshifumi	8	MURAI, Takuya	16	OKUZONO, Chiemi	50
KURIHARA, Tatsuo	34	MURAJI, Koichi	12	OMATSU, Yamato	4
KURODA, Ryo	24	MURATA, Masahiro	14	OMOSU, Takuya	36
KURONISHI, Megumi	58	MURATA, Michihisa	6	ONO, Teruo	18
KUROSAKI, Satoshi	36	MURATA, Naoki	6	ORITA, Takuya	8
KUSUDA, Toshiyuki	18	MURATA, Yasujiro	6	OSAWA, Masatoshi	32
KWON, Oh Min	38	MURAYAMA, Masafumi	76	OTA, Masaki	34
KWON, OII WIII	50		20, 79		30
T 1		MURAYAMA, Tomo		OTSUKA, Yuhei	
[L]		MURDEY, Richard James	40	OTSUKI, Haruka	8, 84
LAI, Ming Wei	46			OUE, Shohei	14
LE, Phuong Quang	56	[N]		OZAKI, Masashi	6, 84
LE, Toan Khanh	26	NAESHIRO, Kazuki	22	OZAKI, Yusuke	52
LEE, Jaehyun	6	NAGAISHI, Masaru	8	OZAWA, Fumiyuki	54
LI, Ruiming	60	NAGANO, Yoshiki	36		
LI, Tian	18	NAKAGAWA, Hiromichi	28	[P]	
· · · · · · · · · · · · · · · · · · ·					1.4
LI, Wenjuan	14	NAKAJIMA, Natsu	81	PARK, Beomsu	14
LI, Xian	22	NAKAJIMA, Sho	50	PATEL, Vijay Kumar	14
LI, Xiaopei	14	NAKAJIMA, Yuuki	12	PE, Kathleen Beverly Alog	26
LI, Yanhui	10	NAKAMICHI, Kimiyo	12	PERRON, Amelie	26
LI, Yanze	58	NAKAMIYA, Yoshihide	44	PI, Wenya	60
LIAN, Zichao	10	NAKAMURA, Masaharu	50	PINCELLA, Francesca	50
LIAO, Youqi	26	NAKAMURA, Tomoya	6	PRODINGER, Florian	58
LIN, Xiaoya	24	NAKAMURA, Yasuyuki	14, 78	PUNZALAN, Louvy Lynn	26
	14			FUNZALAN, LOUVY LYIII	20
LIN, Yiming		NAKANISHI, Yohei	12		
LIU, Pengyu	60	NAKANO, Tomohiro	4	[Q]	
					26
LU, Wenjie	8, 78	NAKASHIMA, Mitsue	26	QIN, Ying	
LU, Yangtian	8, 78 14	NEMOTO, Takashi	26 46	QIN, Ying QU, Rui	14
LU, Yangtian		NEMOTO, Takashi NGUYEN, Canh Hao	46 62	QU, Rui	
LU, Yangtian [M]	14	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo	46 62 8	QU, Rui [R]	14
LU, Yangtian [M] MAENO, Ayaka	14 28	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji	46 62 8 36	QU, Rui	
LU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar	28 4	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe	46 62 8 36 52	QU, Rui [R] RAKKAPAO, Natthida	14
LU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori	28 4 22	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu	46 62 8 36 52 36	QU, Rui [R] RAKKAPAO, Natthida [S]	38
LU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi	28 4 22 62	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo	46 62 8 36 52 36 14	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki	14 38 42
LU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di	28 4 22 62 26	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma	46 62 8 36 52 36 14 38	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo	14 38 42 22
LU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi	28 4 22 62 26 16	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo	46 62 8 36 52 36 14	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki	14 38 42
LU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di	28 4 22 62 26	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma	46 62 8 36 52 36 14 38	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo	14 38 42 22
LU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu	28 4 22 62 26 16 16, 79	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio	46 62 8 36 52 36 14 38 10	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi	14 38 42 22 52, 80
LU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro	28 4 22 62 26 16 16, 79 20, 85	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki	46 62 8 36 52 36 14 38 10 12 58	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki	14 38 42 22 52, 80 44, 83 20
LU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro	28 4 22 62 26 16 16, 79 20, 85 44	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota	46 62 8 36 52 36 14 38 10 12 58	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita	14 38 42 22 52, 80 44, 83 20 12, 78
LU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki	28 4 22 62 26 16 16, 79 20, 85 44 8	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki	46 62 8 36 52 36 14 38 10 12 58 12 52	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou	14 38 42 22 52, 80 44, 83 20 12, 78 20
LU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi	28 4 22 62 26 16 16, 79 20, 85 44 8 50	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki	46 62 8 36 52 36 14 38 10 12 58 12 52 14	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori	14 38 42 22 52, 80 44, 83 20 12, 78 20 10
LU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi	46 62 8 36 52 36 14 38 10 12 58 12 52 14	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 12
IU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUKAWA, Kimihiro MATSUMIYA, Yumi	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 12 10
IU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUKAWA, Kimihiro MATSUMIYA, Yumi MATSUMOTO, Kenshi	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi	46 62 8 36 52 36 14 38 10 12 58 12 52 14	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 12 10 54
IU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUKAWA, Kimihiro MATSUMIYA, Yumi	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 12 10
IU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUKAWA, Kimihiro MATSUMIYA, Yumi MATSUMOTO, Kenshi	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Yuichiro	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 12 10 54 20
IU, Yangtian [M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUKAWA, Kimihiro MATSUMIYA, Yumi MATSUNOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58	QU, Rui [R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Yuichiro SASAMORI, Takahiro	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 12 10 54 20 4, 82
IMJ MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUKAWA, Kimihiro MATSUMIYA, Yumi MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Yuichiro SASAMORI, Takahiro SATO, Ohji	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 12 10 54 20 4, 82 38
IMJ MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUKAWA, Kimihiro MATSUMIYA, Yumi MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Vuichiro SASAMORI, Takahiro SATO, Ohji SATO, Ryota	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 54 20 4, 82 38 10
IMJ MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUKAWA, Kimihiro MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke MINAMI, Tomoharu	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16 30	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento OGATA, Hiroyuki	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Takahiro SATO, Ohji SATO, Ryota SATO, Shinichi	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 54 20 4, 82 38 10 26
IMJ MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUKAWA, Kimihiro MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke MINAMI, Tomoharu MIURA, Tsutomu	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16 30 28	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHIMO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento OGATA, Hiroyuki OGATA, Isamu	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Yuichiro SASAMORI, Takahiro SATO, Ohji SATO, Ryota SATO, Shinichi SCHAPER, Andreas Karl	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 54 20 4, 82 38 10 26 46
IMJ MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUMIYA, Yumi MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke MINAMI, Tomoharu MIURA, Tsutomu MIYAKE, Harumi	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16 30 28 28	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHIMO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIYAMA, Hiroki NIWA, Ryota NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento OGATA, Hiroyuki OGATA, Isamu OGAWA, Hiroki	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Yuichiro SASAMORI, Takahiro SATO, Ohji SATO, Ryota SATO, Shinichi SCHAPER, Andreas Karl SEKI, Toyomitsu	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 54 20 4, 82 38 10 26 46 14, 85
IMJ MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUMIYA, Yumi MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke MINAMI, Tomoharu MIURA, Tsutomu MIYAKE, Harumi MIYASAKA, Yasuhiro	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16 30 28 28 83	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHIMO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIVAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento OGATA, Hiroyuki OGAWA, Hiroki OGAWA, Takuya	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Vuichiro SASAMORI, Takahiro SATO, Ohji SATO, Ryota SATO, Shinichi SCHAPER, Andreas Karl SEKI, Toyomitsu SHANOH, Takafumi	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 54 20 4, 82 38 10 26 46 14, 85 50
[M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUMIYA, Yumi MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke MINAMI, Tomoharu MIURA, Tsutomu MIYAKE, Harumi MIYASAKA, Yasuhiro MIYATAKE, Yuki	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16 30 28 28 83 36	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHIMO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIVAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento OGATA, Hiroyuki OGAWA, Hiroki OGAWA, Takuya OGAWA, Takuya	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Puichiro SASAKI, Takahiro SATO, Ohji SATO, Ohji SATO, Ryota SATO, Shinichi SCHAPER, Andreas Karl SEKI, Toyomitsu SHANOH, Takafumi SHIBAYAMA, Hiromitsu	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 54 20 4, 82 38 10 26 46 14, 85 50 8
IMJ MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUMIYA, Yumi MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke MINAMI, Tomoharu MIURA, Tsutomu MIYAKE, Harumi MIYASAKA, Yasuhiro	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16 30 28 28 83	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHIMO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIVAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento OGATA, Hiroyuki OGAWA, Hiroki OGAWA, Takuya	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Vuichiro SASAMORI, Takahiro SATO, Ohji SATO, Ryota SATO, Shinichi SCHAPER, Andreas Karl SEKI, Toyomitsu SHANOH, Takafumi	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 54 20 4, 82 38 10 26 46 14, 85 50
[M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUKAWA, Kimihiro MATSUMIYA, Yumi MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke MINAMI, Tomoharu MIURA, Tsutomu MIYAKE, Harumi MIYASAKA, Yasuhiro MIYATAKE, Yuki	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16 30 28 28 83 36	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHIMO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIVAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento OGATA, Hiroyuki OGAWA, Hiroki OGAWA, Takuya OGAWA, Takuya	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Puichiro SASAKI, Takahiro SATO, Ohji SATO, Ohji SATO, Ryota SATO, Shinichi SCHAPER, Andreas Karl SEKI, Toyomitsu SHANOH, Takafumi SHIBAYAMA, Hiromitsu	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 54 20 4, 82 38 10 26 46 14, 85 50 8
IMJ MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUDA, Hiroshi MATSUMAYA, Yumi MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke MINAMI, Tomoharu MIURA, Tsutomu MIYAKE, Harumi MIYASAKA, Yasuhiro MIYATAKE, Yuki MIYAWAKI, Eisuke MIYOSHI, Shun-ichiro	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16 30 28 28 83 36 42 14	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHIMO, Hideo NISHIMO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIVAMA, Hiroki NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento OGATA, Hiroyuki OGAWA, Hiroki OGAWA, Takuya OGAWA, Tetsuya OGAWA, Yoshihiro OGUSHI, Masaya	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Takahiro SASAMORI, Takahiro SATO, Ohji SATO, Ryota SATO, Shinichi SCHAPER, Andreas Karl SEKI, Toyomitsu SHANOH, Takafumi SHIBAYAMA, Hiromitsu SHIMAKAWA, Yuichi SHIMAMURA, Ryota	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 54 20 4, 82 38 10 26 46 14, 85 50 8 52, 80 24
IMJ MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUDA, Hiroshi MATSUMAYA, Yumi MATSUMOTO, Kenshi MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke MINAMI, Tomoharu MIURA, Tsutomu MIYAKE, Harumi MIYASAKA, Yasuhiro MIYATAKE, Yuki MIYAWAKI, Eisuke MIYOSHI, Shun-ichiro MIZUHATA, Yoshiyuki	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16 30 28 28 83 36 42 14 4, 78	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHIMO, Hideo NISHIMO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIVAMA, Hiroki NIWA, Ryota NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento OGATA, Hiroyuki OGATA, Isamu OGAWA, Takuya OGAWA, Tetsuya OGAWA, Yoshihiro OGUSHI, Masaya OHKANDA, Junko	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58 58 58 59 40 46 76 20 26, 79	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Takahiro SASAMORI, Takahiro SATO, Ohji SATO, Ohji SATO, Ryota SATO, Shinichi SCHAPER, Andreas Karl SEKI, Toyomitsu SHANOH, Takafumi SHIBAYAMA, Hiromitsu SHIMAKAWA, Yuichi SHIMAMURA, Ryota SHIMAZAKI, Ai	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 54 20 4, 82 38 10 26 46 14, 85 50 8 52, 80 24 6
[M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUDA, Hiroshi MATSUDA, Genshi MATSUMOTO, Kenshi MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke MINAMI, Tomoharu MIURA, Tsutomu MIYAKE, Harumi MIYASAKA, Yasuhiro MIYATAKE, Yuki MIYAWAKI, Eisuke MIYOSHI, Shun-ichiro MIZUHATA, Yoshiyuki MIZUNO, Hayato	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16 30 28 28 83 36 42 14 4, 78 18	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIVAMA, Hiroki NIWA, Ryota NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento OGATA, Hiroyuki OGATA, Isamu OGAWA, Takuya OGAWA, Tetsuya OGAWA, Yoshihiro OGUSHI, Masaya OHKANDA, Junko OHKE, Yoshie	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58 58 52 14 46 76 20 26, 79 34	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Vuichiro SASAMORI, Takahiro SATO, Ohji SATO, Ohji SATO, Shinichi SCHAPER, Andreas Karl SEKI, Toyomitsu SHANOH, Takafumi SHIBAYAMA, Hiromitsu SHIMAKAWA, Yuichi SHIMAKAWA, Yuichi SHIMAMURA, Ryota SHIMAZAKI, Ai SHIMIZU, Hirohiko M.	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 54 20 4, 82 38 10 26 46 14, 85 50 8 52, 80 24 6 42
[M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUDA, Hiroshi MATSUMIYA, Yumi MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke MINAMI, Tomoharu MIVRA, Tsutomu MIYAKE, Harumi MIYASAKA, Yasuhiro MIYATAKE, Yuki MIYAWAKI, Eisuke MIYOSHI, Shun-ichiro MIZUHATA, Yoshiyuki MIZUNO, Hayato MIZUNO, Hiroki	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16 30 28 28 83 36 42 14 4, 78 18 10	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIVAMA, Hiroki NIWA, Ryota NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Hiroshi NOMURA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento OGATA, Hiroyuki OGATA, Isamu OGAWA, Takuya OGAWA, Tetsuya OGAWA, Yoshihiro OGUSHI, Masaya OHKANDA, Junko OHKE, Yoshie OHKITA, Yuya	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58 58 52 14 46 76 20 26, 79 34 38	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Vuichiro SASAMORI, Takahiro SATO, Ohji SATO, Ohji SATO, Shinichi SCHAPER, Andreas Karl SEKI, Toyomitsu SHANOH, Takafumi SHIBAYAMA, Hiromitsu SHIMAKAWA, Yuichi SHIMAKAWA, Yuichi SHIMAMURA, Ryota SHIMAZAKI, Ai SHIMIZU, Hirohiko M. SHIMIZU, Yoshihiko	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 12 10 54 20 4, 82 38 10 26 46 14, 85 50 8 52, 80 24 6 42 12
[M] MAENO, Ayaka MAJHI, Paresh Kumar MAKINO, Katsunori MAMITSUKA, Hiroshi MAO, Di MARUYAMA, Yuuichi MASAI, Hirokazu MASUDA, Toshihiro MASUNO, Shinichiro MATAYOSHI, Aki MATSUDA, Hiroshi MATSUDA, Hiroshi MATSUDA, Genshi MATSUMOTO, Kenshi MATSUMOTO, Kenshi MATSUNAGA, Tsubasa MATSUO, Yutaka MENDOZA, Aileen de-Leon MIMA, Daisuke MINAMI, Tomoharu MIURA, Tsutomu MIYAKE, Harumi MIYASAKA, Yasuhiro MIYATAKE, Yuki MIYAWAKI, Eisuke MIYOSHI, Shun-ichiro MIZUHATA, Yoshiyuki MIZUNO, Hayato	28 4 22 62 26 16 16, 79 20, 85 44 8 50 12 38 10 36 76 26 16 30 28 28 83 36 42 14 4, 78 18	NEMOTO, Takashi NGUYEN, Canh Hao NINOMIYA, Ryo NISHIDA, Koji NISHIMURA, Tomoe NISHIMURA, Yu NISHINO, Hideo NISHINO, Takuma NISHINO, Toshio NISHIUMI, Kosuke NISHIVAMA, Hiroki NIWA, Ryota NIWA, Ryota NIWA, Yasuyuki NOGUCHI, Naoki NOMURA, Hiroshi NOMURA, Kayo NURSHAHIRA, Binti Yusuf [O] O'BRIEN, Harry ODA, Kento OGATA, Hiroyuki OGATA, Isamu OGAWA, Takuya OGAWA, Tetsuya OGAWA, Yoshihiro OGUSHI, Masaya OHKANDA, Junko OHKE, Yoshie	46 62 8 36 52 36 14 38 10 12 58 12 52 14 14 20 58 58 52 14 46 76 20 26, 79 34	[R] RAKKAPAO, Natthida [S] SAEKI, Takayuki SAITO, Shingo SAITO, Takashi SAKABE, Shuji SAKAGAMI, Koki SAKAKIBARA, Keita SAKAMOTO, Kentarou SAKAMOTO, Masanori SAKAUE, Motokazu SARUYAMA, Masaki SASAKI, Daichi SASAKI, Vuichiro SASAMORI, Takahiro SATO, Ohji SATO, Ohji SATO, Shinichi SCHAPER, Andreas Karl SEKI, Toyomitsu SHANOH, Takafumi SHIBAYAMA, Hiromitsu SHIMAKAWA, Yuichi SHIMAKAWA, Yuichi SHIMAMURA, Ryota SHIMAZAKI, Ai SHIMIZU, Hirohiko M.	14 38 42 22 52, 80 44, 83 20 12, 78 20 10 54 20 4, 82 38 10 26 46 14, 85 50 8 52, 80 24 6 42

SHIMOAKA, Takafumi	32, 79	[U]		ZHANG, Xiao-Juan	24
SHIMODA, Shotaro	8	UEDA, Yoshihiro	8, 78	ZHAO, Chenzhou	12
SHIMOGAWA, Hiroyuki	6	UEHARA, Wataru	30	ZHENG, Linjie	30
SHINODA, Kouki	20	UENO, Ryo	50	ZHU, Shanfeng	76
SHIOYA, Nobutaka	32, 79	UESUGI, Motonari	26		
SHITARA, Daichi	14	UKAI, Hironori	32		
SHIZU, Katsuyuki	28, 82	UMETANI, Shigeo	30		
SIMONE, Marini	60				
SIMO-OKA, Takaaki	16	[V]			
SOHRIN, Yoshiki	30, 79	VINCHENT, Alexandra	60		
SUGAHARA, Tomohiro	4, 78, 82, 84	VU, Hue Thi	26		
SUGIMOTO, Saki	34				
SUGIOKA, Koji	76	[W]			
SUGIURA, Miwa	34	WADA, Tamaki	52		
SUN, Liansheng	14	WADA, Yoshimasa	28, 79, 82		
SUWANAWAT, Nittikarn	34	WAKAI, Chihiro	32		
SUZUKI, Katsuaki	28, 82	WAKAMIYA, Atsushi	6, 82		
SUZUKI, Yuko	4, 78	WAKAYAMA, Rin	54		
,	,	WAKIOKA, Masayuki	54		
[T]		WANG, Shuo	8		
TABUCHI, Michimasa	10	WATANABE, Bunta	22		
TAGUCHI, Hiro-omi	54, 85	WATANABE, Hiroshi	38		
TAHARA, Hirokazu	56, 80	WATANABE, Kensuke	20		
TAJIRI, Ken	26	WATARI, Machiko	24		
	54		62		
TAKAHASHI, Rina		WIMALAWARNE, Kishan	02		
TAKAHASHI, Yoshiaki	76	IVI			
TAKAHASHI, Yuuya	16	VIONG Para	50		
TAKANO, Shotaro	30	XIONG, Peng	52		
TAKASHIMA, Ippei	26	XU, Kai	54		
TAKASHIMA, Megumi	8				
TAKAYA, Hikaru	50	[Y]	2.4		
TAKAYA, Junichiro	26	YAGURA, Kazuki	34		
TAKEMOTO, Yasushi	26	YAHATA, Yoshikazu	12		
TAKENAKA, Mikihito	36	YAKUMARU, Shinya	6		
TAKENOYA, Mihoko	26	YAMADA, Kihiro	18		
TAKEUCHI, Katsuhiko	54	YAMADA, Makoto	62, 83		
TAMEMOTO, Naoki	20	YAMADA, Takumi	56, 80		
TAMURA, Takeyuki	60	YAMAGO, Shigeru	14		
TAN, Zhenhong	52	YAMAGUCHI, Atsushi	46		
TANAKA, Asako	34	YAMAGUCHI, Toshiaki	34		
TANAKA, Kensho	18	YAMAMOTO, Rumiko	58		
TANAKA, Yuto	54	YAMAMOTO, Satoru	8		
TANIGUCHI, Takuya	18	YAMASHITA, Kohei	12, 85		
TANISHIMA, Miho	12	YAMASHITA, Natsumi	54		
TANTIPANJAPORN, Ajcha	arapan 26	YAMASHITA, Shohei	58		
TERAMOTO, Kensuke	44, 80	YANADA, Mizuho	12		
TERANISHI, Toshiharu	10	YANAGISAWA, Tatsuya	4		
TEX, David Michael	56	YANG, Baochen	8		
TOHZAKI, Yudai	62	YANG, Jinn-Moon	60		
TOKITA, Shigeki	44	YANO, Yuko	14		
TOKITOH, Norihiro	4, 82	YAO, Cheng	14		
TOKUDA, Youmei	16	YARITA, Naoki	56		
TOKUNAGA, Tomohisa	34	YASUDA, Keiko	24		
TONGU, Hiromu	42	YASUDA, Shuma	12		
TORIMOTO, Aya	16, 79, 85	YATSUZUKA, Kenji	26		
TOSAKA, Masatoshi	14	YELLA, Ramesh	8		
TOYOTAKE, Yosuke	34	YOKOYAMA, Fumiaki	34		
TRINH, Thang Thuy	10	YOSHIDA, Hiroki	26		
TSUGE, Tomohiko	24	YOSHIDA, Keisuke	8		
TSUJI, Shogo	20				
	28	YOSHIDA, Ryota	50, 80 58		
TSUJI, Tomohiko		YOSHIKAWA, Genki	38 79		
TSUJII, Yoshinobu	12, 78	YOSHIMURA, Yoko			
TSUJIOKA, Kouta	36	YOSHIZAWA, Akiyasu	58		
TSUJISAKA, Makoto	30	1721			
TSUKAO, Masahiro	6, 84	[Z]	(70		
TUAN, Watie Binti Tuan M	1at 58	ZHANG, Rui	6, 78		

KEYWORD INDEX

[A]		Homogeneous Catalyst	54	Polymer Physics	36
Accelerator Physics	42	Hybrid Materials	12		, 36
Amorphous Materials	28	Try orie triaterials	12	Polymer Synthesis	14
Analytical Chemistry	30	[I]		Polyunsaturated Fatty Acid	34
Artificial Materials	18	Infrared and Raman Spectroscopy	32	Precision Polymerization	12
Artificial Waterials	10	Inorganic Nanoparticles	10	Protein Complexes	60
(D)		Intense Laser Science	44		48
[B] Bacterial Cold-adaptation Mecha	miam 24	International Linear Collider	42	Protein Crystallography Protein–Protein Interaction Inhibitors	
Beam Physics	42		20	Frotein-Frotein interaction ininoitors	22
Bioinformatics	58, 62	Intracellular Delivery	50	[0]	
Biointerface	12	Iron Catalyst	30	[Q]	28
Boolean Networks	60	DIZ1		Quantum Chemical Calculation	
Boolean Networks	60	[K]	4	Quantum Information Science	16
ICI		Kinetic Stabilization	4	Quantum Materials	16
[C] Cell Therapy	26	IT 1		Quantum Sensing	16 18
Charge Generation and Transpor		[L] Laser Nano-ablation Physics	44	Quantum Transport	10
	1 40 26			(D)	
Chemical Biology		Laser Plasma Radiations (electron	18, 10118, 44	[R]	<i>5</i> 1
Chemical Genetics	26	and THz)		Reaction Mechanism	54
Chemical Graphs	60	Living Radical Polymerization	12, 14	Regioselective Functionalization	8
Chemical Library	26	Low-coordinate Phosphorus Ligan		Rheology	38
Computational Genomics	62	Low-coordinated Species	4	RNA	24
π-Conjugated Polymer	54	0.0		101	
π-Conjugated Systems	6	[M]	(2)	[S]	
COP9 Signalosome	24	Machine Learning	62	Scale-free Networks	60
Crystal	48	Marine Chemistry	30	Selective Organic Reaction	50
Curved π-Conjugated Molecules	14	Membrane Curvature	20	Self Assembly	36
IDI		Membrane Vesicle	34	Semiconductor Nanoparticles	56
[D]	(2)	Membrane-Permeable Peptides	20	Signal Transduction	24
Data Mining	62	Metal Ion Recognition	30	Single Electronics	10
Diabetes Diagnosis	22	Metalated Peptide Catalyst	50	Single Photon Spectroscopy	56
Diamond	16	Metallic Nanoparticle Catalyst	50	Small Molecules	26
Dielectric Spectroscopy	38	Molecular Electronics	40		5, 38
DNA/RNA Binding Protein	20	Molecular Evolution	58	Solar Cells	56
Dynamic Chirality	8	Molecular Recognition	8	Solid State Chemistry	52
OF 1		Morphogenesis	24	Solid-State NMR	28
[E]	4.6	DATI		Solution-Phase XAS Analysis	50
EELS	46	[N]	1.0	Spintronics	18
Electrical Conductivity and	40	Nanocomposite Magnet	10	Stable Isotopes	30
Photoconductivity	40	Nano-fabrication	18	STEM	46
Electronic Structure	46	Neutron Optics	42	Steric Protection	4
Elemental Ratio	46	NV Center	16	Structural Biology	48
Endohedral Fullerenes	6	101		Structure and Function	48
Environmental Genomics	58 22	[0]	20	Surface and Interface Chemistry	32
Enzyme Inhibitors	52	Organic Light-Emitting Diodes	28 32	Surface Plasmon	46
Epitaxial Thin Film Growth	34	Organic Semiconductor		Systems Biology	62
Extremophiles	34	Organic Seniconductors	40	[TP]	
LIE!		Organic Solar Cells	6, 28	[T]	40
[F]	4.4	Organic Synthesis	14	Thin Films	40
Femtosecond Laser Processing	44	Organocatalysis	8	Trace Elements	30
Femtosecond Laser Spectroscopy	y 56 22	ini		Transition Metal Complex	54
Fructosyl Peptide Oxidase		[P]	20	Transition Metal Complexes	4
Functional Materials Functional Transition Metal Oxid	6 des 52	Peptide Design	20	DED	
Functional Transition Metal Oxid	ues 32	Perfluoroalkyl Compounds Perovskite Structured Oxides	32 52	[U] Ultrafast Electron Diffraction (UED)	4.4
ICI		Perovskite-Based Solar Cells		Unusual Amino Acid	44
[G] GenomeNet	58	Perovskites Perovskites	6 56	Oliusuai Allillio ACIU	8
γ-Glutamyl Transpeptidase	38 22	Phase Rotation	36 42	IX 71	
Grammar-based Compression	60	Phase Rotation Phospholipid Acyltransferase	34	[V]	58
Grammar-based Compression	00	Phospholipid Signaling	24	Virology	20
nn		Photocatalysts	10	[V]	
[H] Heteroatom	4	Plasmonics	10	[X] X-ray Crystallographic Analysis	48
Hierarchical Structure	36	pMAIRS	32	A-ray Crystanographic Analysis	40
High Pressure Synthesis	52	Polymer Brush	12		
TITELL LICESCULC DALIMICSIS	34	1 OTYTICE DEUSII	14		

Institute for hemical esearch

Kyoto University